



African Journal of Biological Sciences

Journal homepage: <http://www.afjbs.com>



Research Paper

Open Access

Multi-Location Adaptability Trial for Yield Potential Assessment of Exotic Potato Varieties in Punjab

^{1*}Afshan Afzal, ²Mohammad Aslam, ³Muhammad Ramzan, ⁴Javed Ahmed Abro, ⁵Muhammad Nasir, ⁶Naeem-ud-din, ⁷Tariq Aziz, ⁸Abdul Ghani Langau

^{1*}Department of Botany, University of Education Lahore, Pakistan.

²PARC-Horticulture Research Institute, Khuzdar, Balochistan, Pakistan.

³WWF-Pakistan, Food and Markets Program, Khuzdar Balochistan, Pakistan.

⁴PARC-Balochistan Agricultural Research and Development Centre, Quetta, Pakistan.

⁵Horticulturist at Khaity Technologies PVT Olive Project PSDP, Pakistan.

⁶PARC-Horticulture Research Institute, Khuzdar, Balochistan, Pakistan.

⁷Assistant Development Officer at Pakistan Tobacco Board, Ministry of National Food Security And Research, Islamabad, Pakistan.

⁸Directorate of Agriculture Research Vegetable Seed ARI saryab Quetta, Balochistan, Pakistan.

Corresponding Author: ^{1*} Afshan Afzal

Department of Botany, University of Education Lahore, Pakistan.

Email: afshanafzal43@gmail.com

Article Info

Volume 6, Issue 12, September 2024

Received: 23 June 2024

Accepted: 19 August 2024

Published: 07 September 2024

[doi: 10.33472/AFJBS.6.12.2024.6504-6510](https://doi.org/10.33472/AFJBS.6.12.2024.6504-6510)**ABSTRACT:**

An adaptive trial was conducted by Department of Botany Research Punjab to evaluate the performance of Three exotic potato varieties under the climatic conditions of Punjab. Tubers of varieties Lady Rosita, Sante and Asterix were sown at Seed centers. The experiment laid out using Replicated Complete Block Design (RCBD) with 3 replications. Standard agronomic and plant protection measures were applied. Data of the parameters such as plant height, number of stems per plant, number of tubers per plant, weight of tubers per plant and yield per hectare were observed. Analysis of variance showed that all the varieties were significantly different in all the parameters. LSD test was applied to compare means of all the parameters. Plant height showed significant difference among the varieties with Asterix showing maximum plant height of 49.3 cm. Variety Sante showed least plant height having mean of 20.2cm. Variations were observed among the varieties for number of stems per plant. Sante showed maximum number of stems having mean of 3.8. Least number of Lady Rosita stems were observed in variety had mean value of 2.5. Variations were also observed among the varieties for number of tuber per plant. Sante produced maximum number of tubers having mean value of 12.4. Variety Asterix produced least number of tubers per plant having mean value of 7.3. Maximum tuber weight per plant was produced by Asterix having a mean value of 932.3. Lady Rosita produced least weight of tubers per plants with a mean value of 741.2. Variety Sante was best yielder producing 18.5 tons/ha of seed potato while varieties Lady Rosita showed minimum yield of 13.4 tons.

© 2024 Afshan Afzal, This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made

1. Introduction

Potato (*Solanum tuberosum* L.) belongs to family Solanaceae and is known as king of the vegetables as being part of every meal. It has been extensively cultivated all over the world and is ranked as 3rd after wheat and rice, whereas it is 4th valuable food crop on basis its domestic consumption after rice, maize and wheat (Devaux et al., 2014). Potato is a resourceful food crop being grown from sea level to high mountainous regions and is now under cultivation in almost 100 countries that can be grown in various environments and is currently cultivated globally in over 100 countries (United States Potato Board, 2015). It contains 79% water content in it along with carbohydrates (18%), protein (2%), and vitamins (1%) and many minerals and trace elements that make it a nutritious and energy rich food (Ahmad et al., 2011; Low et al. 2007). It has no cholesterol contents in it, additionally it possesses some antioxidants that can be useful to cure cardiac and cancer disease in human (Isobel Hoskins 2005, Al-Saikhan et al., 1995). Medically it has given keen importance to useful to manage hypertension and high blood pressure due presence of an organic content kukoamine (Anonymous 2017; Akande et al., 2000). It is also named as poor man's food as it is included in every meal because

it provides more calories of per hectare than any food crop grown all over the globe (Amdie et al., 2017).

In Pakistan three crops are being cultivated as in summer (70-75%), autumn (20%), and sprig crop (710%). Summer crop is cultivated from March to April in the mountainous areas of Gilgit Baltistan and Kashmir. Autumn crop is being sown from the month of September to October in plains of Punjab, Sind and Baluchistan whereas, Spring crop season strats from January and ends in late Fabruarty in the lower hilly areas of Balochistan and KP (GOP, 2016-2017). Due to more returns from potato to the farmers during the last decade in Pakistan it has been considered as high value potato cash crop. Potato has been grown on 0.196 m ha⁻¹ with annual production of 4.57 million tonnes having average yield of 19.32 t ha⁻¹ (Pakistan Economic Survey, 20182019). However, the potato production in Pakistan is very low due to several reasons i.e. high incidence of pests, insects, diseases, water scarcity etc along with cultivation of low yielding varieties and use of table potato as seed. Thus the aim to evaluate high yielding and well adapting exotic potato varieties is an integral component of research either in government of private sector. Being an important commodity from the consumption and trade perspectives, increase in potato productivity in future requires promising attention from the concerned stakeholders. Provision of high yielding varieties of the crop imported from other countries to farmers may significantly elevate its yield outputs and production.(Adul Majeed and Zahir Muhammad.,2018).

In the Punjab region, the Department of Agriculture Research has established advanced Potato Tissue Culture Laboratories in areas Sahiwal, Faisalabad & Lahore. These facilities, which include greenhouses and screen houses, are focused on producing pre-basic seed potatoes of both exotic and national varieties. The goal is to produce and test genotypes that are well-suited to the local environmental conditions.

2. Material and Methods

Tubers of three varieties i.e Lady Rosita, Sante, Asterix were cultivated. Experiment was laid using Randomized Complete Block Design having 3 replications. In each replication 33 tuber of each variety were sown Plants on ridges having plant to plant distance of 15 cm and row to row distance of 75cm. Standard agronomic and plant protection measures were applied. Crop was harvested of each variety were randomly selected and data of parameters i.e. plant height (cm), number of stems per plant, number of tubers per plant and weight of tubers per plant (gms) were recorded. Total production (kg) of all plants of each variety in each replication were recorded and yield/hectare were calculated from it. Data was statistically analyzed using Statistix version 8.1 software.

3. Results and Discussions

The results pertaining to the adaptability of these exotic potato varieties under climatic conditions and their analysis of variance depicted significant differences in the selected parameters that are discussed under following headings.

Plant height

The plant height of the potato varieties was recorded when they got maximum growth and development in the field. Highly significant results were studied among the varieties for this parameter. Pameela variety showed maximum plant height Lady Rosita (42.3 cm) & minimum Sante (20.2cm) (Table 1). On the basis LSD values six homogenous groups (a, ab, bc, c, cd) were formed that showed significant variation for plant height among the three potato varieties. The study of Khalid et, al. (2019) is in congruence to our findings that the potato varieties

showed variation for the morphological parameters as it is plant genetic attributes that cause variation. Our results are in accordance with Eaton et al. (2017) who reported difference in plant height of different potatoes genotypes and might be due to plant genetic makeup and environmental effects. The findings of Luthra et al. (2005) and Schittenhelm et al. (2006) also revealed the differences in potato and they concluded that all these differences for plant height are due to plant genotype, variety, nutrients availability and the agroecological factors under area of study.

Number of stems plant⁻¹

The number of main stems plant⁻¹ was also recorded for the potato varieties under investigation. There was significant variation among the tested potato varieties for stems plant⁻¹. Sante variety showed maximum number of shoots/ plant with a mean value 3.8. Minimum number of shoots/ plants was observed in variety Lady Rosita having mean value of 2.5 shoots/plant. The ANOVA also depicted the significant differences among potato varieties for number of stems plant⁻¹ (Table 1). Khalid et, al. (2019) revealed similar results that the potato variety Menato produced maximum (16.2 stems m⁻²). The findings of Eaton et al, (2017) are in strong conformity of our results that the potato varieties showed significant differences in number of stems plant⁻¹ Our results are similar to those reported by Hoque (1990) reported 5.17 main stems in potato variety Diamant, whereas Bashar (1978) th reported 5.2 main stems plant⁻¹ in potato variety Cardinal. Difference in number of stems plant⁻¹ was due to the genetic variability of varieties. Similar results were reported by many researchers who found that different potato cultivars exhibit great variation for different characteristics like number of stems, number of leaves, plant height and number of tubers per plot (Hanan & Lodhi, 1979; Randhawa et al., 1980).

Number of tubers plant⁻¹

The number of tubers plant⁻¹ is amongst most economical parameter that directly measures the yield potential of tuber crops. In the current study this parameter was also recorded for the potato varieties under inquire. The varieties displayed variation for this parameter. The variety Sante produced maximum (12.4) tubers plant⁻¹. The least number of tubers plant⁻¹ were recorded in Asterix that produced 7.3 tubers plant⁻¹ Table 1. The statistical analysis also depicted significant differences amongst the candidate potato varieties. Our conclusions are in line with the findings of Khan et, al (2019) that envisaged great variation among the tested potato genotypes and potato lines. Our results are also in agreement with the assessment of Eaton et al. (2017) that exposed the tuber production in potato crop is a dependant on the genetic makeup of the genotypes, the method of planting used and the environmental conditions (availability of water and soil nutrients. The stolon and tuber formation in potato is critically affected by genetic makeup of varieties and agroecological and environmental features (Subarta and Upadhyya, 1997). Any how the findings of Khalid et, al. (2019) are in contrast to our findings that reported the potato variety Melanto produced maximum (50.76) tubers m² in comparison with Desiree (38.21) tubers m².

Tubers weight plant⁻¹

Tuber weight plant⁻¹ also contributes considerably to the yield attribute of potato crop. The variability in tuber production is an inherited character that is genotype dependant. Currently to determine yield potential, the tuber weight plant⁻¹ was investigated. The potato varieties responded significantly different for this economically important parameter. In the current study, the potato variety Asterix produced more bulky tubers plant⁻¹ with maximum (932.3 g) tubers weight plant⁻¹. Least weight of tubers plant⁻¹ having 741.2 g was observed in variety Lady Rosita Table 1. The ANOVA also displayed significant variation among the potato varieties under study. Our results are supported by the findings of Panday et, al (2004); Panday

et al; (2008) that reported variation in tuber yield among the assessed potato varieties and lines. The findings of Patel et al, (2008) also concluded that the bulky and large size tuber production in potato may be due to good seed germination and better plant growth and development under a favorable agroecological zone. **Yield hectare**

Yield determination is the main parameter that assesses the potential of seed potato production in the varieties under study. In this investigation the potato variety Sante was recorded as the high yielding variety among the compared varieties with a production of 18.5 tons ha⁻¹. On the other hand, the potato variety Lady Rosita emerged as least productive with 13.4 tons ha⁻¹ (Table 1). The findings of Khalid et al, (2019) also depicted that the yield potential of potato crop is dependent on variety, its genetic features and influence of various biotic and abiotic factors in the area of study. The findings of Patel et al, (2008); Patel et al, (2004) are also in strong conformity of our study that potato varieties different from each other on basis of yield that is directly related with maximum seed germination, better plant growth and development, successful tuber formation having good size and shape under available field conditions. Variation in yield of seed potato had been reported by Hossain et al 2003 and Anonymous (1990). In contrast the findings of Shafayet et al who found that a highest yield was observed in Granola (27.82 t/ha) followed by Asterix (26.83 t/ha) and Provento (26.33 t/ha). Our findings are different from those reported by Shafayet et al . (2005) who reported significant variation in the yield potential of various seed potato varieties. Shehroz et al (2018) also find that several exotic varieties of potato with same plant to plant and row to row distance could be adopted for local cultivation to enhance the potato production in the country.

Sante variety shown high yield per hectare as desired, the development and accessibility of new varieties of potato which are suitable for the processing is quite critical for sustaining the rise in processing industries of Pakistan. Moreover, the potato varieties with more dry matter and less reducing sugars are required by potato processing industry to ensure high quality processed products. Enhancement in the production of potato is just due to the application of modern technologies and utilization of new seed varieties. White potatoes and the red potatoes are the major contributors in potato production by volume in Pakistan (Sana, 2019)

S.No	Variety	Plant Height (cm)	No. of Stems/Plant	No. of Tubers/Plant	Weight of Tubers/Plant (gm)	Yield/ha (Tons)
1	Lady Rosita	42.3c	2.5bc	11.2bc	741.2bc	13.4c
2	Sante	20.2d	3.8b	12.4ab	858.3abc	18.5b
3	Asterix	49.3ab	3.6bc	7.3bc	932.3abc	18.0b

Table no 1: Analysis of varieties of Potatos

4. Conclusions

On basis of field performance Sante emerged as best yielding variety for seed potato production at Punjab followed by Asterix whereas; and Lady Rosita responded as least productive in this area.

5. References

1. (½): 84–7 Abbas G, Hafiz IA, Abbasi NA & Hussain AZHAR (2012). Determination of processing and nutritional quality attributes of potato genotypes in Pakistan. Pak J Bot 44: 201-208.

2. Abbasi et al. 2004. Evaluation of Exotic Potato Varieties in Ecological Conditions of Islamabad during Autumn
3. Abdul Majeed and Zahir Muhammad 2018. Potato Production in Pakistan: Challenges and Prospective Management Strategies –a Review. Pak. J. Bot., 50(5): 2077-20 84
4. Ahmad, K.U. (1980) Exotic Potato Varieties for Bangladesh. Proc eedings of the 3rd Workshop of Potato Research Workers , PRC, BARI, Joydebpur, Dhaka, 1-12.
5. Anonymous. 1992. Variety Development from Exotic Sources. Annual Report, TCRC, BARI, Joydebpur, Gazipur, Bangladesh, 21-52.
6. FAO. 2008. International year of the potato. www.Potato2008.org.
7. FAO. 2010. Strengthening potato value chains:Technical and policy options for developing courtiers. Food and Agriculture Organization of the United Nations, Rome.
8. FAOSTAT (2014). www.potatopro.com/world/potatostatistics
9. FAOSTAT. 2012. Faostat.fao.org.
10. GOP (2012). Agriculture Statistics of Pakistan, Ministry of food, Agriculture and Livestock, Govt. of Pakistan, Islamabad. pp 82.
11. Hanan, A. and F. Lodhi, 1979. Relative performance of German potato cultivars in Peshawar. In Proceedings of 26th/27th Pakistan Science Conference, Lahore 1979. Part III Abst. Of the Papers. Lahore, Pakistan. Association for the Advancement of Science
12. Hossain, M.M., Rashid, M.H., Sattar, M.A. and Akhter, M.I. 2003. Preliminary Yield Trial (PYT) of Dutch Varieties in Their 1st Generation. Annual Research Reports on Tuber Crops Improvement, TCRC, BARI, Joydebpur, Gazipur, 9-10
13. IUCN, NAs. 2002. Strategy for Sustainable Development—Background Paper on Agriculture and Food Security. (Second draft), Pakistan.
14. Khalid, S., S. Shaheen, A. Ishfaq and R. Siddique. 2019. Performance study of exotic variety of potato (*Solanum tuberosum* L.) for processing industry under the agroclimatic condition of Lahore-Pakistan. Pure Appl. Biol., 8(4): 2262-2268.
15. Luthra SKJ, Gopal SK, Pandey Singh BP (2005). Genetic parameters and characters associated in *tuberosum* potatoes. Potato J. 32:234-239.
16. Pandey SK, Singh SV, Kumar D, Kumar P & Manivel P (2004). Sustaining potato chipping industry from western and central Uttar Pradesh: Adoption of suitable varieties. Potato J. 31: (3-4).
17. Pandey SK, Singh SV, Kumar D, Manivel P, Marwaha RS, Kumar P & Gupta VK (2008). Kufri Himsona: A chipping cultivar for hill regions of India. Potato J 35(1-2): 1-8.
18. Potato-Cultivation-in-Pakistan. www.valleyirrigationpakistan.com/wp-content.
19. Qasim, M., S. Khalid, A. Naz, M.Z. Khan, S.A. Khan. 2013. Effects of different planting systems on yield of potato crop in Kaghan valley. A mountainous region of Pakistan. Agricultural Sciences. 4(4).
20. Randhawa, K.S., B.S. Joyapa and K.S. Sandhu, 1980. A note on the performance of some new genotype (*Solanum tuberosum* L.) for qualitative and quantitative factors under Punjab condition. Haryana J. Hort. Sci.,
21. Robert, H. 2001. Global distribution of the potato crop. American Journal of Potato Research. 78 (6): 403–12.
22. Salunkhe DK & Kadam SS (1998). Handbook of vegetable science and technology: production, compostion, storage, and processing. CRC press.
23. Sana Khalid., Shabnum Shaheen., Ammara Ishfaq and Riffat Siddique (2019). Performance study of exotic variety of potato (*Solanum tuberosum* L.) for processing industry under the agroclimatic condition of LahorePakistan. Pure Appl. Biol., 8(4): 2262-2268, December, 2019 <http://dx.doi.org/10.19045/bspab.2019.80172>

24. Sayem, M. A., Taslima, N., Sidhu, G. S., Chowdhury, F., Sumi, S. M., Anwar, A. S., & Rowshon, M. AI-Driven Diagnostic Tools: A Survey of Adoption and Outcomes in Global Healthcare Practices.
25. Schittenhelm S, Sourell H, Löpmeier FJ (2006). Drought resistance of potato cultivars with contrasting canopyarchitecture. *Eur. J. Agron.* 24:193-202
26. Season/ *Int. J. Agri. Biol.*, Vol. 6, No. 3, *International Journal of Agriculture & Biology* 1560– 8530/2004/06–3–479–482
27. Shabbir, A., Anwar, A. S., Taslima, N., Sayem, M. A., Sikder, A. R., & Sidhu, G. S. Analyzing Enterprise Data Protection and Safety Risks in Cloud Computing Using Ensemble Learning.
28. Shafayet, H., Golam, R., Khaleque, A.M, Moynul, H. and Mannan, A.K. (2015) Yield Potential of Twelve Potato (*Solanum tuberosum*) Varieties Grown from Different Generations of Seed. *The Agriculturists* , 13,120132.
29. Shehroz M. M. Hussain , I. Ramzan and M. A. Ali.(2018) Evaluation of Exotic Potato Germplasm for High
30. Sidhu, G. S., Sayem, M. A., Taslima, N., Anwar, A. S., Chowdhury, F., & Rowshon, M. (2024). AI AND WORKFORCE DEVELOPMENT: A COMPARATIVE ANALYSIS OF SKILL GAPS AND TRAINING NEEDS IN EMERGING ECONOMIES. *International journal of business and management sciences*, 4(08), 12-28.
31. Subarta M., Upadhyya, M.O. 1997. Potato production in western Bengal. *Env. Ecol.* 15: 646–9.
32. Touria E. Eaton, Abul Kalam Azad, Humayun Kabir, Abu Bakkar Siddiq. 2017. Evaluation of Six Modern Varieties of Potatoes for Yield, Plant Growth Parameters and Resistance to Insects and Diseases. *Agricultural Sciences.* (8) : 1315-1326.
33. Yield and Disease Resistance under local conditions. *The Journal of Animal & Plant Sciences*, 28(5): 2018, Page: 1413-1417 ISSN: 1018-7081