



ALLELOPATHIC EFFECT OF CYANOBACTERIAL STRAINS ON PHYTOPATHOGENIC FUNGI

RICHHA MISHRA¹ and ASIT KUMAR²

¹Department of Microbiology, A.P. S.G. M. N. S. Govt. P.G, College, KABIRDHAM (C.G.) INDIA

²Department of Zoology, Govt. R. V. R. S. Kanya Mahavidyalaya Kawardha , (C.G.) INDIA

*Corresponding Author: E. mail.richa.micro10@gmail.com

Volume 6, Issue 15, Sep 2024

Received: 15 July 2024

Accepted: 25 Aug 2024

Published: 05 Sep 2024

[doi: 10.48047/AFJBS.6.15.2024.5993-6005](https://doi.org/10.48047/AFJBS.6.15.2024.5993-6005)

Abstract:

Allelochemicals are subsets of secondary metabolites that no longer required for metabolism of the allelopathic organism and their poor allelopathic results are vital part of organism defense against antagonists. Allelopathic interactions involving Cyanobacterial flora are being explored for their pharmaceutical and environmental importance. Cyanobacterial allelopathy may be appeared as one of the massive elements influencing their dominance in numerous habitats and as particular producers of a selection of allelochemicals that can be applied as green biocontrol marketers. In present work detrimental (dreadful allelopathy) effects of locally isolated Cyanobacterial strains had been evaluated in opposition to plant pathogenic three fungal isolates (*Fusarium oxysporum*, *Sclerotium rolfsii* and *Rhizoctonia solani*). It turned into discovered that the crude extracts of 4 Cyanobacterial isolates (*Microcystis aeruginosa*, *Oscillatoria boryana*, *Anabaena sphaerica* and *Nostoc calcicola*), had been able to diminishing the boom and similarly development of phyto-pathogenic fungal isolates. While *M. Aeruginosa* showed greater allelopathic interest compared to different cyanobacterial strains. Methanol crude extracts have been extra efficient. Crude extract (100 % in Methanol) of *Microcystis aeruginosa* changed into determined extra efficient towards neighborhood isolates *Fusarium oxysporum* (% MI – 88.42 ± 0.24) and *Sclerotium rolfsii* (83.42), in comparison to *Rhizoctonia solani* (78.33). Allelopathic potentiality of cyanobacteria have need to be further investigated which could provide promising answers in bio-control in opposition to pathogenic Fungi.

Keywords: Allelopathy, Cyanobacteria, Phyto-pathogens, Crude extracts, Biocontrol agents.

Introduction:

An expanding international population and urgency of removing hunger and malnutrition call for decided rules and powerful actions to make certain sustainable growth in agricultural productiveness. India is a biggest Country in South Asia that includes 70 percent of the total nearby populace. After green Revolution big progress with cutting-edge irrigation and fertilizer software has been made. The concern of pesticide use with respect to human fitness and surroundings has added increasing hobby in options by using fending off terrible results on the environment. Secondary metabolites from cyanobacteria are related to poisonous, hormonal, antineoplastic and antimicrobial

effects (Shweta et al., 2011). Currently algae are one of the leader organic dealers which have been studied for the manipulate of plant pathogenic fungi, especially soil-borne disorder (Hewedy, 2000). Cyanobacteria and eukaryotic algal produce biologically active compounds which have antifungal and antibacterial pastime (Kulik, 1995) against plant pathogens. Allelopathy is a unique adaptation to gain a aggressive gain over different organism inhabiting the equal microbial network, Allelopathic impact of Cyanobacteria on pathogenic fungi was also studied (Tiwari and Kaur, 2014). Cyanobacteria are known to supply intracellular and extracellular metabolites with diverse biological activities consisting of antialgal, antibacterial, antifungal and antiviral pastime (Mohamed El-anwar H. Osman, 2011; Shrivastava et al., 2012; Shrivastava et al., 2014., Shrivastava et al., 2017). The antimicrobial substances may additionally target diverse sorts of microorganisms. Secondary metabolites influence different organisms inside the location and are idea to be of phylogenetically crucial. Kulik, (1995), stated some of reasons, cyanobacteria and algae are suitable candidates for exploitation as biocontrol agents of plant pathogenic fungi. The ability for the use of Cyanobacteria is the biocontrol of plant pathogenic bacteria and fungi. Tiwari & Sharma (2013), has found the antifungal interest of *Anabaena variabilis* in opposition to plant pathogens. Screening of cyanobacteria for antibiotics has opened a new horizon for discovering new capsules. Exploring allelopathic efficacy of cyanobacteria to govern plant pathogenic fungi can prove to be a fantastic option as they're easy to develop with minimum vitamins, price effective, no aspect results and surroundings friendly. This work describes the allelopathic impact of cyanobacterial strains towards phytopathogenic microorganisms.

MATERIALS AND METHODS:

Isolation and identification of microbes:

For the choice of the experimental materials distinct species of fungi and cyanobacteria have been remoted applying right media and methods. Identification of remoted microbes became finished with the aid of microscopic statement and biochemical checks.

Cyanobacteria:

Soil samples had been amassed from diverse agro-fields of Bilaspur division. The samples had been introduced to laboratory in plastic vials /zipped polythene bag and washed with distilled water to save potential contaminants. Isolation of cyanobacterial strains became carried out in particular nutrient media (Allen & Arnon, BG-one hundred ten and Chu-10) and changed into saved as suspension way of life in boom medium below prescribed situation (Shrivastava, 2000 and Shrivastava et al., 2005). Isolated strains have been recognized by way of microscopic observation with the help of Key standardized by means of Desikachary (1959) and Anand (1989).

Fungal lines:

The fungal strains have been remoted from agronomic soil samples. They were grown in the laboratory in suitable media, for fungal evaluation media used which includes Potato dextrose agar media (PDA) and Sabouraud's agar media and were identified on the basis of their morphological traits. Taxonomic guides (Alexopoulos, C. J., et al., 1996) have been used to become aware of the remoted fungal culture.

Extraction:

The bloom samples were collected from various zones of Bilaspur division. The bloom samples had been accumulated the use of plankton net of 20µm mesh size. Samples were saved in sterile zipped polythene luggage. Cyanobacterial cells had been focused by way of the usage of centrifugation. A portion of the concentrated samples turned into filtered thru a 0.45 µm glass fiber clear out (What's men-41) and air dried in an oven at 60 °C.

Dried cellular mass-2g/25ml (w/v) of Cyanobacteria had been extracted with 75% methanol (HPLC grade), for one hour then centrifuged at 5000 rpm for 7 min. The supernatant changed into separated in clean glass vials and filtered with 0.45 µm pore length. Specific Dilutions have been organized for toxicity check in-vitro.

Toxicity assessment:

For checking out the allelopathic activities of crude extract of cyanobacteria against diagnosed fungal isolates with the help of special methods and Carbendazim and Mancozeb (as fungicides) had been used as control. The following techniques had been hired to confer the toxicity of crude extract of various Cyanobacterial isolates on identified micro organism & fungi.

Percent mycelial inhibition:

Petri plates with medium were aseptically inoculated on the centre with mycelial disc of 5mm diameter taken from ninety six hrs vintage way of life. Plates were then incubated at 26 ± 1°C in BOD incubator and mycelial growth pattern became found. Radial diameter turned into recorded twice perpendicularly after 96 hrs incubation using a transparent millimeter ruler. Percent inhibition of mycelia boom changed into calculated.

RESULTS AND DISCUSSION:

Through exam of samples overall sixteen species of cyanobacteria have been remoted, out of which 9 heterocystous shape and 7 non-heterocystous bureaucracy have been recognized as nitrogen fixing and non nitrogen fixing traces respectively (table -1, PLATE -1; Fig. – I to XVI).

Table - 1: Morphologically distinct Cyanobacteria isolated at some point of the survey.

TABLE - 1: Morphologically dissimilar Cyanobacteria isolated during the survey.

	Non Nitrogen fixing	Nitrogen fixing
Unicellular	1. <i>Aphanocapsa elachista</i> 2. <i>Aphanothece saxicola</i>	NIL
Colonial	3. <i>Microcystis aeruginosa</i>	NIL

Filamentous	4. <i>Lyngbya birgei</i>	8. <i>Anabaena oryzae</i>
	5. <i>Lyngbya shackletoni</i>	9. <i>Anabaena sphaerica</i>
	6. <i>Oscillatoria boryana</i>	10. <i>Anabaena unispora</i>
	7. <i>Oscillatoria cortiana</i>	11. <i>Aulosira prolifica</i>
		12. <i>Calothrix elenkinii</i>
		13. <i>Cylindrospermum indicum</i>
		14. <i>Gloeotrichia echinulata</i>
		15. <i>Nostoc calcicola</i>
		16. <i>Nostoc ellipsosporum</i>

Simultaneously total 6 fungal species (*Aspergillus niger*, *Penicillium notatum*, *Mucor sp.*, *Sclerotium rolfii*, *Fusarium oxysporum* and *Rhizoctonia solani*) were isolated and identified on the basis of their morphological features (TABLE -2, PLATE -2; Fig. – I to VI).

TABLE-2: Characteristic features of fungal isolates.

NAME OF FUNGI	TAXONOMIC STATUS	COLOUR	MICROSCOPIC OBSERVATION
<i>Aspergillus niger</i>	Ascomycetes	dark to pale yellow in early stage while became black at maturity	Jet black conidia, reverse usually grey, spherical conidia, rough with maturity.
<i>Penicillium notatum</i>	Ascomycetes	velvety or woolly colonies initially white and become green	Conidiophore green in colour, repeatedly branched, brushy head.
<i>Mucor sp.</i>	Zygomycetes	filamentous colonies, fluffy appearance, white in colour	Globular sporangia round black in colour that was supported by elevated column like columella. Non-septate or partially septate hyphae .
<i>Sclerotium rolfii</i>	Basidiomycetes	It was characterized as felty white appearance	Silky white hyphae tend to aggregate into rhizomorphic cord as well as characteristic sclerotium – fan shape was found.
<i>Fusarium oxysporum</i>	Ascomycetes	It was characterized as white to pink colony	Moist appearance, red with cottony and orange brown mycelium, with light brown exudates hyphen septet, small conidia in chain, sickle shaped long conidia were also observed.
<i>Rhizoctonia solani</i>	Basidiomycetes	Buff to black colony	The mycelium is generally hyaline, irregular in shape. Hyphae produce branches at right and acute angles to the main hyphae. specialized hyphae composed of moniloid cells, that fuse together to produce hard structure called sclerotia.

The efficacy of Cyanobacterial allelopathy changed into screened in opposition to the increase of *E. Coli* (ATCC-25922). The effect of crude extracts of all Cyanobacterial isolates was tested on growth pattern of test strain via measurement of optical density (600nm). And located information had been referred to in table–four and PLATE -four.




PLATE- 3

On the idea of toxic potentiality of crude extracts non- heterocystous (*Microcystis aeruginosa* & *Oscillatoria boryana*) and two heterocystous (*Nostoc calcicola* & *Anabaena sphaerica*) cyanobacterial isolates had been selected for similarly test in view of the evaluation of the allelopathic effect of cyanobacterial isolates on phytopathogenic Fungal isolates. To assess the Allelopathic impact of cyanobacterial became evaluated through percent mycelia inhibition and involved graph indicated that the extract of *Microcystis aeruginosa* have great potential as fungicides Carbendazim and Mancozeb compound against pathogenic fungi which turned into examined (desk – five, PLATE -five; Fig. I - IV.). Poisonous potentiality of *Nostoc calcicola* was observed lesser than *Oscillatoria boryana* however extra than different cyanobacterial isolates, whereas minimum doubtlessly of growth inhibition turned into discovered in case of *Anabaena sphaerica*.

TABLE – 5 Allelopathic effect of crude extracts of Cyanobacterial isolate on Fungal isolates (*Fusarium oxysporum*, *Sclerotium rolfii* and *Rhizoctonia solani*).

Cyanobacterial Isolates	Crude extracts	Conc. of crude extract	Percent of Mycelial Inhibition (Mean value ±SD)		
			<i>Fusarium oxysporum</i>	<i>Sclerotium rolfii</i>	<i>Rhizoctonia solani</i>
<i>Anabaena sphaerica</i>	Ethanollic extract	25	21.42 ±0.25	14.35 ±0.18	12.23 ±0.22
		50	30.61 ±0.32	21.25 ±0.43	14.27 ±0.52
		75	42.37 ±0.27	28.58 ±0.26	25.62 ±0.35
		100	60.25 ±0.22	56.16 ±0.15	53.24 ±0.28
	Methanollic extract	25	23.51 ±0.24	17.48 ±0.35	13.26 ±0.26
		50	33.52 ±0.22	23.28 ±0.43	16.17 ±0.52
		75	45.56 ±0.26	30.88 ±0.36	27.65 ±0.41
		100	62.25 ±0.33	58.35 ±0.38	55.41 ±0.29
<i>Microcystis aeruginosa</i>	Ethanollic extract	25	53.37 ±0.32	45.25 ±0.24	39.67 ±0.25
		50	71.56 ±0.16	63.21 ±0.23	51.45 ±0.21
		75	76.24 ±0.34	69.51 ±0.28	65.66 ±0.19
		100	84.52 ±0.35	81.34 ±0.42	75.35 ±0.28
	Methanollic extract	25	56.23 ±0.42	49.52 ±0.34	40.34 ±0.27
		50	71.24 ±0.36	66.23 ±0.45	53.45 ±0.28
		75	79.43 ±0.27	73.34 ±0.35	67.28 ±0.32
		100	88.46 ±0.24	83.42 ±0.28	78.33 ±0.45
<i>Nostoc calcicola</i>	Ethanollic extract	25	43.65 ±0.35	40.18 ±0.43	33.21 ±0.56
		50	58.25 ±0.32	47.32 ±0.24	41.62 ±0.29
		75	65.28 ±0.32	60.35 ±0.46	56.75 ±0.34
		100	81.52 ±0.25	70.81 ±0.21	63.33 ±0.33
	Methanollic extract	25	40.25 ±0.17	38.4 ±0.25	32.56 ±0.21
		50	56.18 ±0.42	44.67 ±0.16	39.28 ±0.32
		75	63.3 ±0.32	57.23 ±0.18	54.28 ±0.15
		100	72.26 ±0.26	66.53 ±0.34	61.36 ±0.42
<i>Oscillatoria boryana</i>	Ethanollic extract	25	43.65 ±0.23	41.56 ±0.28	35.57 ±0.23
		50	62.21 ±0.35	48.26 ±0.27	43.55 ±0.26
		75	72.24 ±0.44	64.51 ±0.35	62.66 ±0.45
		100	80.32 ±0.35	78.24 ±0.42	71.44 ±0.28
	Methanollic extract	25	48.35 ±0.15	42.51 ±0.35	36.23 ±0.22
		50	64.25 ±0.12	52.56 ±0.17	43.45 ±0.25
		75	76.27 ±0.27	68.33 ±0.35	59.23 ±0.32
		100	83.34 ±0.34	79.36 ±0.28	73.31 ±0.45
Solvent control			0	0	0
Fungicides					
Carbendazim		100 ppm.	94.70 ±0.13	90.64 ±0.16	88.23 ±0.17
Mancozeb		100 ppm.	89.76 ±0.21	87.70 ±0.24	85.29 ±0.36







PLATE- 5: Fig I - IV

Discussion:

The consequences obtained all through present course of research, concerning Cyanobacterial toxicity, revealed that the boom of fungi. Fungal isolates have been inhibited once they had been subjected to crude extracts (Ethanol or Methanol) on solid nutrient media. Boom inhibition potency of crude extracts of cyanobacterial isolates have additionally been as compared with wellknown fungicides towards fungal boom respectively. The end result of toxic potentiality as acquired in opposition to fungal pathogens has additionally been supported the findings. Cyanobacteria produce biologically energetic compounds that have allelopathic and toxic hobby in opposition to plant pathogens (Bonjouklian, R. Et al. 1991; Kiviranta, J., et al. 2006). Kim and Kim, 2008, said inhibition of *F.Oxysporum f. Sp. Lycopersici* by way of extracts of *N. Commune FA-103*. Biological control of *Fusarium oxysporum f. Sp. Lycopersici* (FOL) inflicting wilt disorder of tomato was studied in vitro as well as beneath pot situations and became concluded that *N. Linkia*, can be utilised for the organic manage of wilt ailment of tomato which may additionally help to gain a higher yield and suitable fitness in agriculture (Hend, A. 2012). Mohamed et al. 2011, investigated the suppression effect of cyanobacterial species- *Nostoc endophytum* and *Nostoc muscurum* against, the causal agent of soyabean root rot *Rhizoctonia solani*.

The prevailing studies work correlates the findings of various people as suggested earlier concerning concerned research paintings. On this have a look at Cynobacteria confirmed tremendous allelopathic activities. This type of investigation produce a miles generalized view that cynobacteria are able to inhibiting phytopathogens. This work makes a speciality of the ability of Canobacteria to be used as allelopathic biocontrol agent further research on Cyanobacterial metabolites are vital for clinical employees to pick out the poisonous effect on phytopathogenic fungi which might also lead to the formula of considerable bio-energetic compounds of organic origin.

ACKNOWLEDGMENT:

Writer is thankful to the fundamental, Head, branch of Botany & Microbiology, government. E. Raghavendra Rao Postgraduate technological know-how college, Bilaspur (C.G.) for presenting research facilities to done the existing research and gratefully acknowledged to Ph. D. Manager Dr. D. K.. Shrivastava for his treasured steerage, large criticism and moral help.

REFERNCE:

Anand, N. (1989): Hand book of Blue Green Algae. Pub. Bisen Singh and Mahendrapal Singh. Dehradun.

Alexopoulos, C. J.; Mims, C. W. and Bleckwell, M. (1996): Introductory Mycology 4thEdt. John wiley and Sons, New York, USA.

Bonjouklian, R. Smitka, T.A., Doolin, L. E. Molloy, R. M., Deebono, M. Shaffer, S. A., Moore, E., Stewart, J.B. Patterson, G.M.L. (1991): Tjipanazoles, new antifungal agents agents from the blue-green algae *Tolypothrix tjipansensis*. *Tetrahedron*Vol. 47; Pp-7739-7750.

Desikachary, T.V. (1959, 1972):Cyanophyta. Pub. Indian Council of Agricultural Research New Delhi.

Hewedy, M. A., Rhhah, M.M.H. and Ismail, I. A. (2000): Pathological studies on soyabean damping off disease. *Egypt J. Applied Sci. Vol. 15; Pp-88-102.*

Hend A. A. and Perveen K. (2012): Biological control of *Fusarium* wilt of tomato by antagonist fungi and cyanobacteria. *African Journal of Biotechnology.* Vol.11; Pp-1100-1105.

Kulik, M. M.(1995): The potential for using cyanobacteria (blue-green algae) and algae in the biological control of plant pathogenic bacteria and fungi. *Eur. J. Plant Pathol.*Vol.101; Pp-585-599.

Kim, J. Kim, J.D.(2008): Inhibitory effect of algal extracts on mycelia growth of the tomato-wilt pathogen,*Fusarium oxysporum* f. sp. *lycopersici*. *Mycobiol.* Vol. 36; Pp- 242-248.

Kiviranta, J. Abdel-Hameed, A. Sivonen, K. Niemelä, S.I. Carlberg, G.(2006): Toxicity of cyanobacteria to mosquito larvae- screening of active compounds. *Environ. Toxicol. Water Qual. Vol. 8; Pp-63-71.*

Mohamed El-anwar H. Osman, Mostafa M. El-Sheekh, Metwally A. Metwally, Abd

El whab A. Ismail and Mona M. Ismail (2011): Antagonistic Activity

of Some Fungi and Cyanobacteria Species against *Rhizoctonia solani*. *International Journal of Plant Pathology Vol.2*; Pp- 101-114.

Mishra B. N. and Mishra , M. K. (1983): Introductory Practical Biostatistics, NayaPrakash, Calcutta-2006, BidhaSarni.

Shrivastava, D. K. (2000): Cyanobacteria from paddy fields of Durg district of Chhattisgarh State. *Phycos*. Vol. 39(1&2), p-125-128.

Shrivastava, A.K.; Pandey, F. K. and Shrivastava, D. K. (2005): Nitrogen fixing cyanobacteria flora and properties of paddy field soils of four districts of Chhattisgarh state. *Nat. J. Life Sc.* 2(Supp.), p-501-506.

Shrivastava, D. K. and Saluja, T. (2012): Antibacterial properties of exo-toxins released by Cyanobacteria- *Phormediumcaldicola* and *Oscillatoriapriniceps* isolated from Bilaspur (C. G.). *National Journal of life Science*. Vol. 09, Pp. – 229-231.

Shrivastava, D. K.; Mishra, R. and Chandra T. P. (2014): Evaluation of Cyanobacterial toxicity against corresponding field soil Bacteria. *International Journal of Pharmacy & Life Sciences* Vol- 5(7) Pp- 3694- 3700.

Shrivastava, D. K.; Mishra, R. and Chandra T. P. (2017): Allelopathic effect of cyanobacterial strains on phytopathogenic Bacteria *Research journal of pharmaceutical biological and chemical sciences* Vol-8(7) Pp- 2361-2371

Tiwari, A. and Sharma, A. (2013): Antifungal activity of *Anabaena variabilis* against plant pathogens. International Journal of Phrma and Biosciences. *Int J Pharm Bio Sci. Vol. 4; Pp-1030-1036.*

Tiwari, A. and Kaur, A. (2014): Allelopathic impact of Cyanobacteria on pathogenicfungi. *Int. J. Pure App. Biosci. Vol. 2(3); Pp-63-70.*

Yadav, S., Sinha, R. P., Tyagi, M. B. and Kumar, A. (2011): Cyanobacterial secondarymetabolites. *Int. J. Pharm. Bio Sci. Vol.2(1) (B); Pp-144-167.*

===***===