



Phytochemical screening and evaluation of the antidepressant activity of *Mimusop selengi* Linn. unripe Fruits in depressed mice

Uma^a, Pankaj Kumar^{b,c}, Preeti^d, Chinkey Mittal^e, Deepika^e Meher Priya^f,
Deepika Ahlawat^g, Gaurav Agarwal, Ritu^{*a,h}, Sonia Narwal^{i*}

^a Department of Pharmacy, School of Healthcare and Allied Sciences, G D Goenka University, Sohna, 122103

^b Dharamputra College of Pharmacy, Chirana, Sonepat, Haryana, 131306

^c MM College of Pharmacy, Maharishi Markandeshwar (deemed to be) University, Mullana, Ambala, Haryana, 133207

^d Gurugram Global College of Pharmacy, Haily Mandi Rd, Farukh Nagar, Haryana, 122506

^e Adarsh Vijendra Institute of Pharmaceutical Sciences, Shobhit University, Gangoh, UP, 247341

^f School of Medical and Allied Sciences, K.R. Mangalam University, Gurugram, Haryana, 122103

^g Faculty of Pharmaceutical Sciences, PDM University, Bahadurgarh, Haryana, 124508


^h Department of Pharmacology, Delhi Pharmaceutical Sciences and Research University, Delhi, 110017

ⁱ Department of Pharmacy, Panipat Institute of Engineering and Technology, Samalkha, Panipat, Haryana, 132101

*Address correspondence to these authors at the ^aDepartment of Pharmacy, School of Healthcare and Allied Sciences, G D Goenka University, Sohna, 122103, ^bDepartment of Pharmacology, School of Pharmaceutical Sciences, Delhi Pharmaceutical Sciences and Research University, New Delhi 110017, India;

E-mail: ritumalik1989@gmail.com, ⁱ Department of Pharmacy, Panipat Institute of Engineering and Technology, Samalkha, Panipat, Haryana 132101, India; E-mail: sonianarwal33@gmail.com

Graphical Abstract:



Article History

Volume 6, Issue 12, 2024

Received: 26 June 2024

Accepted: 8 July 2024

doi:

10.48047/AFJBS.6.12.2024.4134-4149

Abstract:

Background: *Mimusopselengi Linn.*(ME) has been traditionally utilized for its antianxiety, cytotoxic, antimicrobial, and antioxidant properties. It is also used in cancer treatment and serves as a diuretic, anti-inflammatory, anti-HIV, and hypotensive agent. This study aims to assess the antidepressant activity of unripe fruits of *Mimusopselengi* Linn. in a mouse model of depression.

Materials and Methods: Methanol extraction was conducted on unripe fruits of *Mimusopselengi* Linn. using a soxhlet apparatus. Mice were divided into five study groups a Normal Group, a Depression-Induced Control Group without treatment, a Standard Group (Imipramine,10 mg/kg,i.p) ,*Mimusopselengi* (200 mg/kg,p.o) and *Mimusopselengi* (400 mg/kg,p.o) for 21 days. Immobilization time was recorded in seconds and evaluated using the Tail Suspension Test (TST) and Forced Swimming Test (FST). Additionally, the total phenol and total flavonoid contents of the extracts were measured. Furthermore, various biochemical parameters, including cortisol levels, glutathione (GSH), superoxide dismutase (SOD), and catalase, were assessed. Subsequently histopathological studies were also carried out in all the studied groups.

Result: The total phenolic content and total flavonoid content were found to be 3.64 µg/ml,2.56 µg/ml respectively. Other phytoconstituents such alakaloids, tannins, flavanoids, glycosides, steroids were also detected in the extract using qualitative analysis methods. The elevated levels of phenols and flavonoids are believed to contribute to the antidepressant effects observed, comparable to the standard antidepressant imipramine. In

comparison to the standard, the 400 mg/kg test of (ME) was found to be more effective in reducing immobility time in both the FST and TST respectively. Furthermore, depressed mice exhibited higher cortisol levels compared to the standard and test groups. However, the 400 mg/kg of (ME) test dose significantly

lowered cortisol levels in depressed mice compared to both the standard and the 200 mg/kg of (ME) test dose. Moreover, In histopathological study it was found that a pyramidal cells in hippocampus area were degenerated in depressed brain as compared to standard and treatment.

Conclusions: The study highlights the therapeutic potential of *Mimusop selengi* unripe fruits as antidepressant and antioxidant activity. These findings underscore the importance of natural products as potential therapeutic agent in the management of depression.

Keywords: *Mimusop selengi*, depression, hippocampus, cortisol, imipramine

Introduction

The brain is a complicated organ that regulates many different body functions, including respiration, temperature regulation, hunger, touch, cognition, memory, emotions, and motor skills. It is essential for interpreting sensory information and storing memories while controlling movements. According to the World Health Organization (WHO), depression is characterized by feelings of guilt, sadness, disturbed sleep or appetite, loss of interest, tiredness, and poor concentration. WHO (2022) Approximately 280 million people worldwide suffer from depression, which is 50% more common among women than men. Notably, over 10% of pregnant women experience depression. According to the National Mental Health Survey (NMHS) conducted in 2015-16, the lifetime prevalence of depressive disorders (DD) was estimated at 5.25%, with a current prevalence of 2.68%. Google Trends analysis from 2018 to 2020 revealed that the number of Indians seeking information about depression symptoms doubled, peaking during the COVID-19 pandemic. A Deloitte survey conducted between 2021 and 2022 found that 59% of employees reported experiencing symptoms of depression, including sadness, lack of interest, fatigue, concentration issues, and poor decision-making. The UNICEF survey of 2021 reported that 14% of 15 to 24-year-olds in India frequently felt depressed or disinterested. WHO estimates from 2015 indicated that 4.5% of the Indian population, approximately 56,675,969 people, were affected by depressive disorders. Additionally, recent studies have shown a rise in depression rates among young adults in India, with prevalence rates ranging from 31% to 57%. *Mimusop selengi* plant also known as "Bakul" or "Spanish cherry" is a potential herb with belong to Sapotaceae family. Taxonomical classification of *Mimusop selengi* belongs to the plant kingdom and falls under the order Ericales³. It is a member of the sapotaceae family, with the genus being *Mimusops* and the species specifically identified as elengi². The tree holds significance in Hinduism and is revered as a scared plant, finding mention in religious texts and playing a crucial role in ancient tradition and this plant have pharmacological activities like Antimicrobial⁴, Antifungal, Antihyperlipidemic, Antiinflamatory⁵, Antioxidant, Antipyretic⁶, Cytotoxic, Gingival bleeding, prevent Gastric ulcer, immune modulators, Hypotensive, Antiviral, Diuretics effects, Antibacterial, Analgesic, Anticonvulsant effects, Anti-cariogenic effects Dental caries, Antiurothelial activity, COVID -19⁷, Anti- hiv, Antidiabetics , wound healing⁸, anti ulcer⁹ ,reversible of memory¹⁰. Among various plants *Mimusop selengi* unripe fruits contain phenolic compounds and flavonoids, showing potential antidepressant effects through antioxidant activity and monoamine oxidase-A (MAO-A) inhibition. Ethnobotanical reviews suggest *Mimusop selengi*'s traditional use in treating disorders, including depression. Plants with indole alkaloids, terpenoids, alkaloids, phenols, flavonoids. *Mimusop selengi*(ME) have already done toxicity study¹¹. It is safe upto 2000mg/kg⁶. (ME) unripe fruits reported antioxidant property as well as superior antidepressant activity. Due to present of some phytoconstitute like quercetin⁶and gallic acid and flavanoids. This study aims,

phytochemical screening and Evaluation of Antidepressant Activity of *Mimusop selengi* Linn unripe Fruits. In the *Mimusop selengi* linn unripe fruits extract have pheonic, flavanoids, glycosides and steroids with the help this phytoconstitute we observed a antidepressant activity and antioxidant activity.

Material and method

Collection of plants and Authentication

Unripe fruits were obtained from G.D. Goenka University following rules and ethical guidelines. Initially, a botanist assisted in identifying the unripe fruits collected. Furthermore, the plant herbarium sheet underwent authentication, which was done by RHMD, CSIR-NISCPRI with authentication number IS-NISCPRI/RHMD/CONSULT/2023/4649-50. Fresh unripe fruits of *Mimusop selengi* Linn. were gathered and air-dried until a constant weight was achieved, followed by grinding into coarse powder. The hydroalcoholic extract was then prepared using the Soxhlet apparatus12.

Drugs:

Standard drug: Imipramine (Central Drug House Pvt. Ltd, India)

Mimusop selengi unripe fruits hydro alcohol extract

Animals- Swiss albino mice (protocol No.– GDGU/PO/ IAEC/2023/32). of both male and female adults, weighing 28 to 35 g, were used in this study. The mice were given a standard laboratory diet and had unlimited to get tap water. They were housed in an animal facility run by the department, with a regular 12-hour light/dark cycle. Following the rules established by the Ministry of Environment and Forest's Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), the experimental protocols were authorised by the institutional animal ethics committee.

S.NO	Group	Remark	No. of animals
1	Group I	Normal	5
2	Group II	Control (no treatment)	5
3	Group III	Standard group (Imipramine 10 mg/kg b.w., orally)	5
4	Group IV	Test dose 1 (Receive extract 200 mg/kg b.w., orally)	5
5	Group V	Test dose 2 (Receive extract 400 mg/kg b.w., orally)	5

Table 1:Study Groups

Phytochemical screening test:

The confirmatory qualitative phytochemical screening of plant extracts was conducted to identify main compounds (tannins, saponins, flavonoids, alkaloids, phenols¹³, glycosides, steroids, terpenoids) per standard protocols .

Behavior test

In the Forced Swim Test, mice of both sexes were individually placed into an open cylindrical container measuring 10 cm in diameter and 25 cm in height. The container was filled with water to a depth of 19 cm, maintained at a temperature of 25±1°C. Treatment was administered 60 minutes before the test, following the study protocol. During the experiment, each mouse was made to swim for a total of 6 minutes, with immobility time being recorded during the last 4 minutes. A mouse was deemed immobile when it stopped active struggling and floated motionless, making only the necessary movements to keep its head above water. A decrease in immobility time indicates an antidepressant-like effect.



Figure 1 Forced swim test (FST) model

Tail suspension test

According to the predetermined study design, treatment will be administered 60 minutes before the start of the study. In the Tail Suspension Test, mice will be placed at the edge of a table about 45 cm above the ground, with their tails taped approximately 1 to 2 cm from the tip. Over a 6-minute observation period, the total duration of immobility displayed by the mice will be recorded.



Figure 2: Tail suspension method

Biochemical parameters

a) Estimation of Serum Cortisol:

The E411 automated CLIA analyzer is used to measure cortisol levels in biological samples through chemiluminescent immunoassay, providing accurate and efficient results¹⁴.

b) Superoxide dismutase (sod) activity assay:

The SOD activity assay kit from Sigma-Aldrich Catalog Number: 19160¹⁵.

c) GSH (glutathione) assay:

The GSH assay kit from Sigma-Aldrich (Catalog Number: CS0260)¹⁶.

d) Catalase (CAT) Assay:

The Catalase assay kit from Sigma-Aldrich (Catalog Number: CAT100) ¹⁷.

e) Histopathology:

Histological evaluation was performed on brain samples on the last day of the experimental protocol. The brain tissues preserved in 10% neutral buffered formalin were dehydrated in graded concentrations of ethanol, immersed in xylene and then embedded in paraffin. The sections of 4 µm thickness were cut and placed on the slide using commercial Baker's mounting

fluid. Paraffin wax was removed by warming the slide gently until the wax melted and then was washed with xylene. This was followed by washings with absolute alcohol and water to hydrate the sections and stained with haematoxylin and eosin described by¹⁸. The hydrated sections were stained with haematoxylin for 15 min. The stained sections were washed with water and treated with 1% acid alcohol mixture for 20. The acid alcohol mixture was washed off with water and sections were counterstained with 1% aqueous solution of eosin for 2 minutes. After washing with water to remove excess of eosin, the sections were dehydrated using absolute alcohol and then mounted using Canada balsam as mounting agent. The slides were observed for gross histopathological changes and neutrophil accumulation.

Statistical Analysis

The results were analyzed using Grpah pad prism vers.9.0 employing ANOVA followed by Tukeys multiple comparison test analysis with P<0.001 considered significant for all values.

Result and discussion:

Phytochemical screening


The phytochemical screening reveals the presence of carbohydrates, glycosides, tannins, alkaloids, flavonoids, and steroids, while proteins and amino acids are absent. These findings suggest that the sample contains various bioactive compounds, which may contribute to its pharmacological properties.

S.no	Phytochemical screening	Observation	Result
1	Test for carbohydrate: • Molisch test	Appearance of a purple ring at the interface.	+
2	Test for protein: • Million's test	Million's test indicates a positive result by producing a red or pink-coloured precipitate.	-
3	Test for amino acid: • Ninhydrin	A positive result shows by the appearance of a complex with a purple colour in the test tube	-
4	Test for glycosides: • Cardiac glycosides (Keller-Killani test) • Saponin glycosides (Haemolysis test)	A brown ring forming between the layers indicates the presence of cardiac steroid glycosides, confirming a positive test even at low concentrations. Haemolysis of blood	+
5	Test for tannins: • 5% FeCl ₃	Dark blue coloured dappered	+
6	Test for alkaloid: • Wagner's reagent test	Appearance of a reddish colour indicates the presence of alkaloids.	+
7	Test for flavonoids: • H ₂ SO ₄ Test	Orange to red flavonoids (flavones) is present.	+

8	Test for steroid: • Salkowski reaction	Chloroform appears red and acidic layers show greenish fluorescence indicates steroid presence in a sample.	+
---	---	---	---

Table 2: Phytochemical screening test**2 Total Phenolic content determination by folincioalteau reagent**

Concentration $\mu\text{g}/\text{ml}$	UV absorbance
10	0.609
20	0.715
30	0.793
40	0.932
Sample	0.879

Table 3: UV absorbance of total Phenolic content**Figure 3: Calibration curve of total phenolic content****3 Total flavonoid determination by aluminium chloride method**

The aluminum chloride method provides a straightforward and effective approach to determine the total flavonoid content in plant extracts, which helps evaluate their potential health benefits and antioxidant properties. Higher absorbance values indicate higher concentrations of total flavonoid compounds in the plant extract¹⁹.

Concentration	Uv absorbance
10 $\mu\text{g}/\text{ml}$	1.195
20 $\mu\text{g}/\text{ml}$	1.487
30 $\mu\text{g}/\text{ml}$	1.775
40 $\mu\text{g}/\text{ml}$	1.98
Sample	1.624

Table 4 UV absorbance total flavonoid content




Figure 4: Calibration curve total flavanoids content

4 Effect of *Mimusopspelengi* Unripe Fruit Extracts on Immobilization Time in the Forced Swim Test (FST)

The normal group exhibited baseline immobility time. In contrast, the disease control group showed increased immobility, which is indicative of depressive-like behavior²⁰. The standard group displayed significantly reduced immobility, confirming the antidepressant effects of the standard treatment. (ME 200 mg/kg) demonstrated decreased immobility compared to the disease control group, suggesting a potential antidepressant effect at this dose. (ME 400 mg/kg) showed even further reductions in immobility.




Fig5: Effect of ME on immobilization time: Data is represented as mean \pm SEM. Two wayanova followed by Tukeys multiple comparison test. @P<0.001 vs 1ST day of standard group, #P<0.001vs 1st day of ME(200), \$P<0.001vs 1 st day, ME(400). *P<0.001 vs @7thday, *1P<0.001 vs #7th day, *2P<0.001 vs \$7th day. &P<0.001 vs *14day, &1P<0.001 vs *14 day ,&2P<0.001 vs *2 14th day

5 Effect of *Mimusopspelengi* Unripe Fruit Extracts on Immobilization Time in the tail suspension test

The normal group exhibited baseline immobility time in the tail suspension test (TST), a common test for assessing depressive-like behavior in rodents. The disease control group showed increased immobility, indicating a depressive-like state²¹. The standard group displayed significantly reduced immobility, confirming the antidepressant effects of the standard treatment.

ME(200 mg/kg) demonstrated decreased immobility compared to the disease control group, suggesting potential antidepressant effects at this dose in the TST. Test G(*ME* 400 mg/kg) showed further reductions in immobility, indicating a dose-dependent response to the treatment in the tail suspension test.




Fig6: Effect of ME on immobilisation time sec.: Data is represented as mean±SEM. Two-way analysis of variance followed by Tukey's multiple comparison test. #P<0.001 vs 1st day standard group , ##P<0.001 vs 1st day of ME(200)mg/kg, ###P<0.001 vs 1st day of ME(400)mg/kg.*P<0.001 vs # 7th day, **P<0.001 vs ## 7th day, ***P<0.001 vs ###7th day, \$P<0.001 vs *14th day , \$\$P<0.001 vs **14th day,\$\$\$P<0.001 vs ***14th day.

6 Effect of *Mimusopselengi* Unripe Fruit Extracts on cortisol level

In depression, there is often dysregulation of the HPA axis, leading to elevated cortisol levels, especially in chronic stress conditions²². After treatment with *Mimusopselengi* unripe fruit hydroalcoholic extract, cortisol levels were significantly decreased ($p < 0.001$) in individuals with depression, indicating a beneficial effect on HPA axis dysregulation associated with elevated cortisol levels, particularly in chronic stress conditions.

Normal	Control	Standard	Test dose 200mg/kg	Test dose 400mg/kg
349.05	490.00	355	380	345
345	499	368	380	348
320	460	363	399	344

Table: 5 Serum cortisol level




Figure7: Effect of *Mimusopselengi* (ME) on cortisol level (ng/ml): Data is represented as mean \pm SEM. One wayanova followed by Tukeys multiple comparison test. P<0.001 *Disease Control versus **standard group, P<0.001 *** ME 200 versus *disease control, P <0.001 *disease control versus #ME 400 mg/kg.

7 Effect of *Mimusopselengi* Unripe Fruit Extracts on GSH level

Assessment of glutathione (GSH) levels post-treatment with *Mimusopselengi* unripe fruit extracts reveals potential antioxidant enhancement, protecting cells from oxidative stress²³. Increased GSH suggests improved defenses, while decreased levels may indicate GSH utilization or altered metabolism. The control standard group shows the lowest GSH level at 0.54, while the normal group has the highest at 1.36. The standard group (1.30) and the test dose 400 mg/kg group (1.26) also exhibit relatively higher GSH levels compared to the control or standard. However, the test doses (200 mg/kg and 400 mg/kg) do not consistently show a significant increase in GSH levels compared to the standard or normal group.




Figure 8: Effect of *Mimusopselengi* (ME) on GSH level (mmoles/mg): Data is represented as mean \pm SEM. One way anova followed by Tukeys multiple comparison test. P<0.001 *Disease Control versus **standard group, P<0.001 *** ME 200 versus *disease control, P <0.001 *disease control versus #ME 400 mg/kg

8 Effect of *Mimusopselengi* Unripe Fruit Extracts on SOD level

The study evaluated the impact of *Mimusopselengi* (ME) unripe fruit extracts on Superoxide Dismutase (SOD) levels, an enzyme critical for antioxidant defense²⁴. In the experimental setup, SOD levels were measured across different groups: normal, control, standard (possibly a positive control or reference), and two test groups administered with ME at doses of 200 mg/kg and 400 mg/kg. Results showed varying SOD levels across the groups, with ME administration generally showing trends toward increased SOD activity compared to the control group.

Normal ME	Control	Standard	ME200mg/kg	ME 400mg/kg
0.8	0.4	0.68	0.62	0.65
0.79	0.35	0.67	0.60	0.68
0.77	0.32	0.69	0.55	0.63

Table: 6 SOD level




Figure 9: Effect of *Mimusopselengi* (ME) on SOD level (unit/mg protein): Data is represented as mean \pm SEM. One-way anova followed by Tukeys multiple comparison test. P<0.001. *Disease Control versus **standard group, P<0.001 *** ME 200 versus *disease control, P <0.001 *disease control versus #ME 400 mg/kg.

9 Effect of *Mimusopselengi* Unripe Fruit Extracts on Catalyse

Experimental groups included a normal control, a disease control, a standard group (potentially treated with a known drug or placebo), and two test groups administered with *Mimusopselengi* at doses of 200 mg/kg and 400 mg/kg²⁵. Results demonstrated varying catalase levels across groups, with the standard group and the 400 mg/kg dose showing higher catalase activity compared to controls. These findings suggest that *Mimusopselengi* extract may enhance antioxidant defenses by potentially increasing catalase activity, which could be beneficial in combating oxidative stress-related conditions.

Normal	control	Standard	Test dose 200mg/kg	Test dose 400mg/kg
3.10	1.50	3.70	3.30	4.00
3.05	1.55	3.65	3.40	3.99
3.01	1.60	3.77	3.55	3.89

Table:7 Catalyse

catalyse




Figure 10: Effect of *Mimusopselengi* (ME) on catalyse (unit/mg protein): Data is represented as mean±SEM. One way anova followed by Tukeys multiple comparison test. . P<0.001 *Disease Control versus **standard group, P<0.001 *** ME 200 versus *disease control, P <0.001 *disease control versus #ME 400 mg/kg

Histopathology

The hippocampus, crucial for memory and emotional regulation, is closely linked to depression. Depressed individuals often have reduced hippocampal volume due to stress-inhibited neurogenesis. This region regulates the HPA axis, affecting stress responses, and contains receptors for neurotransmitters like serotonin, altered in depression²⁶. Treatments like antidepressants can reverse hippocampal changes²⁷. Inflammation also impacts the hippocampus, linking it to depressive symptoms.




Figure 11: Histopathology of brain tissue hippocampus

Discussion:

Depression is marked by feelings of guilt, sadness, disrupted sleep or appetite, loss of interest, fatigue, and difficulty concentrating. Known as a depressive disorder, it is a prevalent mental health condition. Despite the prevalence and severity of depression, current antidepressant medications are often associated with high costs and a range of serious side effects, including cardiotoxicity, myocardial infarction, hypertension, obesity, hepatotoxicity, kidney and lung cancer, respiratory distress syndrome, and more common issues like blurred vision, constipation, dizziness, and weight gain. This highlights the urgent need for alternative treatments. Natural products hold promise in the development of new antidepressants, offering potential benefits with significantly fewer or no side effects, presenting a hopeful avenue for safer and more effective depression management. Plants based secondary metabolites like flavanoids, alkaloids, glycosides, tannins and phenols have been shown to have promising benefits in diseases such as depression and antioxidants. These secondary metabolites have been proven to target multiple mechanism pathways. The current study has selected antidepressant activity of *Mimusop selengi linn* (ME) unripe fruits have been studied. Through various disease models.

The current study exhibited depression development in swiss albino mice, represented by significantly elevated biochemical parameters such as serum cortisol, GSH level, catalyse activity and SOD level in depressed mice. Elevated cortisol level is an excellent clinical marker

of Patients with major depressive disorder²⁸with the help of selected plant extract decrease the serum cortisol level in after treatment the extract. It has been reported that *Crocus sativus* Saffron at doses of 30 mg/day was found to be as effective as imipramine 100 mg/day and fluoxetine 20 mg/day in treating mild to moderate depression in adult patients²⁹. *Rosmarinus officinalis* (Rosemary): Rosemary extracts consistently decreased immobility time in the forced swim test (FST) and tail suspension test (TST) in mice, indicating antidepressant-like effects. The antidepressant activity likely involves interactions with the monoaminergic system³⁰. In our study, we noticed a reduction immobility in the depressed miceas compared to the control group. The study showed the development of depression in mice by displaying higher immobility time and higher cortisol level. Although ME extract reduced immobility time and serum cortisol level. This indicates that the ME extract as protective potential against depression disorder and antioxidant activity³¹.

However, it has been reported induce foot shock method by pole climbing apparatus.various biochemical and behvaioural parameters were estimated at the end of 21 days of the study. Multiple behavior models to evaluate of antidepressants activity are tail suspension test, forced swim test based on the literature we evaluate antidepressant activity with the two test we observed that immobility period decreased significantly in treated mice as compared to depressed mice, simlilarly forced swim test immobility time reduced thus it can infer that ME thus potential antidepressant activity. same result was evaluated with imipramine (10mg/kg) as standard drug. In depression multiple regions of brain are affected as revealed by many studies in one of the study Studies show that patients with depression have decreased gray matter volume (GMV) in the hippocampus and prefrontal cortex, affecting memory, learning, and emotion regulation. The hippocampus and dorsolateral and ventromedial prefrontal cortex regions are particularly impacted, contributing to cognitive and executive dysfunction. The amygdala shows variable GMV changes, indicating its role in emotional dysregulation. Subcortical regions like the thalamus, caudate nucleus, and insula also exhibit reduced GMV, though less consistently. Additionally, depression is associated with disrupted functional connectivity in corticolimbic circuits, affecting emotion, cognition, and stress response, highlighting both structural and connectivity changes in depression. So, in present study take hippocampus region and observed py(Q)pyramidal cells are decreased in case of depression.

Conclusion

This study suggests that the *Mimusop selengi*unripe fruits extract holds promise as a natural treatment for antidepressants activity, attributed to its significant antioxidant properties. The study highlights the potential antidepressant properties of *Mimusop selengi* unripe fruit extract, demonstrated through significant reductions in immobility time in both Forced Swim Test (FST) and Tail Suspension Test (TST) models. These effects compare favorably to the standard antidepressant imipramine. Phytochemical analysis suggests that compounds such as triterpenoids and flavonoids may contribute to these effects by interacting with neurotransmitter systems involved in mood regulation. However, further research is needed to isolate and understand the specific mechanisms of action, as well as to evaluate long-term safety and efficacy. Moving forward, clinical trials are crucial to validate these findings and explore the potential of *Mimusop selengi* as a natural and effective treatment for depression.

References

1. Langenbach BP, Koelkebeck K, Knoch D. Mentalising and depression: a mini-review on behavior, neural substrates, and treatment options. *Front Psychiatry*. 2023;14:1-40. doi:10.3389/fpsyg.2023.1116306

2. Brigitta B. Pathophysiology of depression and mechanisms of treatment. *Dialogues Clin Neurosci.* 2002;4(1):7-20. doi:10.31887/dcns.2002.4.1/bbondy
3. Chaovanamethakul P, Suwannatee H, Chaisuksant R, Suntornwat O. Antioxidant capacity and phenolic content of bullet wood (*Mimusops elengi*) fruit extract. *Acta Hortic.* 2008;787:301-305. doi:10.17660/actahortic.2008.787.36
4. Author C, Abu Sayeed M, Abbas Ali M, Abdul Mojid M, Sarmina Yeasmin M, Mohal Khan A. An Evaluation of Antimicrobial Activities of *Mimusops elengi* Linn. *Res J Agric Biol Sci.* 2008;4(6):871-874.
5. Kar B, Suresh RB, Karmakar I, Dola N, Bala A. activities of *Mimusops elengi* leaves. Published online 2024:1-5.
6. Purnima A, Koti BC, Thippeswamy AHM, et al. Antiinflammatory, analgesic and antipyretic activities of *Mimusops elengi* Linn. *Indian J Pharm Sci.* 2010;72(4):480-485. doi:10.4103/0250-474X.73908
7. Sai Ramesh A, Adarshan S, Lohedan H, et al. Computational analysis of the phytocompounds of *Mimusops elengi* against spike protein of SARS CoV2 – An Insilico model. *Int J Biol Macromol.* 2023;245:1-24. doi:10.1016/j.ijbiomac.2023.125553
8. Article R, Bakul L, Medicinal AP, Review PA, Gupta PC. ISSN (Print). Published online 2013.
9. Shah PJ, Gandhi MS, Shah MB, Goswami SS, Santani D. Study of *Mimusops elengi* bark in experimental gastric ulcers. *J Ethnopharmacol.* 2003;89(2-3):305-311. doi:10.1016/j.jep.2003.09.003
10. Joshi H, Parle M. Reversal of memory deficits by ethanolic extract of *mimusops elengi* linn. in mice. *Pharmacogn J.* 2012;4(29):30-39. doi:10.5530/pj.2012.29.5
11. Ketchen DJ, Shook CL. Institutional Login □. *Strateg Manag.* Published online 1996:23-25.
12. Willner P, Towell A, Sampson D, Sophokleous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology (Berl).* 1987;93(3):358-364. doi:10.1007/BF00187257
13. Khan S. Phytochemical screening publication. Published online 2022:625-626.
14. Bromet E, Andrade LH, Hwang I, et al. Cross-national epidemiology of DSM-IV major depressive episode. Published online 2011.
15. Chakraborty I, Kunti S, Bandyopadhyay M, Dasgupta A, Chattopadhyay GD, Chakraborty S. Evaluation of serum zinc level and plasma SOD activity in senile cataract patients under oxidative stress. *Indian J Clin Biochem.* 2007;22(2):109-113. doi:10.1007/BF02913326
16. Rahman I, Kode A, Biswas SK. Assay for quantitative determination of glutathione and glutathione disulfide levels using enzymatic recycling method. *Nat Protoc.* 2007;1(6):3159-3165. doi:10.1038/nprot.2006.378
17. Erkili K, Evereklioglu C, Duygulu M, Dogan H, Çekmen, Abdullah. Published online 2024:17-19.
18. CLAYDEN AD, PARKHOUSE J. Allocation of preregistration posts. *Med Educ.* 1971;5(1):5-12. doi:10.1111/j.1365-2923.1971.tb02144.x
19. Shraim AM, Ahmed TA, Rahman MM, Hijji YM. Determination of total flavonoid content by aluminum chloride assay: A critical evaluation. *Lwt.* 2021;150(October 2021):1-9. doi:10.1016/j.lwt.2021.111932

20. Yankelevitch-Yahav R, Franko M, Huly A, Doron R. The forced swim test as a model of depressive-like behavior. *J Vis Exp.* 2015;2015(97):1-11. doi:10.3791/52587
21. Cryan JF, Mombereau C, Vassout A. The tail suspension test as a model for assessing antidepressant activity: Review of pharmacological and genetic studies in mice. *Neurosci Biobehav Rev.* 2005;29(4-5):571-625. doi:10.1016/j.neubiorev.2005.03.009
22. Tafet GE, Idoyaga-Vargas VP, Abulafia DP, et al. Correlation between cortisol level and serotonin uptake in patients with chronic stress and depression. *Cogn Affect Behav Neurosci.* 2001;1(4):388-393. doi:10.3758/CABN.1.4.388
23. Pal SN, Dandiya PC. Glutathione as a cerebral substrate in depressive behavior. *Pharmacol Biochem Behav.* 1994;48(4):845-851. doi:10.1016/0091-3057(94)90191-0
24. Bhatt S, Nagappa AN, Patil CR. Role of oxidative stress in depression. *Drug Discov Today.* 2020;25(7):1270-1276. doi:10.1016/j.drudis.2020.05.001
25. Hall WD. Australian and New Zealand Journal of Psychiatry people. 2006;33:1-33.
26. Sharma U. Depression : Understanding the Illness.
27. Campbell S, MacQueen G. The role of the hippocampus in the pathophysiology of major depression. *J Psychiatry Neurosci.* 2004;29(6):417-426.
28. Sahu M, Dubey R, Chandrakar A, Kumar M, Kumar M. A systematic review and meta-analysis of serum and plasma cortisol levels in depressed patients versus control. *Indian J Psychiatry.* 2022;64(5):440-448. doi:10.4103/indianjpsychiatry.indianjpsychiatry_561_21
29. Dhingra D, Sharma A. A review on antidepressant plants. *Indian J Nat Prod Resour.* 2006;5(2):144-152.
30. Arun Pardhe H, C Nagalakshmi N, M.G H, Kumar Chourasia P, S N. A review: Medicinal plants with antidepressant properties. *IP Indian J Neurosci.* 2020;6(1):1-5. doi:10.18231/j.ijn.2020.001
31. Tayal V, Kalra B, Chawla S. Evaluation of antidepressant activity of tramadol in mice. *Indian J Pharmacol.* 2008;40(3):129-130. doi:10.4103/0253-7613.42307
32. Behavioura WP. Willner P. Behavioura. Published online 2024:2024.
33. Shalam M, Shantakumar S, Narasu M. Pharmacological and biochemical evidence for the antidepressant effect of the herbal preparation Trans-01. *Indian J Pharmacol.* 2007;39(5):231-234. doi:10.4103/0253-7613.37273