



African Journal of Biological Sciences



Original article

Assessment of *in vitro* Antioxidant activity, Total Phenolic and Total Flavonoid in *Careya arborea* Fruit Pulp Extracts

Daya L. Chothani¹*, Dipen R. Bhimani

¹B. K Mody Government Pharmacy College, Rajkot-360003, Gujarat, India

* *Correspondence:*

Dr. Daya L. Chothani

B. K. Mody government Pharmacy College,
Near Aji dam chokdi, Polytechnic campus,
Rajkot-360003

Email id: daya.herb@gmail.com

+91-9924214110

ABSTRACT

In the current investigation, *Careya arborea* fruit pulp was used to assess total phenolic, flavonoid and *in vitro* antioxidant potential. The ethyl acetate extract (EAE), methanolic extract (ME), water extract (WE) of fruit pulp were studied. Using the Folin-Ciocalteu reagent, total phenolics were measured, and the colorimetric method was used to determine the flavonoid concentration. Antioxidant activity was evaluated using DPPH assay and reducing power by FeCl₃. A positive control was ascorbic acid. The fruit pulps from *C. arborea* with the highest phenolic, flavonoid, and antioxidant potential were extracted using ethyl acetate. When compared to a positive control, the *C. arborea* fruit showed a substantial *in vitro* antioxidant activity ($P < 0.05$). The Finding concluded that the *C. arborea* fruit pulp exhibited antioxidant activity.

KEYWORDS: DPPH free radical scavenging activity, *Careya arborea*, Fruit pulp, Total phenolic

Article History

Volume 6, Issue 11, 2024

Received: 02 Jun 2024

Accepted: 15 Jun 2024

doi: [10.48047/AFJBS.6.11.2024.38-49](https://doi.org/10.48047/AFJBS.6.11.2024.38-49)

INTRODUCTION

The presences of unpaired electrons on atoms, molecules or ions are known as free radical. Chemically, they are unstable and reactive. ^[1] The body contains free radicals, which are produced during metabolic activities. Bacteria and viruses can be eliminated by free radicals generated by the immune system. They also have a part in the synthesis of hormones, energy, and enzyme activation. Excessive formations of free radical damage tissue Radiation, the sun, medical X-rays, pollution, etc. can all boost the production of extra free radicals. High-fat diets have the potential to promote the production of free radicals. The free radical can be destroyed first line body defense system. Free radicals are neutralized by enzymes such as superoxide dismutase, glutathione peroxides, vitamin C, vitamin E, selenium mineral, and hormone melatonin. In cases of severe or persistent oxidative stress, protection might not be sufficient. Therefore, to keep free radicals in balance, a certain amount of exogenous antioxidants or free radical scavengers is always needed. ^[2-4] Free radicals are responsible for development of disorders like diabetes, stroke, arteriosclerosis, cancer, the aging process and cardiovascular diseases. ^[5-7] Different types of Phytoconstituents are found in plants. The phenolic compound is an effective reducing agent ^[8-9]

Careya arborea Roxb is a member of the Lecythidaceae family, commonly known as wild Guava and Kumbhi (Because the fruits resemble earthen pots). ^[10] The calyx limb is topped with big, globose, meaty, indehiscent fruits. Many seeds, deeply buried in soft pulp. In Ayurveda, the fruits are erotic and caustic, and they heal kapha. ^[11] It is found in the Sub-Himalayan region, extending eastward from Jammu to West Bengal, Madhya Pradesh, and Tamil Nadu. ^[12] Fruits are uses as an astringent, decoction aids digestion, ^[13] astringent, demulcent, ^[14-17] edible and aromatic. ^[18] Fruit powder is used to treat obesity and prevent coughing. Applying fruit pulp to the scalp promotes hair development. ^[19-20] *Careya arborea* bark has reported Central nervous system depressant, ^[21] antimicrobial and antioxidant, ^[22] antidiarrhoeal, ^[23] hepatoprotective, ^[24] antitumor, ^[25] analgesic, ^[26] and anticonvulsant activity. ^[27] *C. arborea* fruits pulp remains unexplored. Therefore, the current study was planned to determining their total phenolic components, flavonoid concentration, and antioxidant potential of *C. arborea* fruit pulp

MATERIALS AND METHODS

Plant material

Plant materials in June 2012 of *C. arborea* were procured from Vadodara. Plant was identified and authenticated at Botany Department of The M. S. University, Vadodara. Voucher specimen (DC-CA-2) was kept in our lab's herbarium.

Reagent and Chemicals

Analytical grade reagents and chemicals were all utilized. Folin Ciocalteu reagent, Quercetin, Aluminum chloride, Gallic acid, Potassium acetate, Sodium carbonate α , α Diphenyl – β Picryl Hydrazyl (DPPH), Ascorbic acid were procured from E. Merck (Darmstadt, Germany), Hi-Media lab. Ltd (Mumbai) and Sigma (Chemical Co, St. Louis, MO, USA). A Shimadzu UV-1800 was used to record each and every UV-Vis measurement.

Preparation of extracts

About 20 grams of fruit pulp powder were extracted using methanol, water, and ethyl acetate individually for duration of 24 hours. After filtering, the extracts underwent evaporation to concentrate on water bath and dried.

Phytochemical screening ^[28]

Chemical test were performed for the presence of phytoconstituents such as alkaloid, phenolic, flavonoid, saponin and sterols, by employing chemical tests.

Thin layer chromatographic profile of phenolic compounds ^[29-30]

The mobile phase used to create the TLC profile of the extracts was CHCl_3 : ethyl acetate: Formic acid (2:8:0.4) as mobile phase. Alcoholic FeCl_3 was sprayed onto the plate. The R_f values of observed compounds were recorded.

Determination of total phenolic content ^[31]

Folin-Ciocalteu reagent was used to quantify total phenolic (Singleton and Rossi, 1965). The amount of total phenolic was measured in the fruit pulp of *C. arborea* in the methanol extract (ME), ethyl acetate extract (EAE), and water extract (WE). The standard was gallic acid. Methanol was used to take the gallic acid calibration curve.

In a volumetric flask, 1.0 ml of test sample containing 1 mg extract was diluted with 10 ml of distilled water. It was mixed with 1.5 ml of folin ciocalteu reagent. After letting the mixture for five minutes, 4 ml of a 20% sodium carbonate solution was added, and distilled water was used to bring the volume up to 25 ml. After this mixture was left for thirty minutes, and the absorbance measured with a Shimadzu 1800 spectrophotometer at 765 nm of the blue color developed. The calibration curve of gallic acid was used to determine the percentage of total phenolics. Gallic acid percentage was used to express the total phenolic content.

Determination of total Flavonoid content ^[32-35]

The method was performed on methanol extract, ethyl acetate extract and water extract (WE) of *C. arborea* fruit pulp.

Aluminum chloride colorimetric method

The standard utilized was quercetin. 1 mg of Quercetin was dissolved in 100 ml methanol to produce (10 μ g/ml). Methanol was used to take the quercetin calibration curve. The standard solutions and extracts were separately mixed with 1.5 ml of 95% methanol, 0.1 ml of 10 % aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. The absorbance of the reaction mixture was measured at 415 nm in a Shimadzu 1800 spectrophotometer following a 30-minute incubation period at room temperature. The same volume of distilled water was used in place of 10% aluminum chloride in the blank.

The percentage of total flavonoids was calculated from calibration curve. Flavonoid content was expressed as % Quercetin.

DPPH free radical scavenging activity ^[36-39]

Ascorbic acid was dissolved in methanol to get the concentration of 30, 45, 60, 75, 90 and 100 μ g/ml. The test samples were dissolved in methanol to give stock solution of 1000 μ g/ml. 50, 100, 150, 200, 250 and 300 μ g/ml and ethyl acetate extract 25,50, 75,100, 125 and 150 μ g/ml concentration were prepared with methanol. Accurately 1.3mg of DPPH was dissolved in 1 ml of methanol, the test covered with aluminum foil to protected from the light.

DPPH free radical scavenging activity

A 75 µl DPPH solution was added to 3 ml methanol and the absorbance was measured immediately at 516 nm for control reading. Different concentrations of standard and test samples were diluted with methanol up to 3 ml and added 75 µl of DPPH. Using a Shimadzu 1800 spectrophotometer, the absorbance was measured right away at 516 nm after the addition of the DPPH solution, with methanol serving as a blank. After 30 minutes, the absorbance decreased in the presence of test samples at various concentrations, and the percentage reduction and IC₅₀ were calculated using the following formula:

$$\% \text{ Free radical scavenging activity} = \frac{\text{Control absorbance} - \text{Sample absorbance}}{\text{Control absorbance}} \times 100$$

Reducing power by FeCl₃ [40-42] ethanol was used to dissolve the extracts and ascorbic acid (standard). The reducing power of methanolic extract and ethyl acetate extracts of *C. arborea* fruits were determined according to the method of Oyaizu (1986). Extracts were mixed with 2.5ml 0.2 M phosphate buffer (pH 6.6) and 2.5ml potassium ferricyanide (1%). After the mixture was incubated at 50 °C for 20 min, 2.5ml of TCA (10%) were added and the mixture was centrifuged at 3000 rpm for 10 min. Supernatant (2.5ml) was mixed with distilled water (2.5 mL) and 0.5 ml ferric chloride (0.1%), kept for 10 min. A similar procedure was used to generate the control, excluding the extracts, and the absorbance at 700 nm was measured and compared to the standard. Improved reducing power is indicated by a higher sample absorbance.

RESULTS AND DISCUSSION

Phytochemical screening

Fruits that underwent phytochemical screening revealed the presence of flavonoids, proteins, sterols, carbohydrates, saponins, and phenolic compounds

TLC profile for phenolics compounds

Phytoconstituents are identified and separated using thin layer chromatography.



Figure 1. TLC profile of phenolics compounds of *C. arborea* Fruit pulp extracts.

A-Ethyl acetate extract and B- methanolic extract

Total phenolics content

Chemically, Folin-Ciocalteu reagent is phosphomolybdate and phosphotungstate. Under alkaline circumstances, phenolic compounds reduce the phosphotungstate-phosphomolybdate complex to blue reaction products. It measures the amount the quantity of the material required to prevent the reagent from oxidizing. Table 1 presented the findings

Flavonoid content

Aluminum chloride colorimetric method

A colorimetric technique for aluminum chloride, with the C-4 keto group and either the C-3 or C-5 hydroxyl group of flavones, isoflavons, and flavonols, aluminum chloride (AlCl_3) forms acid stable complexes. Furthermore, AlCl_3 combines with the ortho-dihydroxyl groups in the flavonoid's A or B ring to generate acid-labile complex. ^[38] Results were reported in Table 1.

Table 1: Total phenolic, Flavonoid content and DPPH assay IC₅₀ of *C. arborea* Fruit pulp extracts.

Extracts	% W/W Flavonoid compounds	% W/W (phenolic content)	IC ₅₀ (µg/ml) Value DPPH Assay
Water extract(WE)	0.254±0.012	2.70±0.2804	----
Methyl extract	0.128±0.002	3.956±0.3606	94±0.5292
Ethyl acetate extract(EAE)	0.258±0.011	4.55±0.5292	26.5±0.5
Ascorbic acid(AA)	---	----	53± 0.7638
Values represent mean ± standard deviation (n=3)			

The results indicated that the *C. arborea* fruit pulp methanol and water extract had lower amounts of phenolic and flavonoid compounds than the ethyl acetate extract.

Anti oxidant activity *in vitro*

The methanol and ethyl acetate extracts of *C. arborea* fruit pulp were used to study the *in vitro* free radical scavenging activity and reduction power by FeCl₃.

DPPH free radical scavenging activity

DPPH (1, 1-diphenyl-2-picrylhydrazyl) is a stable free radical and deep violet color. Antioxidant creates a stable, yellow-colored diphenyl picrylhydrazine by reducing DPPH and accepting an electron or hydrogen radical. [39] The free radical scavenging capability of DPPH radical is determined by the decrease in absorbance at 516 nm. The drop in absorbance at 516 nm indicates the DPPH radical's capacity to scavenge free radicals. The activity was represented as a drop in sample absorbance at various concentration points.

The IC₅₀: radical-scavenging activity (concentration in µg required for 50% inhibition of DPPH radical) was calculated from the graph (Figure 1). The greater radical scavenging activity is indicated by the lower IC₅₀ value. The following is the order in which extracts and standards scavenge the DPPH radical: ME > AA > EAE. When compared to methanol, EAE have more radical-scavenging activity (Figure 2). Table 1 presents comparison results between several

extracts of *C. arborea* fruit pulp with respect to their ability to scavenge free radicals using DPPH. Furthermore, there exists a clear proportionality between the activity in the extracts and the concentration of total phenolics and flavonoid content.




Figure 2: Comparison of DPPH free radical scavenging activity of *C. arborea* fruit pulp.

Where, AA: Ascorbic acid, EAE: Ethyl acetate extract, ME, Methanolic extract

Compared to methanolic extract, the ethyl acetate extract of *C. arborea* fruit pulp has shown a greater capacity for DPPH radical scavenging.

Reducing power by FeCl_3

The reduction of Fe^{3+} /ferricyanide complex to the ferrous form in presence of antioxidants served as the basis for the reducing activity. The Fe^{2+} forms the Perl's Prussian blue at 700 nm. The reducing power *C. arborea* was studied using the potassium ferricyanide reduction method. As concentration increased, so did the extracts' decreasing power. Compared to the methanol extract of *C. arborea*, the ethyl acetate extract exhibited more activity. The outcome is shown in Figure.3.




Figure 3. Reductive potential of *C. arborea* fruits

Statistical Analysis

The mean \pm SD for each of the three determinations ($n = 3$) was presented in the results. One way analysis of variance (ANOVA) was used to estimate the variation, and Graph Pad Prism version 6.00 and Microsoft Excel 2007 were used. Value of $p < 0.05$ was considered as significant difference.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgment

The authors are grateful to B. K. Mody Government Pharmacy College, Rajkot for providing necessary facility to complete this research work. The authors also thank Dr. P. S. Nagar for helping with the plant collection and authentication

CONCLUSION:

The *Careya arborea* fruit pulp has showed *in vitro* antioxidant activity when performed using DPPH free radical scavenging assay and reducing power by FeCl_3 . Greater activity was shown by the ethyl acetate extract. Total phenolic and total flavonoid concentrations in the ethyl acetate extract were found to be higher than in the methanol extract. The study concludes that the antioxidant potential of *C. arborea* fruits was comparable ($P < 0.05$) to standard and that they might be utilized as a natural source of antioxidants.

REFERENCES

1. Stadtman ER. Protein oxidation and aging. *Free Radic Res* 2006; 40(12):1250-8.
2. Balch, Phyllis A. Prescription for Nutritional Healing, Penguin, 2006, p 62-64
3. Schulz– Aellen M-F. Aging and Human Longevity, pp 113-118, Birkhauser, 1997
4. Lobo V, Patil A, Phatak A and Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010 Jul-Dec; 4(8): 118–126.
5. Prior RL, Cao G. Analysis of botanicals and dietary supplements for antioxidant capacity: a review. *J. AOAC Int.* 2000; 83: 950–956.
6. Yamaguchi F, Saito M, Ariga T, Yoshimura Y, Nakazawa H. Free radical scavenging activity of garcinol from *Garcinia indica* fruit rind. *J. Agri. Food Chem.* 200; 48: 2320–2325.
7. Verma AR, Vijayakumar M, Rao V, Mathela CS. In vitro and in vivo antioxidant properties and DNA damage protective activity of green fruit of *Ficus glomerata*. *Food and Chemical Toxicology.* 2010;48: 704–709
8. Anderson D. Antioxidant defences against reactive oxygen species causing genetic and other damage. *Mutat. Res.* 1999; 350: 103–108.
9. Anagnostopoulou MA, Kefalas P, Papageorgiou VP, Assimepoulou AN, Boskou D. Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chem.* 2006; 94: 19–25.
10. Patil MV, Patil DA. Ethnobotany of Nasik District Maharashtra. Daya Books, 2006, p.338
11. Kirtikar KR, Basu BD. Indian Medicinal Plants. Vol. 2, 2nd edn, Bishen Singh Mahendra Pal Singh, Dehradun; India, 1975: 894-5.
12. Khare CP. Indian Medicinal Plants: An Illustrated Dictionary, Springer, 2007: 121-122.
13. Panda H. National Institute Of Industrial Re, 01-Jan-1999 – Herb gardens. Pp-218.
14. Pandey CN. Medicinal plants of Gujarat, Gujarat Ecological Education and Research Foundation, Gujarat, India, 2005: 146
15. Nadkarni KM. Indian material medica. Vol.2, Popular Prakashan, 1994: 273
16. Chopra RN, Chopra IC. Indigenous Drugs of India. Academic Publishers, 2006: 584
17. Pullaiah T. Medicinal Plants in Andhra Pradesh, India Daya Books, 2002 : 125
18. Negi S S .Forests and Forestry in Nepal. APH Publishing, 1994 : 141
19. Pawar S, Patil DA. Ethnobotany of Jalgaon District, Maharashtra Daya Books, 2008: 175.
20. Chothani DL, Patel NM. Quantification of gallic acid in fruit and leaves of *Careya arborea* by highperformance thin layer chromatography. *American Journal of Physiology, Biochemistry and Pharmacology.* 2014;3(4):1-5.

21. Kumar RS, Sundram RS, Sivakumar P, Nethaji R, Senthil V. Murthy NV , Kanagasabi R. CNS activity of the methanol extracts of *Careya arborea* in experimental animal model, *Bangladesh J Pharmacol* 2008; 3: 36-43 l.
22. RS Kumar, T Sivakumar, RS Sundaram, Sivakumar P, Nethaji R and Gupta M. Antimicrobial and Antioxidant Activities of *Careya arborea* Roxb. Stem Bark, *Iranian journal of pharmacology & therapeutics*, 2006, 5:35-41, 2006.
23. Rahman MT, Khan OF, Saha S, Alimuzzaman M. Antidiarrhoeal activity of the bark extract of *Careya arborea* Roxb. *Fitoterapia* 2003;74:116-8
24. Senthilkumar N, Badami S, Dongre SH, Bhojraj S. Antioxidant and hepatoprotective activity of the methanol extract of *Careya arborea* bark in Ehrlich ascites carcinoma-bearing mice. *J Nat Med.* 2008 Jul; 62(3):336-9.
25. Natesan S, Badami S, Dongre SH, Godavarthi A. Antitumor activity and antioxidant status of the methanol extract of *Careya arborea* bark against Dalton's lymphoma ascites-induced ascitic and solid tumor in mice. *J Pharmacol Sci.* 2007 Jan; 103(1):12-2.
26. Ahmed M, Rahman MW, Rahman MT, Hossain CF. Analgesic principle from the bark of *Careya arborea*. *Pharmazie.* 2002 Oct; 57(10):698-701.
27. Shinde GS, Karadi RV, Khedkar AS, Dere PJ, Mandavkar YD, Khalure PR. Screening of *Careya arborea* Roxb for their anticonvulsant properties in experimental animals. *Int J Green Pharm* 2013;7:29-33
28. Kokate CK. *Practical Pharmacognosy*, 4th ed, New Delhi: Vallabh Prakashan 2005, p. 107-111.
29. Wagner H, Blade S, Zgainsky GM. *Plant Drug Analysis*. Springer Verlag: Britain, 1984.
30. Harbone JH. *Phytochemical methods*. 2nd ed, London: Chapman and Hall. 1984:37-39.
31. Singleton VL and Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 1965; 16:144- 158.
32. Chang CC, Yang MH, Wen HM, Chern JC. Estimation of Total Flavonoid content in propolis by two complementary colorimetric methods, *J. Food Drug Anal.* 2002; 10(3):178-182.
33. Chothani DL, Patel NM. Anti-allergic potential of methanolic extract of leaves and fruits of *Careya arborea*. *Journal of Pharma Sci Tech.* 2014;4(1):29-31.
34. Chothani DL, Patel NM. Phytochemical screening and quantification of phytoconstituents in *Gmelina arborea* fruits extracts. *J. Med. Plant Res.* 2018;6(4):31-5.
35. Plummer DT. *Introduction to practical Biochemistry*, 2nd edition, 1988: 212-13.
36. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn J. Sci. Technol*, 2004; 26(2): 211-219.
37. Shimada K, Fujikawa K, Nakamura T. Antioxidative properties of xanthan on the auto oxidation of soyabean oil in cyclodextrin emulsion, *J. Agric. Food. Chem.* 1992; 40: 945-948.

38. Chothani DL, Mishra SH. In vitro anti-oxidant activity of *Ruellia tuberosa* root extracts, *Free Radicals and Antioxidants* 2012; 2(4):38-44.
39. Wang ZH, Hsu CC, Yin MC. Antioxidative characteristics of aqueous and ethanol extracts of glossy privet fruit. *Food Chemistry*.2009; 112: 914–918.
40. Oyaizu M. Studies on product of browning reaction: antioxidative activities of products of browning reaction prepared from glucosamine, *Japanese Journal of Nutrition*. 1986; 44: 307-315.
41. Mabry TJ, Markham KR and Thomas MB. *The Systematic Identification of Flavonoids*. Springer- Verlag. 1970. New York, U.S.A.
42. Soares JR, Dins TCP, Cunha A P, & Ameida LM. Antioxidant activity of some extracts of *Thymus zygis*. *Free Radical Research*, 1997; 26, 469–478