



Enhancing Cassava Productivity Through Optimal Sowing Density In Central-Western Côte D'Ivoire

Gbotto Ahou Anique^{1*}, Gnamien Yah Gwladys¹, Anzara Gnigouan Kadio Guy Roland¹, Yao Kouakou Abessika Georges¹, Yao Kouamé Célestin¹, Yao Jean Jaurès¹, Doubi Bi Tra Serge², Kouassi N'dri Jacob¹

¹Department of Agronomy and Forestry, Genetics Laboratory, University of Jean Lorougnon Guede, BP 150 Daloa, Côte d'Ivoire

²Department of Genetics, Marc Delorme Research Station, National Centre for Agronomic Research (CNRA), Côte d'Ivoire, 07 BP 13 Abidjan 07

*Corresponding author: aniquegbotto@yahoo.fr

Volume 7, Issue 6, Jun 2025

Received : 15 April 2025

Accepted : 05 May 2025

Published : 09 Jun 2025

[doi :10.48047/AFJBS.7.6.2025.607-619](https://doi.org/10.48047/AFJBS.7.6.2025.607-619)

Abstract Cassava is a crucial food crop with significant economic and nutritional importance, particularly in tropical regions. Despite its potential, cassava yields remain insufficient in Côte d'Ivoire, largely due to poor cultivation practices such as inappropriate planting densities. This study investigates the impact of sowing density, location, and variety on cassava production in the Vavoua department of central-western Côte d'Ivoire. The research focused on two cassava varieties, Yavo and Bonoua, grown under three planting densities (1m x 80 cm, 1m x 1m, and 80 cm x 80 cm) at two locations, N'Guessankro and Sanpedro2. Results indicate that the Yavo variety achieved the highest yield when planted at a low density (1m x 1m) in N'Guessankro, benefiting from favorable environmental and soil conditions. Optimal spacing facilitated efficient resource use and enhanced photochemical activity, boosting cassava growth and yield. Conversely, close planting led to increased competition for nutrients, limiting productivity. The findings underscore the importance of selecting appropriate planting density, variety, and location to maximize cassava productivity. Further studies are recommended to explore the impact of organic fertilizers and their optimal application rates on cassava yields in this region.

Keywords: Cassava, Yavo variety, sowing density, yield

Introduction

Since its independence, the agricultural sector has been the cornerstone of Côte d'Ivoire's economic development. To harness this potential, the Ivorian government has prioritized the development of industrial crops, particularly coffee and cocoa. However, food crops, critical to Côte d'Ivoire's diet, have not received significant public investment or targeted institutional support (Boni et al., 2015). As a result, the production of certain crops, such as cassava, remains low. Cassava is cultivated widely across the globe and ranks as the third most cultivated food crop after rice and maize, with an estimated global production of 302.7 million tons in 2020 (Bilong et al., 2022). Rich in carbohydrates, starch, and fiber, cassava offers significant nutritional benefits (Wang et al., 2011). During times of famine, it serves as a vital food reserve (Ceballos et al., 2006) and provides an essential source of calories in the diets of tropical regions (Borku, 2025; Montagnac et al., 2009). In Côte d'Ivoire, cassava is a key food crop, mainly grown for fresh and dried consumption. However, national production remains insufficient to meet the population's needs, which exceed 2.4 million tons annually (Kintché et al., 2017; Hillocks et al., 2002).

Several factors contribute to low cassava yields in Côte d'Ivoire, including poor soil quality, excessive reliance on chemical fertilizers, unskilled labor, the use of unsuitable varieties, and, most critically, poor cultivation practices such as suboptimal planting densities (Adjei-Nsiah et al., 2007). Planting density, which directly correlates with plant spacing, plays a pivotal role in maximizing crop yields (Gnamien et al., 2024). Recognizing the importance of these variables, a study was conducted in Vavoua, in central-western Côte d'Ivoire, to enhance cassava yields. The research aimed to test the hypothesis that the varieties used are well-suited to the Vavoua region and that the chosen planting density could optimize cassava production.

2. Material and methods

2.1. Experimental site

The study was conducted from April to July 2022. The study area is located in the centralwestern part of Côte d'Ivoire, in the department of Vavoua. It was carried out in the village of Pelezi, specifically in the Sanpedro2 and N'Guessankro areas. The study site is 91 km far from the town of Daloa and is located at 7°16'50" north latitude and 6°49'35" west longitude. This area experiences four seasons: a major rainy season from April to mid-July, a minor dry season from mid-July to mid-September, a minor rainy season from mid-September to November, and a major dry season from December to March.

2.2. Plant material, treatments, and experimental design

The plant material used in this study consists of two varieties of cassava (Yavo and Bonoua). Cassava tubers used were obtained from farmers who had previously cultivated these varieties. The experiment was laid out in a split-plot design with three replicates (blocks), implemented across two sites: Sanpedro 2 and N'Guessankro, which served as the main plot factor (environmental condition). The sub-plot factor was a combination of two cassava varieties (Yavo and Bonoua) and three planting densities (1 m × 80 cm, 1 m × 1 m, and 80 cm × 80 cm), resulting in six treatment combinations. Each block was divided into six sub-plots, randomly assigned one of the six treatments, for a total of 18 plots across the experiment, covering a total surface area of 1200 m². A two-meter buffer was maintained between plots to reduce inter-plot interference. Randomization of treatments within each block was conducted using a computergenerated random sequence to ensure unbiased allocation. Blocking was used to control for spatial variability across the field.

The treatments applied in both sites (Sanpedro 2 and N'Guessankro) are combinations of density and cultivar. These treatments are as follows:

YD1: Yavo planted at a medium density (1 m x 80 cm),

YD2: Yavo planted at a low density (1 m x 1 m),

YD3: Yavo planted at a high density (80 cm x 80 cm),

BD1: Bonoua planted at a medium density (1m x 80 cm),

BD2: Bonoua planted at a low density (1m x 1m),

BD3: Bonoua planted at a high density (80 cm x 80 cm)

Land preparation was done manually by ploughing using a hoe. Preparation began with clearing. This was followed by burning and stump removal at each site. The elementary plots were marked out with ropes to indicate the planting points according to the different densities of the plants. In order to ensure good germination conditions, the cuttings were planted after a rain and at three spacings: 80 cm x 80 cm; 1 m x 80 m and 1 m x 1 m. Manual weeding of individual plots was carried out as part of plot maintenance. A total of three weeding operations were carried out, the first of which was for weed control two weeks after sowing. The second was done 1 month after sowing and the third 2 months after planting. There were no phytosanitary treatments applied to the plots. In this study, data were collected from 20 plants per variety at different densities. Parameters such as plant height, stem diameter, number of tubers, number of branches, tuber length, tuber diameter, tuber weight and yield were recorded.

2.3. Statistical analysis of data

Standard errors (SE) are provided for all treatment means in the data tables to reflect variability. Before conducting ANOVA, assumptions of normality and homogeneity of variances were tested using the Shapiro-Wilk and Levene's tests, respectively. Data that did not meet these assumptions were transformed as needed to meet the requirements of the analysis. Analysis of variance (ANOVA) was performed on the data collected. For each variable studied, the means were compared using a 3-factor analysis of variance (ANOVA 3), factoring in locality, density, and cultivar. Test significance was determined by comparing the probability (p) associated with the statistic at the threshold of $\alpha = 0.05$. Maps were generated using QGIS version 2.8 software. All data were processed using the statistical software STATISTICA version 7.1.

3. Results and discussion

3.1. Results

3.1.1. Effect of locality on cassava agronomic parameters

The impact of locality on cassava agronomic parameters is illustrated in Table 1. The data reveals that locality significantly influenced all recorded parameters ($p < 0.05$). Among the localities assessed, N'Guessankro demonstrated the most favorable conditions for cassava growth and yield, with production exceeding 26020 kg/ha, compared to 14016.13 kg/ha observed in San pedro 2.

3.1.2. Variety response on agronomic parameters

Table 2 indicates that variety significantly influenced 75% of the agronomic parameters of cassava ($p < 0.05$), including plant height, collar circumference, number of branches, tuber length, tuber weight, and overall yield. The variety Bonoua demonstrated the highest values for vegetative parameters, while Yavo exhibited the highest mean tuber weight and yield. Notably, a discrepancy of approximately 2 kilograms in mean tuber weight and 2 kg/ha in mean yield was observed when comparing Yavo with the Bonoua variety.

3.1.3. Effect of planting density on cassava agronomic parameters

The impact of cassava planting density on agronomic parameters is summarized in Table 3. The results indicate that the applied densities significantly affected cassava production ($p < 0.05$). The highest values across all recorded parameters were achieved at the low density (1 cm x 1 cm), while the high density (80 cm x 80 cm) produced the lowest values. Meanwhile, the medium density (80 cm x 1 m) exhibited intermediate averages. Notably, the yield at the low

density surpassed the high density by over 13 kg/ha and exceeded the medium density by 3 kg/ha.

3.1.4. Effect of location * density interaction on cassava agronomic parameters

The results presented in Table 4 demonstrate that location, variety, and planting density significantly influence cassava productivity ($p < 0.05$). These factors impacted over 87.5% of the measured parameters, including yield. Optimal cassava production was observed in the N'Guessankro locality with the Yavo variety planted at a low density (1 m x 1 m). Conversely, high-density planting in Sanpedro2 markedly reduced cassava productivity.

3.2. Discussion

3.2.1. Effect of locality on cassava agronomic parameters

The results demonstrate that cassava agronomic traits varied significantly between the two study sites, with superior performance observed in N'Guessankro. Two key factors may account for this disparity. First, the soil fertility in N'Guessankro is likely more favorable for cassava growth. Cassava is highly responsive to soil nutrient availability, particularly nitrogen, phosphorus, and potassium, which influence root and shoot development (Howeler et al., 2017; Trung et al., 2022). Soil analyses in similar Ivorian regions have shown that even moderate differences in organic matter or pH can significantly affect cassava yield potential (Gnahoua et al., 2016). Second, climatic conditions in N'Guessankro, such as rainfall distribution and solar radiation, may have been more conducive to plant development. Hounzinme et al. (2020) emphasized that cassava growth is sensitive to environmental stressors, especially water availability during the early growth phase. Therefore, N'Guessankro's edaphic and climatic profile likely allowed for more effective biomass accumulation and root bulking, resulting in higher productivity.

3.2.2. Variety response on agronomic parameters

Significant varietal differences were observed, indicating strong genotype $\times$ environment interactions. The Bonoua variety generally exhibited higher vegetative growth, but the Yavo variety achieved better root yields in N'Guessankro under wider spacing. This divergence suggests that genotypic plasticity plays a critical role in varietal performance under varying environmental and management conditions. Yavo's superior performance under wider spacing may be due to its root system architecture and photosynthetic efficiency, which are enhanced when interplant competition is reduced. Studies by Punyasu et al. (2025) and Ceballos et al. (2006, 2020, 2021) support this, noting that modern cassava genotypes with deep or spreading

root systems can better exploit soil resources when provided sufficient space, particularly in marginal or transitional agroecological zones.

3.2.3. Effect of sowing density on cassava production and yield

Sowing density significantly influenced yield-related traits. Wider spacings (1 m × 80 cm and 1 m × 1 m) led to better performance compared to closer spacing (80 cm × 80 cm). This is likely due to enhanced light interception, reduced canopy competition, and improved nutrient and water availability per plant, which are critical for root development (Gnamien et al., 2024; Sultana et al., 2023). Dense planting creates shading that hampers lower leaf photosynthesis and increases competition for belowground resources. These results confirm previous findings (Fondio et al., 2000; Assefa et al., 2022), emphasizing the importance of spacing as a management lever to optimize cassava productivity.

3.2.4. Effect of location × density interaction on cassava agronomic parameters

The interaction between location and density was significant, particularly for the Yavo variety in N'Guessankro at the 1 m × 1 m spacing, which yielded the best results. This suggests a synergistic effect between genotype adaptability, environmental suitability, and reduced interplant competition. Yavo may possess traits such as drought avoidance, efficient water use, or deeper rooting, which make it better suited to wider spacing in low-competition environments like those found in N'Guessankro. Hussain et al. (2023) reported that organic matter-rich soils amplify such genotype-specific advantages by improving nutrient uptake and soil structure, which could further explain Yavo's superior performance. These findings underscore the importance of location-specific agronomic recommendations, taking into account the combined effects of environment, variety, and management practices.

4. Conclusion and recommendations

This study highlights the complex interplay between planting density, varietal performance, and agroecological conditions in determining cassava productivity in the Vavoua region. The Yavo variety achieved the highest yield when planted at a 1 m × 1 m spacing in N'Guessankro, suggesting that both genotype-specific spacing and site conditions must be considered in optimizing cassava production.

Implications:

- For farmers: Adoption of Yavo with a 1 m × 1 m spacing in agroecological zones similar to N'Guessankro could significantly boost yields.
- For extension Services: Development of variety-specific planting guides and site-based recommendations should be integrated into farmer training and rural advisory programs.
- For policymakers: These findings can inform regional cassava development strategies, especially in national programs targeting food security and agricultural productivity.

Recommendations

- Incorporate optimized spacing for specific varieties into national cassava production guidelines.
- Support soil fertility mapping and varietal zoning to tailor interventions based on local conditions.
- Promote participatory trials with farmers to validate and refine these recommendations under real-world conditions.

Future research directions

- Evaluate the impact of different organic and inorganic fertilization regimes on varietal performance in N'Guessankro.
- Investigate the long-term soil health effects of varied planting densities.
- Explore genotype-specific root traits and their relationship to spacing efficiency under different stress environments.

5. Acknowledgements

The authors wish to express their sincere gratitude to the local communities for their cooperation, availability, and active participation throughout the course of this study. Their support was instrumental to the successful completion of the fieldwork.

5. References

- Adjei-Nsiah, S., Kuyper, T.W., Leeuwis, C., Abekoe, M., K. and Giller, K., E. (2007). Evaluating sustainable and profitable cropping sequences with cassava and four legume crops: effects on soil fertility and maize yields in the forest/savannah transitional agro-ecological zone of Ghana. *Field Crop Research*, 103(2), 87-97. <https://doi.org/10.1016/j.fcr.2007.05.001>.
- Assefa, Amelework & Bairu, Michael. (2022). Advances in genetic analysis and breeding of Cassava (*Manihot esculenta* Crantz): A Review. *Plants*. 11. 1617.

DOI: [10.3390/plants11121617](https://doi.org/10.3390/plants11121617).

- Bilong, E.G., Abossolo-Angue, M., Ajebesone, F.N., Anaba, B.D., Madong, B.A, Nomo, L.B. and Bilong P. (2022). Improving soil physical properties and cassava productivity through organic manures management in the southern Cameroon. *Heliyon*, 8(6), e09570. <https://doi.org/10.1016/j.heliyon.2022.e09570>.
- Boni, N., Okoma, M.P., Kouakou, A.M., Dibi, K.E.B., Zohouri, G.P., Essis, B.S., and Dansi, A.A. (2015). Morphological Characterization of Cassava (*Manihot esculenta* Crantz) Accessions Collected in the Centre-west, South-west and West of Côte d'Ivoire. *Greener Journal of Agricultural Sciences*. 4. DOI: [10.15580/GJAS.2014.6.050614224](https://doi.org/10.15580/GJAS.2014.6.050614224).
- Borku, A.W. Cassava (*Manihot esculenta* Crantz): its nutritional composition insights for future research and development in Ethiopia. *Discover Sustainability* 6, 404 (2025). <https://doi.org/10.1007/s43621-025-00996-2>.
- Ceballos, H., Sánchez, T., A., L., Chávez, C., Iglesias, D., Debouck, G., Mafla, J. and Tohme J. (2006). Variation in crude protein content in cassava (*Manihot esculenta* Crantz) roots. *Journal of Food Composition and Analysis*, 19(6,7), 589-593. <https://doi.org/10.1016/j.jfca.2005.11.001>.
- Ceballos H., Rojanaridpiched C., Phumichai C., Kittipadakul P., Iglesias C., Gracen V.E. (2020) Excellence in cassava breeding: perspectives for the future. *Crop Breeding Genetics and Genomics*, 2(2):e200008. DOI: <https://doi.org/10.20900/cbgg20200008>.
- Ceballos, H., et al. (2021). Fifty years of a public cassava breeding program: evolution of breeding objectives, methods, and decision-making processes. *Theoretical and Applied Genetics*, 134(11), 2335–2353. DOI: [10.1007/s00122-021-03852-9](https://doi.org/10.1007/s00122-021-03852-9).
- Fondio, L., Djidji, A., H., N'gbesso, M., F., D., P. and Koné D. (2013). Evaluation de neuf variétés de tomate (*Solanum lycopersicum* L.) par rapport au flétrissement bactérien et à la productivité. *International Journal of Biological and Chemical Sciences*, 7(3), 10781086. DOI: [10.4314/ijbcs.v7i3.15](https://doi.org/10.4314/ijbcs.v7i3.15).
- Fondio, L., Kouamé, C., Djidji, A. and Traoré D. (2000). Densité de semis, croissance et production de deux lignées de gombo (*Abelmoschus spp*) en Côte d'Ivoire. *Cahiers Agricultures*, 8, 413-415. <https://revues.cirad.fr/index.php/cahiersagricultures/article/view/30200>.
- Gnahoua, J. B. G., Ettien, D. J. B., N'zué, B., De Neve, S., & Boeckx, P. (2016). Assessment of low-input technologies to improve productivity of early harvested cassava in Côte d'Ivoire. *Agroecology and Sustainable Food Systems*, 40(9), 941–964. <https://doi.org/10.1080/21683565.2016.1209610>.

- Gnamien Y. G., Kouassi N. J., Nanti B. T. J-I., Kouame N., Ayolié K., and Kouadio Y. J. 2024. Influence of number of seedling per planting hole on niebe (*Vigna unguiculata* l. Walp) agronomic parameters cultivated in Daloa, Côte d'Ivoire. *International Journal of Zoology and Applied Biosciences*, 9(4), pp: 59-67.
<https://doi.org/10.55126/ijzab.2024.v09.i04.010>.
- Hillocks, R., J. (2002). Cassava in africa. In: Hillocks, R.J., Thresh, J.M., Bellotti, A.C. (Eds.), Cassava: Biology, Production and Utilization. CABI Publishing, Wallingford, UK, pp. 41–54. <https://doi.org/10.1079/9780851995243.0041>.
- Hounzinme, S., S., Chanhoun, C., S., L., Teka O. and Oumorou, M. (2020). Effets de la variabilité climatique sur le rendement de quelques cultures vivrières dans le nord-est du Benin. *European Scientific Journal*, 12(16), 1857-7881.
<https://doi.org/10.19044/esj.2020.v16n12p137>.
- Howeler, R.H. (2017). Cassava cultivation and soil productivity. DOI: [10.19103/AS.2016.0014.25](https://doi.org/10.19103/AS.2016.0014.25).
- Hussain, A., Bashir, H., Zafar, S., A., Rehman, R., S., Khalid, M., N., Awais, M., Sadiq, M., T. and Amjad, I. (2023). The importance of soil organic matter (SOM) on soil productivity and plant growth. *Biological and Agricultural Sciences Research Journal*, 2: 11. DOI: <https://doi.org/10.54112/basrj.v2023i1.11>.
- Kintché, K., Hauser, S., Mahungu, N., M., Ndonda, A., Lukombo, S., Nhamo, N., Uzokwe, V., N., E., Yomeni, M., Ngamitshara, J., Ekoko, B., Mbala, M., Akem, C., Pypers, P., Matungulu, K., P., Kehbila, A. and Vanlauwe, B. (2017). Cassava yield loss in farmer fields was mainly caused by low soil fertility and suboptimal management practices in two provinces of the Democratic Republic of Congo. *European Journal of Agronomy*, 89, 107-123. <https://doi.org/10.1016/j.eja.2017.06.011>.
- Montagnac, J., Davis, C. and Tanumihardjo, S. (2009). Nutritional value of cassava for use as a staple food and recent advances for improvement. *Comprehensive Reviews in Food Science and Food Safety*, 8, 181 - 194. DOI: [10.1111/j.1541-4337.2009.00077.x](https://doi.org/10.1111/j.1541-4337.2009.00077.x).
- Radhouane, L., Aissa, N. and Romdhane L. (2014). Effets d'un stress hydrique appliqué à différents stades de développement sur l'aspect quantitatif et qualitatif des chez un écotype autochtone de sorgho grain (*Sorghum bicolor*), semences. *Journal of Applied Biosciences*, 74, 6149 – 6156. DOI: [10.4314/jab.v74i1.6](https://doi.org/10.4314/jab.v74i1.6).
- Sultana F, Dev W, Xin M, Han Y, Feng L, Lei Y, Yang B, Wang G, Li X, Wang Z, Xing F, Xiong S and Li Y. 2023. Competition for light interception in different plant canopy

- characteristics of Diverse Cotton Cultivars. *Genes (Basel)*, 14(2): 364. DOI: <https://doi.org/10.3390/genes14020364>.
- Trung, N. T., Bui, V. L., Lesueur, D., Atieno, M., Nguyen, C. T., Sarobol, E., Wongkaew, A., and Nakasathien, S. (2022). Cassava-cowpea intercropping system for controlling soil erosion in the Northern mountainous areas of Vietnam. *Asia-Pacific Journal of Science and Technology*, 27(05), APST–27. <https://doi.org/10.14456/apst.2022.74>.
- Punyasu, N., Thaiprasit, J., Kalapanulak, S., Saithong, T., and Postma, J.A. (2025). Modeling cassava root system architecture and the underlying dynamics in shoot–root carbon allocation during the early storage root bulking stage. *Plant Soil* 507, 863–880. <https://doi.org/10.1007/s11104-024-06771-y>.
- Wang, G., Oh, I. S., Courtright, S., H. and Colbert, A., E. (2011). Transformational leadership and performance across criteria and levels: A meta-analytic review of 25 years of research. *Group & Organization Management*, 36(2), 223-270. <http://dx.doi.org/10.1177/1059601111401017>.

Table 1: Influence of locality on cassava agronomic parameters

Locality	H	Circ	NB	NT	TL	TD	TW	Y
Sp	2.88±21.28a	22±6.18a	1.52±0.66a	3.78±1.61a	23.77±6.27a	48.3±8.33a	1.69±1.23a	14016.13±10314.44a
NG	5.99±0.37b	34.44±8.73b	2.152±0.12b	4.85±2.64b	26.86±5.87b	62.48±11.26b	3.21±2.46b	26020±21112.09b
F	0.473	81.675	0.006	7.118	0.008	61.412	0.005	15.659
p	0.029	0.013	0.032	0.049	0.010	0.035	0.000	0.000

For each trait, values with the same letters in the column are statistically similar

H : plant height (cm) ; **Circ** : collar circumference (cm) ; **NB**: number of branches; **NT**: Number of tubers; **TL**: Tuber length; **TD**: Tuber diameter; **TW**: Tuber weight; **Y**: yield (kg/ha); **Sp** : Sanpedro2; **NG**: N'Guessankro; $p < 0.05$

Table 2I: Variety response on agronomic parameters

Variety	H	Circ	NB	NT	TL	TD	TW	Y
Bonoua	5.21±21.24b	26.13±9.48a	1.96±1.42b	4.31±2.65a	24.25±6.55a	55.46±12.33a	2.29±1.96a	18665.33±15296.98a
Yavo	2.65±0.569a	30.35±9.730b	1.7±1.08a	4.31±1.76a	26.37±5.74a	55.32±12.09a	4.60±2.20b	21370.8±19688.99b
F	0.869	5.77	0.805	0.234	3.544	0.004	0.654	0.706
p	0.035	0.018	0.037	1.246	0.042	0.95	0.020	0.040

For each trait, values with the same letters in the column are statistically similar

H : plant height (cm) ; **Circ** : collar circumference (cm) ; **NB**: number of branches; **NT**: Number of tubers; **TL**: Tuber length; **TD**: Tuber diameter; **TW**: Tuber weight; **Y**: yield (kg/ha); $p < 0.05$

Table 3. Influence of density on agronomic parameters

Density	H	Circ	NB	NT	TL	TD	TW	Y
Low	2.62±0.50a	29.46±10.02b	1.6±1.21a	4.93±2.49ab	29.33±5.55a	57.94±11.54a	3.1325±2.33b	25060±18656.75c
Medium	2.55±0.43a	27.27±8.09a	1.95±2.08ab	4.15±1.88ab	25.021±6.77a	55.771±10.17a	2.2175±1.94a	22175±19411.72b
High	4.64±0.34b	28±5.34a	1.95±1.45ab	3.87±2.23a	25.59±6.43a	52.45±14.11a	2.003±1.83a	12819.2±1113.03a
<i>F</i>	0.972	0.510	0.614	2.408	0.083	2.104	3.439	5.699
<i>p</i>	0.038	0.041	0.043	0.044	0.032	0.026	0.0354	0.004

For each trait, values with the same letters in the column are statistically similar

H : plant height (cm) ; **Circ** : collar circumference (cm) ; **NB**: number of branches; **NT**: Number of tubers; **TL**: Tuber length; **TD**: Tuber diameter; **TW**: Tuber weight; **Y**: yield (kg/ha); *p* < 0.05

Table 4: Influence of the interaction of location * density on the agronomic parameters of cassava

Locality	Variety	Density	H	Circ	NB	NT	TL	TD	TW	Y
Sp	Bonoua	Medium	1.94±0.35a	21.99±8.71b	1.8±0.7b	3.8±1.54ab	25.198±4.62c	47.083±5.29ab	1.89±0.97ab	15120±7760.41ab
Sp	Bonoua	Low	2.24±0.328b	20.06±3.33b	1.3±0.67ab	3.7±1.49ab	29.598±8.62cd	51.911±8.007b	1.58±0.88ab	15800±8854.38ab
Sp	Bonoua	High	1.81±2.29a	17.52±1.67a	2.5±0.97bc	2.2±1.31a	26.06±6.94c	42.897±3.46a	0.66±0.42a	4224±251.92a
Sp	Yavo	Medium	2.79±0.17b	29.72±3.87c	1±0.34a	5.1±1.28bc	30.026±4.8cd	55.361±7.34b	3.33±1.44d	26640±11562.7bc
Sp	Yavo	Low	2.254±0.4b	21.65±3.94ab	1.1±0.316a	3.5±1.17a	24.432±5.2ab	52.821±6.41b	1.43±0.71ab	14300±7196.45ab
Sp	Yavo	High	1.914±0.43a	21.08±6.02ab	1.4±0.52ab	4.4±1.57ab	25.307±5.6b	39.754±6.96a	1.25±0.95a	8012.8±6092.29ab
NG	Bonoua	Medium	2.965±0.29b	29.25±7.13c	1.8±1.87b	5.9±4.04c	21.97±6.53ab	64.43±11.33	3.76±3.01e	30080±24127.66
NG	Bonoua	Low	2.904±0.317b	31.59±4.74cd	1.5±0.52ab	5.2±2.09bc	19.66±3.23a	58.056±9.80bc	2.52±1.07bc	25200±10768.27bc
NG	Bonoua	High	3.101±0.404c	36.42±11.09d	2.9±2.33bc	5.1±3.03bc	23.064±4.13b	68.404±10.87e	3.37±2.38e	21568±15273.98ab
NG	Yavo	Low	2.79±0.37b	36.88±13.03d	1.8±1.31b	4.9±2.02bc	24.16±2.84b	64.923±11.33d	3.55±3.00d	28400±24068.79c
NG	Yavo	Medium	2.802±0.208b	35.81±6.32d	3.9±3.51bc	4.2±2.34b	26.39±5.41c	60.298±13.85c	3.34±3.35d	33400±33503d
NG	Yavo	High	3.398±0.30c	36.98±5.87d	1±3.8a	3.8±1.75ab	<u>27.94±8.29d</u>	<u>58.78±8.75bc</u>	<u>2.73±1.55bc</u>	<u>17472±11234bc</u>
<i>F</i>			0.889	11.047	3.556	2.189	2.699	9.644	3.01221	3.159
<i>p</i>			0.554	0.004	0.001	0.020	0.004	0.002	0.001548	0.001

For each trait, values with the same letters in the column are statistically similar

H : plant height (cm) ; **Circ** : collar circumference (cm) ; **NB**: number of branches; **NT**: Number of tubers; **TL**: Tuber length; **TD**: Tuber diameter; **TW**: Tuber weight; **Y**: yield (kg/ha); Sp : Sanpedro2 ; NG: N'Guessankro; *p* < 0.05