



African Journal of Biological Sciences



Exploration of unique microbial communities in oral and denture microbiomes

Meenakshi S^{1*}, Pavitra T², Raghunath N³, Sumana K⁴, Praveen Rai⁵, Ramith Ramu⁶

¹ Department of Prosthodontics, JSS dental college and Hospital, Mysuru – 570015, Karnataka, INDIA

² Department of Microbiology, Cauvery Institute of Health Sciences, Mysuru - 570028, Karnataka, INDIA

³ Department of Orthodontics, JSS Dental college and Hospital, Mysuru – 570015, Karnataka, INDIA

⁴ Department of Microbiology, JSS Academy of Higher Education & Research, Mysuru – 570015, Karnataka, INDIA

⁵ Nitte (Deemed to be University), Department of Infectious Diseases & Microbial Genomics, Nitte University Centre for Science Education and Research (NUCSER), Mangalore – 575018, Karnataka, INDIA

⁶ Department of Biotechnology and Bioinformatics, JSS Academy of Higher Education & Research, Mysuru – 570015, Karnataka, INDIA

* Corresponding Author:

Dr. Meenakshi S

Associate Professor, Department of Prosthodontics, JSS dental college and Hospital, Mysuru, Karnataka

Email: dr.meenakshis@jssuni.edu.in

ABSTRACT

Background: Microorganisms that have unique traits and play important roles in both health and disease can flourish in the human mouth because of its favorable environment. As science progresses, there is mounting evidence that the oral microbiome is important in conditions other than oral diseases. The oral cavity's ease of monitoring and manipulation of the microbiome opens up new possibilities for microbiome-based diagnostics and therapeutics.

Methods: An electronic search on PubMed, Google Scholar, Scopus for bibliographic and available literature was done using different databases by following keywords oral microbiome and oral bio film, oral hygiene, were used.

Results: This article examines the current state of research on the oral microbiome, briefly summarizing the most modern techniques for characterizing the microbiome, such as cutting edge omics, next-generation sequencing, microbiome databases, and areas where there are now gaps in our understanding.

Conclusions: The deregulation in the oral microbiome play a crucial role in oral polymicrobial diseases, the development and use of new diagnostic technologies emphasizes the early identification, effective treatment, prevention of dental and periodontal diseases.

Keywords: Oral microbiome, Oral artificial substitutes, Oral hygiene, Oral biofilm, Probiotics

BACKGROUND

Term "oral microbiome, "oralmicro biota ", "oral micro flora" refers to microbial population lives in the human oral cavity. Oral microbiome was first discovered by Dutchman Antony van Leeuwenhoek (Huang et al., 2011) did so using a microscope he had built. His observations of his own dental plaque in 1683, which he described as "tiny live animalcules," served as the foundation for the earliest written accounts of the existence of numerous microbes. Joshua Lederberg, a 1958 Nobel Prize winner, is credited with coining the term "microbiome" (Deo and Deshmukh, 2019) (Bacali et al., 2022). In the mouth, there is 700 different types of bacteria, fungus, viruses, and protozoan's (Lu et al., 2019). The oral cavity is a perfect complicated environment for the growth of complex many microorganisms because it has a variety of microbial habitats, including the teeth, buccal mucosa, soft and hard palate, and tongue. Any alteration in the organisms' ordinarily harmonious connection with the host could result in mouth infections (Khor et al., 2021).

TYPES OF ORAL MICROBIOME

A core microbiome and a variable microbiome can be used to categorise the human microbiome. All humans have a core microbiome that is made up of the predominate species that live at various locations throughout the body in a healthy environment. The unique lifestyle, phenotypic, and genotypic determinants, as well as the individual's own unique lifestyle, have all contributed to the evolution of the varied microbiome. Although people have comparable micro biota at different locations on their bodies, there are some distinctions between species and strains of bacteria that can make individual's microbiome as unique to them as their fingerprint (Zarco et al., 2012).

HOW AND WHEN ORAL MICROBIOME DEVELOP?

From birth, the oral micro biota is acquired mostly through vertical transmission from mother to the child. From the first meal onward, the mouth is regularly injected with microbes, and the process of acquiring resident oral micro flora starts (Deo et al., 2019) (Kilian et al., 2016). The tongue&cheek, palate, tooth surfaces, and gingival crevice are just a few of the diverse settings found in the oral cavity that are home to various bacteria. Host factors including tooth eruption or tooth loss, dental health, and hormonal changes in the host all have an impact on these micro flora (Wu et al., 2000). Also, as people age, the types and numbers of oral microbes alter, particularly during the initial stages of dentition development (Percival et al., 1991). In a child's mouth, the emergence of teeth produces new surface for microbial infection and constitutes a significant ecological event. The oral microbial environment is significantly altered when primary teeth are replaced by an adult dentition (Kilian et al., 2016). In the retrospective study by PaweB J. Zawadzki, et al. to explore the micro flora. The existence of diverse microorganisms from distinct families, species, and bacterial strains, protozoan's, and fungus was shown by microscopic studies of samples derived from oral mucosal swab, culture, and in vitro tests in the patient groups examined (Zawadzki et al., 2017).

BIOFILMS IN ORAL MICROBIOME

Bio films are extracellular microbial aggregations that are highly structured and frequently self-produced. Bacteria in biofilms interact with one another, lead a synergistic lifestyle, and exhibit a variety of distinctive traits that define them apart from free-living cells (Muhammad et al., 2020). A biofilm covering the oral structures made by a large number of bacteria that are impacted by their surface and composition composition The biofilm's self-produced extracellular matrix is

mostly made up of extracellular DNA. *S. salivarius* with *E. faecalis* The most significant species in dental caries, biofilm, and endodontic infection are *Fusobacterium nucleatum*, *Treponema spp.*, Body that contains forsythensis, *P. gingivalis*, and *Aggregatibacter actinomycetem comitans*. Bacteria living in biofilm compete fiercely with one another because they are a sophisticated and diverse civilization. Although the influence of the oral microbiome on depression and anxiety has not been widely studied, two recent studies have highlighted alterations in the salivary microbiome composition in relation to depressive-like symptoms. They found that the abundance of certain bacterial taxa, such as Spirochaetaceae, Actinomyces, Treponema, Fusobacterium, and Leptotrichia species, correlated with the severity of depressive and anxiety symptoms in adolescents (Simpson et al., 2020, Patil et al., 2022). For resources, binding, and the chance to survive, the bacterial species compete. Bacitracin production, the two most frequently utilized competitiveness mechanisms by bacteria are quorum sensing and hydrogen peroxide excretion. Bacteriocins. Certain bacteria in biofilms express bacteriocins, which are specialised or non-specialized proteins that can affect other bacteria. Bacteriocins differ from conventional antibiotic in that they often only affect strains of the generating species or strains that are similar to it (Burcham et al., 2020, Marsh et al., 2006).

Although fewer than 100 phlotypes are observed in a typical human, the species represented as adherent cells in biofilm communities within the oral cavity have been estimated to represent a varied population of more than 700 phlotypes, including bacterial and archaeal subdomains (McCleane et al., 2014). Commensal organisms make up the vast majority of oral microorganisms. Oral infections and occasionally systemic disorders can be brought on by harmful bacteria. It is difficult to manage disease-cause biofilms that include microorganisms (Berger et al., 2018).

PREDOMINANT MICROBIAL FLORA IN ORAL CAVITY

Protista	Fungi	Bacteria	Viruses
<ul style="list-style-type: none"> • <i>Trichomonas.</i> • <i>TenaxEntamoeba.</i> • <i>Gingivalis</i> 	<ul style="list-style-type: none"> • <i>Candida albicans</i> • <i>Candida glabrata</i> • <i>Candida spp</i> 	<p>Gram-positive bacteria strains</p> <ul style="list-style-type: none"> • <i>Enterococcus faecalis</i> • <i>Enterococcus faecium</i> • <i>Staphylococcus epidermidis</i> • <i>Staphylococcus aureus</i> • <i>Micrococcus luteus</i> <p>Gram-negative bacteria strains</p> <p><i>Enterobacteriaceae:</i></p> <p><i>Escherichia coli</i></p> <p><i>Klebsiellaoxytoca</i></p> <p><i>Klebsiellapneumoniae</i></p> <p><i>NonEnterobacteriaceae:</i></p> <p><i>Acinetobacterbaumann</i></p> <p><i>Pseudomonas aeruginosa</i></p>	<ul style="list-style-type: none"> • <i>Herpesvirus</i> • <i>Retroviruses</i> • <i>Coronaviruses,</i> • <i>poxviruses,</i> • <i>Norovirus,</i> • <i>human papillomavirus</i> • <i>, Epstein–Barr virus,</i> • <i>Herpes simplex virus,</i> • <i>Hepatitis C virus, and HIV</i>

Adapted references: (Lee et al., 2021, Foster et al., 2004, Rostamifar et al., 2021, Kumari et al., 2022)

THE TOOLS USE TO DEFINE ORAL MICROFLORA

SAMPLE COLLECTION:

The oral microbiome is receiving a lot of interest, and there are reliable, practical, and efficient sample techniques for isolating oral microorganisms that are mostly found in saliva, the supragingival area, the sub gingival sub mucosal region, affected root canal system, and mucosal surface. The collection, transportation, processing, and storage of samples are all parts of the oral microbiome sampling strategy (Lu et al., 2022).

MICROSCOPY

Microscopic investigations enable count or detection of physically distinct structures, such as various cell shapes or gram-stain responses, and can immediately provide the investigator with a great deal of knowledge about the physical structure of a material. A wide range of stain techniques can be employed to designate certain features of interest in the absence of visually discernible features. For many years, the taxonomic differentiation of bacteria has been based on frequently indefinable characteristics of their cells thanks to the employment of straightforward staining techniques like the gram-stain, acid fast stain, or capsule stain. Similar techniques are still effective today, although they are typically learned through error and trial and only allow for the differentiation of stark structural changes (Tripathi and Sapra, 2023).

CULTURE AND MICROSCOPY:

The separation of pure cultures to enable the investigation of individual taxa is a key step in the microbiological analysis process, which is deeply anchored in culture-based techniques. Culture-based analyses are still significant, as well as a pure culture still is necessary before a new taxon

can be granted a legitimate taxonomic name, despite the fact that molecular-based approaches are getting more and more sophisticated. The viable count, which is the most often used culture-based metric, is typically determined by serially diluting a bacterial solution and then plating it onto an agar growth medium. For some microbe groups, viable counts are available.

The goal has traditionally been to find and characterise cultivated area bacteria that were connected to specific oral illnesses, guided by Koch's postulates. Culture, however, is not always able to accurately depict the heterogeneity of a oral microbiome. Miller acknowledged his inability to cultivate every bacterium he saw in 1890 (Benn et al., 2018).

There are still a lot of uncultivated microorganism from the human mouth cavity. Bacteria and viruses from human oral cavity were cultured utilising novel techniques of Isolation of Prior Uncultivated Oral Bacteria, with an emphasis on anaerobic species, by M. V. Sizova et al. In vivo culturing to selectively enhance on organisms actively developing inside the mouth (the "mini-trap" method) is one of these breakthroughs.

To reduce the impact of fast-growing microbes, (ii) single-cell long-term cultivation was used, and (iii) adaptations of traditional enrichment procedures were made utilizing medium which had no sugar, including glucose.maintained tight anaerobic conditions in the majority of their cultivation studies in order to facilitate the growing of obligatory anaerobes.

According to their findings, mini traps enrichment (11%), only one cultivation (3%) and standard plating (1%), had the highest successful recovery rates per cell. The single-cell cultivation approach was used to create the taxonomically richest collection (Li et al., 2005). The separation and preservation in pure culture of 10 strains, including members of what are probably three new microbial taxa, that were earlier only recognized by their molecular fingerprints, is a significant outcome. they concluded that their mix of cutting-edge methods will probably help bridge the

divide among wild and domesticated species that originate from human mouths (Sizova et al., 2012). This DGGE-guided method could lead to the creation of novel medium of various complex microbial populations (Tian et al., 2010). Many bacterial species detected in biological specimens can't be grown in culture, which is the fundamental issue with conventional culture and culture-based analytical tools, rendering them unsuitable for research (Deo and Deshmukh, 2019).

CULTUREINDEPENDENT:

Before cultivation, many molecular techniques have been employed can physically reduce the quantity and variety of bacteria in mixed samples. With the development of culture-independent technologies, these include filtering techniques like density-gradient centrifuged and elutriation and extinction-dilution in which materials were diluted, preferably down to single cells, before being cultured in isolation (Vartoukian et al., 2010).

DNA MICROARRAYS:

FeiTeng mechanical lysis-based DNA extraction techniques out performed all other experimental protocols studied when it came to characterising the oral micro biome (Teng et al., 2018).

Teng F, Darveekaran Nair SS,et al. concluded that orally microbiota structure is mostly influenced by the type of DNA extraction used, whereas the influence of 16S rRNA variable region areas is only marginally significant. Enzymatic

POLYMERASE CHAIN REACTION (PCR)

Inside the human oral cavity, bio films are made up of a vast variety of microbes. By presenting PCR-generated 16S rDNA fragment that migrate at varying distances, indicating the variations in the base-pair, PCR-based denaturing gradients gel electrophoresis (PCR-DGGE) examines microbial diversity (Benn et al., 2018, Li et al., 2005, Patil et al., 2021)

NEXT-GENERATION SEQUENCING (NGS) TECHNOLOGY

Ya Zhang et al. evaluated 12 studies before reaching their findings. Prior to intervention, the most common genera were Porphyromonas, Treponema, Tannerella, & Prevotella, although Streptococcus and Actinomyces often expanded and were the top genera identified by NGS technology (Zhang et al., 2021). The NGS technology showed promise in monitoring and analysing the danger of probable laboratory contamination by being able to track the nucleic acids bacterial contamination via various sources in the lab. While analysing data and interpreting results, it is important to take the possibility of contamination from chemicals, residual DNA, and the environment into account. In order to comprehend the complicated oral microbial population in clinical samples, researchers have been able to efficiently gather vast volumes of DNA fingerprint data in a single instrument run thanks to NGS technology. For point-of-care testing, it is essential to develop quick, easy, and sensitive detection methods due to the growing clinical significance of oral infections. The biggest problem right now and a serious worldwide health emergency is the corona virus disease (COVID-19) pandemic (Patil et al., 2020, Pradeep et al., 2023, Patil et al., 2020, Patil and Ramu, 2021).

METAGENOMICS:

A technique for locating bacteria that are impossible to grow is metagenomic analysis. It also uncovers the genomic variety in microbes by applying the power for genomic studies to the full society for microorganisms & their functional structures through a review of metabolic pathway genes. Also, it provides information on using databases on protein coding sequences. Have between 120 gigabases and 1.5 terabases of sequence data each run, met genomics, the genetic

analysis of the microbiome, provides sequence alignment analysis for studying microbial diversity, population organisation, and functional activity (Huang et al., 2021).

THE HUMAN ORAL MICROBIAL IDENTIFICATION MICROARRAY (HOMIM)

Actinobacteria, Bacteroidetes, Chlamydiae, Chloroflexi, Euryarchaeota, Firmicutes, Fusobacteria, Proteobacteria, Spirochaetes, (Olsen et al., 2015, Byrne et al., 2022) and are a few of the more prevalent oral bacterial species that can be detected simultaneously with HOMIM. This analysis of Human oral microbiome (HOM) advances knowledge of the oral microbiome in healthy subjects (Caseli et al., 2020).

Standards strains, clinical frequent isolates (containing bacteria and fungi) were identified using molecular approaches in this investigation that involved a partial sequence of 16S rDNA or ITS2. Also, throughout the past three years in our lab, molecular techniques were used to identify isolates that had eluded conventional laboratory testing (Wade et al., 2013).

Bacterial and fungal isolates : A Laboratory Medicine Centre of Nanfang hospitals, a 2,200-bed tertiary-care facility connected with a university, provided the bacterial and fungal isolates. Infectious diseases caused by bacteria and fungi, including those caused by anaerobic bacteria, aerobes, Mycobacteria, Candida, Aspergillus, and other isolates, are frequently diagnosed at this laboratory using culture-based methods. All fungi were found in deep mycosis isolates. With the help of a MicroScanWalkAway 96 plus System and the BD Phoenix 100 Automatic Microbiological Systems, both manufactured by Siemens Healthcare Diagnostics Inc. in West Sacramento, California, all isolates were phenol typed according to conventional laboratory practises (Wade et al., 2013).

DNA extraction: Using the TaKaRaMiniBEST Universal Genomic DNA Extraction Kit Ver.5.0 (TaKaRa) and the Lysis Buffer for Microorganism to Direct PCR kit, nucleic acids were extracted in accordance with the manufacturer's instructions (TaKaRa Bio Corp., Tokyo, Japan). Colonies have to be obtained from fresh cultures of bacteria or fungi in Sabouraud's or blood plate media. Moreover, the lysis buffer's colony and hyphae concentrations should be within a certain range (visible turbidity). Direct PCR amplification of previously obtained DNA was performed (Martiz et al., 2023).

Ingredients utilised in PCR amplifications included: 1.25 U of ex Taq DNA polymerase (TaKaRa), 5 L of template, 5 L of 10 PCR buffer, 4 M of each primer stock solution, 4 mM of each dNTP, and 50 L of sterile distilled water. For the amplification, a Mastercycler® PCR System was employed (Eppendorf International, Hamburg, Germany). The 35 cycles of 94°C for 30 s, 55°C for 1 min, and 72°C for 2 min that made up the thermocycling conditions also included a final extension step of 72°C for 5 min. Gel electrophoresis (100 V through a 1.5% agarose gel with 0.5 TBE (Tris-borate-EDTA) running buffer) was used to check the PCR amplification products, which were then stained with ethidium bromide and analysed using the GelDoc XR Gel Documentation System (Bio-Rad, USA.). PCR amplicon sizes were only evaluated at the Beijing Genomics Institute using TaKaRa comparisons of molecular size markers, followed by purification and sequencing (Sanger capillary sequencing) (Shenzhen, China) (Wade et al, 2013).

WHAT ARE THE ROLE OF ORAL MICROBIOME IN HEALTH

The human body contains a unique microbiome that really is essential for maintaining health. These microbes give significant biological benefit to their host, such as promoting the

development of a healthy gut microbiome. Controls the saliva's ability to buffer (at high PH) to neutralized oral acids, the kind that lead to tooth erosion. the bacterial acid load that causes tooth decay and cavities by decreasing the quantity of pathogenic acid-producing microbes through competitive exclusion. aids in avoiding gum disease reduces gum inflammation, starts the digestive process, turns nitrates into nitrites, which is a vital chemical for lowering blood pressure, and prevents plaque prevents halitosis (Cancan et al., 2014, Khadri et al., 2024).

- Encouraging healthy gastrointestinal and immunological systems.
- Certain bacteria assist to nitrate metabolism, that helps promote healthy blood pressure, as well as digestive and metabolic activities that support a normal metabolism.
- Facilitating the procedure of using saliva to break down meals and convert food's nutrients into energy
- Circulating the mouth with ionic minerals that are carried by saliva
- Supporting tooth remineralisation
- Delivering oxygen to the soft tissues and gums
- Shielding us from hazardous environmental germs while preventing diseases
- Preventing inflammation and oxidative stress
- Removing trash from the mouth's outside (Cho et al., 2021).

HOW TO IMPROVE ORAL HEALTHY MICROBIOME?

Probiotics are "live microorganisms that, when administered in sufficient proportions, impart a the host benefits from," according to the WHO definition. By balancing the oral flora, probiotic strains aid in the management of gingivitis. Acidogenic probiotic bacteria like Lactobacilli, Streptococci, and *Bifidobacterium* (Kumari et al., 2023) are the most frequently used bacteria for probiotics because they release antimicrobial substances that have an inhibition activity against pathogens through co-aggregation, producers of toxic by-products, and competition for substrates (Martiz et al., 2023). Studies have looked into using the probiotic genera Lactobacillus and *Bifidobacterium*, which are often used in formulations for digestive health, to prevent dental caries (Huligere et al., 2023).

The oral microbiome is a term used to describe the bacteria that inhabit the oral cavity. It is, after the gut. When comparing to other bodily parts, they exhibit an amazing variety of expected protein actions. The human microbiome is composed of both the fundamental microbiome and a changeable microbiome. Everyone has the same basic microbiome, but because everyone lives differently and has various physical characteristics, everyone also has a unique variable microbiome. There are two sites inside the oral cavity wherein germs might invade: the hard and soft tissues that make up teeth, in addition to the oral mucosa. Microbes can flourish in the oral cavity.

The oral cavity and the nasopharyngeal regions that are linked with it provide an environment that is perfect for the growth of bacteria. The typical temperature of the mouth cavity is 37 degrees Celsius, and this temperature rarely shifts much. This provides bacteria with a favourable habitat in which to thrive. In addition, the pH of saliva is consistently between 6.5 and 7, which is the ideal range for the majority of bacterial species. It fulfils the role of the a

medium for transfer of nutrients to the microorganisms, in addition to maintaining the hydration of the bacteria (Priya and Deshmukh, 2019). In the human oral microbiome, the DPP-4 activity of *Prevotella* is regarded as a virulence factor that contributes to biofilm formation and the development of periodontal disease (Ftieta et al., 2017). Hence, DPP-4 could be inhibited through phytochemicals and nanoemulsions in combination with phytochemicals which would decrease the biofilm formation of *Prevotella* and improve the bioavailability of phytochemicals to modulate oral microbiome (Sajal et al., 2022, Mahadev et al., 2022).

SIGNS OF POOR ORAL MICROBIOME:

Oral diseases, including dental cavities and periodontal disease, are among the most common diseases in the world, and they impact practically all age groups and geographic regions [41] The bacteria that live in the mouth can create metabolites, which in turn can have an effect on the progression of a variety of oral disorders. Significant difficulties may be produced by strains of *Staphylococcus aureus* that also take part in the creation of biofilm; biodegradable polymer infections are most commonly caused by this kind of bacterium. The bacteria that live in the mouth can create metabolites, which in turn can have an effect on the progression of a variety of oral disorders. Significant difficulties may be produced by strains of *Staphylococcus aureus* such as MRSA that also take part in the creation of biofilm; biodegradable polymer infections are most commonly caused by this kind of bacterium (Kumar et al., 2022).

Excessive plaque just on teeth can leave a morning film on your tooth that really is thick, sticky, unpleasant, and off-white. poor breath, Gum recession and bleeding gums delicate teeth oral sores A fungal ailment known as oral candidiasis or oral thrush occurs when the candida fungus (often *C. albicans*) overgrows in the mouth. Cavities and tooth decay, endodontic infections (root

canals), and alveolar osteitis are all examples of gum disease (periodontitis) (dry socket), respiratory infections, heart illness, stroke, tonsillitis, and more (Griffen et al., 2019, Johansson et al., 2000).

ORAL MICROBIOME FOR SYSTEMIC DISEASES

- **Endocarditis.** Infections of the inner layer of your heart chambers or valves frequently arise when bacteria or even other organisms from another part of your body, such your mouth, travel through your bloodstream and stick to particular spots there (endocardium).
- **Cardiovascular disease.** Although while the connection between oral infections and cardiovascular disease, clogged arteries, and stroke is not fully understood, certain research have suggested a link (Wu et al., 2000).
- **Pneumonia.** Pneumonia as well as other respiratory illnesses can be brought on by specific oral cavity bacteria that can be drawn into your lungs.
- **Diabetes.** By reducing the body's capacity to fight against infection, diabetes raises a risk of gum disease. Gum disease appears to be more prevalent and severe in people with diabetes (Ramu et al., 2017, D'Auito et al., 2017).
- **HIV/AIDS.** HIV-positive individuals are especially susceptible to a number of oral microbial infections. In persons with HIV/AIDS, oral issues such uncomfortable mucosal sores are frequent (Griffen et al., 2019).

- **Osteoporosis.** This bone-weakening disorder is linked to both periodontal loss and tooth loss. There is a slight possibility that a few osteoporosis drugs could affect the jaw bones (Wade et al., 2013).
- **Alzheimer's disease.** When Alzheimer's disease advances, dental health is shown to deteriorate (Gao et al., 2020, Martiz et al., 2022).

Anorexia and bulimia, rheumatoid arthritis, certain malignancies, and Sjogren's syndrome, an immune system ailment that produces dry mouth, are additional conditions that may be connected to oral health.

ORAL FLORA ON ARTIFICIAL SUBSTITUTES

CHANGES IN ORAL FLORA BEFORE AND AFTER DENTURES INSERTION

Because implants are now being used widely to treat edentulism, it is becoming more and more important to study that oral flora of missing teeth people. The study by Saeed Abdul Latteef Abdul-Kareem, B, shows found there was no statistically significant difference in the types of microorganisms during the post-insertion period while the overall number of bacteria reduced. While *E. coli*, *Klebsiella*, and *Moraxella* (Branhamella) have been observed after one month of wearing dentures, the mouth of the newly edentulous patient had previously harboured *Neisseria* species, which disappeared after dentures were inserted. *Candida* and streptococci were two other microorganisms that reduced., *Diphtheroids*, *Veillonella*, and *Acinetobactor*, on the other hand, were thought to be a part of the typical flora of edentulous patients that remained unaffected by denture use. Even while different germs can colonise dentures, when worn for a brief amount of time and with appropriate oral hygiene, the number of microorganisms in the mouth cavity does not rise (Muhammad et al., 2020). As people aged, so did the isolation of various non-resident oral microorganisms, possible opportunist pathogens, and viability count and proportion of lactobacilli in saliva (yeasts and staphylococci). With age, there were also changes in the ratio of *Actinomyces spp.* predominance. R. S. Percival et al. examine the level of the inherent and specialised host defences between healthy persons in the various age groups outlined here and patients of the same age who have different disorders. In some people, changes to the host defences' effectiveness may disturb the stability of a native oral microbiota and increase the possibility of colonisation by possibly hazardous species (Percival et al., 1991). In addition, it was shown that the oral microbiomes of young men and women differed (Burcham et

Meenakshi S / Afr.J.Bio.Sc. 6(5) (2024). 5111-5135

al., 2020). Extraction of teeth unavoidably changes the oral microbiota, and edentulous people have significantly lower microbial diversity, albeit some species can recover when wearing dentures (Marsh et al., 2006).

Salivary substitutes can be used continuously and are sometimes beneficial for senior citizens and radiation patients. It's probable that this ecological change will have an impact on the oral flora (Johansson et al., 2000).

Bridges-Porcelain is typically glued to precious metal to create bridges. For added strength, different non-precious metals are occasionally utilised in the base. There are also modern bridges that are fully constructed of a unique, durable porcelain.

Understanding the makeup, the oral flora and the bacteria that make it up has been studied. The microbes are typically found on the surface tissues of all humans, such as the mouth cavity. These microorganisms come in different numbers and types according on an individual's age, food, and level of personal hygiene (Deo and Deshmukh, 2019).

Many systemic illnesses, including bacterial endocarditis, pulmonary pneumonia, paediatric osteomyelitis, premature low birth weight, & cardiovascular disease, and diabetes mellitus are brought on by these oral bacteria (Li et al., 2000, Ramu et al., 2014).

To compete inside the ecosystem, one bacterium uses a lengthy chain of peptides called bacteriocin, which is synthesised by bacterial ribosomes. The oral community produces bacteriocins, which contribute to the diversity and ecological appropriateness of bacteria. Many species of naturally occurring oral cavity occupants create bacteriocin through quorum sensing, which controls the development of oral flora (Yang et al., 2014).

Gram-positive bacteria make bacteriocin, which is a shorter substance with 60 amino acids but a broad spectrum of activity among the gram positive bacteria. Aiming to define the production of bacteriocin while taking into account the aforementioned information, this current study were created to investigate & compared the bacteria fauna of the both healthy and unhealthy dental samples (Jack et al., 1995, Ramu et al., 2015).

CONCLUSION:

Understanding the diversity of the oral microbiome helps improve health and control systemic and oral disorders. The development of more specialised methods for treating related disorders has been substantially hastened using improved technical tools. These technologies have greatly accelerated the ability to detect the oral microbiome in a variety of samples taken from of the oral cavity.

REFERENCES

1. Huang, R., Li, M., & Gregory, R. L. (2011). Bacterial interactions in dental biofilm. *Virulence*. 2(5), 435-444.
2. Deo, P. N., & Deshmukh, R. (2019). Oral microbiome: Unveiling the fundamentals. *Journal of oral and maxillofacial pathology*. 23(1), 122-128.
3. Bacali, C., Vulturar, R., Buduru, S., Cozma, A., Fodor, A., Chiş, A., ... & Moldovan, M. L. (2022). Oral microbiome: getting to know and befriend neighbors, a biological approach. *Biomedicines*. 10(3), 671.
4. Lu, M., Xuan, S., & Wang, Z. (2019). Oral microbiota: A new view of body health. *Food Science and Human Wellness*. 8(1), 8-15.
5. Khor, B., Snow, M., Herrman, E., Ray, N., Mansukhani, K., Patel, K. A., ... & Machida, C. A. (2021). Interconnections between the oral and gut microbiomes: reversal of microbial dysbiosis and the balance between systemic health and disease. *Microorganisms*. 9(3), 496.
6. Zarco, M. F., Vess, T. J., & Ginsburg, G. S. (2012). The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral diseases*. 18(2), 109-120.
7. Kilian, M., Chapple, I. L. C., Hannig, M., Marsh, P. D., Meuric, V., Pedersen, A. M. L., ... & Zaura, E. (2016). The oral microbiome—an update for oral healthcare professionals. *British dental journal*. 221(10), 657-666.
8. Wu, T., Trevisan, M., Genco, R. J., Dorn, J. P., Falkner, K. L., & Sempos, C. T. (2000). Periodontal disease and risk of cerebrovascular disease: the first national health and nutrition examination survey and its follow-up study. *Archives of internal medicine*. 160(18), 2749-2755.
9. Percival, R. S., Challacombe, S. J., & Marsh, P. D. (1991). Age-related microbiological changes in the salivary and plaque microflora of healthy adults. *Journal of medical microbiology*. 35(1), 5-11.
10. Zawadzki, P. J., Perkowski, K., Padzik, M., Mierzwińska-Nastalska, E., Szaflik, J. P., Conn, D. B., & Chomicz, L. (2017). Examination of oral microbiota diversity in adults and older adults as an approach to prevent spread of risk factors for human infections. *BioMed research international*. 2017.
11. Muhammad, M. H., Idris, A. L., Fan, X., Guo, Y., Yu, Y., Jin, X., ... & Huang, T. (2020). Beyond risk: bacterial biofilms and their regulating approaches. *Frontiers in microbiology*. 11, 928.

12. Simpson, C. A., Adler, C., du Plessis, M. R., Landau, E. R., Dashper, S. G., Reynolds, E. C., ... & Simmons, J. G. (2020). Oral microbiome composition, but not diversity, is associated with adolescent anxiety and depression symptoms. *Physiology & Behavior*. 226, 113126.
13. Patil, S. M., Al-Mutairi, K. A., Firdose, N., Ramu, R., Martiz, R. M., & Ashwini, P. (2022). Pharmacoinformatics based screening discovers swertianolin from *Lavandula angustifolia* as a novel neuromodulator targeting epilepsy, depression, and anxiety. *South African Journal of Botany*. 149, 712-730.
14. Burcham, Z. M., Garneau, N. L., Comstock, S. S., Tucker, R. M., Knight, R., & Metcalf, J. L. (2020). Patterns of oral microbiota diversity in adults and children: a crowdsourced population study. *Scientific reports*. 10(1), 2133.
15. Marsh, P. D., & Percival, R. S. (2006). The oral microflora—friend or foe? Can we decide?. *International dental journal*. 56, 233-239.
16. McLean, J. S. (2014). Advancements toward a systems level understanding of the human oral microbiome. *Frontiers in cellular and infection microbiology*, 4, 98.
17. Berger, D., Rakhamimova, A., Pollack, A., & Loewy, Z. (2018). Oral biofilms: development, control, and analysis. *High-throughput*. 7(3), 24.
18. Lee, Y. H., Chung, S. W., Auh, Q. S., Hong, S. J., Lee, Y. A., Jung, J., ... & Hong, J. Y. (2021). Progress in oral microbiome related to oral and systemic diseases: an update. *Diagnostics*. 11(7), 1283.
19. Foster, J. S., & Kolenbrander, P. E. (2004). Development of a multispecies oral bacterial community in a saliva-conditioned flow cell. *Applied and environmental microbiology*. 70(7), 4340-4348.
20. Rostamifar, S., Azad, A., Bazrafkan, A., Modaresi, F., Atashpour, S., & Jahromi, Z. K. (2021). New strategy of reducing biofilm forming bacteria in oral cavity by bismuth nanoparticles. *BioMed Research International*. 2021.
21. Kumari, V. C., Huligere, S. S., Ramu, R., Naik Bajpe, S., Sreenivasa, M. Y., Silina, E., ... & Achar, R. R. (2022). Evaluation of probiotic and antidiabetic attributes of lactobacillus strains isolated from fermented beetroot. *Frontiers in Microbiology*. 13, 911243.
21. Lu, H., Zou, P., Zhang, Y., Zhang, Q., Chen, Z., & Chen, F. (2022). The sampling strategy of oral microbiome. *iMeta*. 1(2), e23.
22. Tripathi, N., & Sapra, A. (2020). Gram Staining.[Updated 2020 Aug 28]. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.

23. Benn, A. M. L., Heng, N. C. K., Broadbent, J. M., & Thomson, W. M. (2018). Studying the human oral microbiome: challenges and the evolution of solutions. *Australian dental journal*. 63(1), 14-24.
24. Li, Y., Ku, C. Y. S., Xu, J., Saxena, D., & Caufield, P. W. (2005). Survey of oral microbial diversity using PCR-based denaturing gradient gel electrophoresis. *Journal of dental research*. 84(6), 559-564.
25. Sizova, M. V., Hohmann, T., Hazen, A., Paster, B. J., Halem, S. R., Murphy, C. M., ... & Epstein, S. (2012). New approaches for isolation of previously uncultivated oral bacteria. *Applied and environmental microbiology*. 78(1), 194-203.
26. Tian, Y., He, X., Torralba, M., Yooseph, S., Nelson, K. E., Lux, R., ... & Shi, W. (2010). Using DGGE profiling to develop a novel culture medium suitable for oral microbial communities. *Molecular oral microbiology*. 25(5), 357-367.
27. Vartoukian, S. R., Palmer, R. M., & Wade, W. G. (2010). Strategies for culture of 'unculturable' bacteria. *FEMS microbiology letters*. 309(1), 1-7.
28. Teng, F., Darveekaran Nair, S. S., Zhu, P., Li, S., Huang, S., Li, X., ... & Yang, F. (2018). Impact of DNA extraction method and targeted 16S-rRNA hypervariable region on oral microbiota profiling. *Scientific reports*. 8(1), 16321.
29. Patil, S. M., Kumari, V. C., Sumana, K., Sujay, S., Tejaswini, M., Shirahatti, P. S., & Ramu, R. (2021). Sustainable development of plant tissue culture industry: The Indian scenario. *Journal of Applied Biology and Biotechnology*. 9(2), 18-27.
30. Zhang, Y., Qi, Y., Lo, E. C., McGrath, C., Mei, M. L., & Dai, R. (2021). Using next - generation sequencing to detect oral microbiome change following periodontal interventions: A systematic review. *Oral diseases*. 27(5), 1073-1089.
31. Patil, S. M., Kumari, V. C., Shirahatti, P. S., Sujay, S., Tejaswini, M., Ranganath, L. V., ... & Ramu, R. (2020). COVID-19 infection: The prospects of pharmacotherapy. *International Journal of Health & Allied Science*. 9(5), 111-113.
32. Pradeep, S., Patil, S. M., Dharmashekara, C., Jain, A., Ramu, R., Shirahatti, P. S., ... & Glossman-Mitnik, D. (2023). Molecular insights into the in silico discovery of corilagin from *Terminalia chebula* as a potential dual inhibitor of SARS-CoV-2 structural proteins. *Journal of Biomolecular Structure and Dynamics*. 41(20), 10869-10884.
33. Patil, S. M., VB, C. K., Shirahatti, P. S., Sa, S., Tejaswini, M., & Mallikarjunaswamy, C. (2020). Pharmacotherapy of COVID-19: A perspective of pathogenicity and life cycle. *Biomedical and Pharmacology Journal*. 13(3), 1579-1594.
34. Patil, S. M., & Ramu, R. (2020). The COVID-19 vaccine saga: A perspective. *Journal of Research in Pharmacy Practice*, 9(4), 218-219.

35. Huang, Y., Zhao, X., Cui, L., & Huang, S. (2021). Metagenomic and metatranscriptomic insight into oral biofilms in periodontitis and related systemic diseases. *Frontiers in microbiology*. 12, 728585.
36. Ramu, R. A. M. I. T. H., Shirahatti, P. S., Zameer, F. A. R. H. A. N., Lakkapa, D. B., & Nagendra, M. (2015). Evaluation of Banana (*Musa* sp. var. Nanjangud Rasa bale) flower and pseudostem extracts on antimicrobial, cytotoxicity and thrombolytic activities. *Int J Pharm Pharm Sci*. 7(1), 136-140.
37. Olsen, I. (2016). The oral microbiome in health and disease. *Oral Infections and General Health: From Molecule to Chairside*. 97-114.
38. Byrne, S. J., Chang, D., Adams, G. G., Butler, C. A., Reynolds, E. C., Darby, I. B., & Dashper, S. G. (2022). Microbiome profiles of non-responding and responding paired periodontitis sites within the same participants following non-surgical treatment. *Journal of Oral Microbiology*. 14(1), 2043595.
39. Caselli, E., Fabbri, C., D'Accolti, M., Soffritti, I., Bassi, C., Mazzacane, S., & Franchi, M. (2020). Defining the oral microbiome by whole-genome sequencing and resistome analysis: the complexity of the healthy picture. *BMC microbiology*. 20, 1-19.
40. Wade, W. G. (2013). The oral microbiome in health and disease. *Pharmacological research*. 69(1), 137-143
41. Martiz, R. M., Kumari VB, C., Huligere, S. S., Khan, M. S., Alafaleq, N. O., Ahmad, S., ... & Ramu, R. (2023). Inhibition of carbohydrate hydrolyzing enzymes by a potential probiotic *Levilactobacillus brevis* RAMULAB49 isolated from fermented *Ananas comosus*. *Frontiers in Microbiology*. 14, 1190105.
42. Cheng, C., Sun, J., Zheng, F., Wu, K., & Rui, Y. (2014). Molecular identification of clinical "difficult-to-identify" microbes from sequencing 16S ribosomal DNA and internal transcribed spacer 2. *Annals of clinical microbiology and antimicrobials*. 13, 1-7.
43. Khadri, M. N., Ramu, R., Simha, N. A., & Khanum, S. A. (2024). Synthesis, molecular docking, analgesic, anti-inflammatory, and ulcerogenic evaluation of thiophene-pyrazole candidates as COX, 5-LOX, and TNF- α inhibitors. *Inflammopharmacology*. 32(1), 693-713.
44. Cho, Y. D., Kim, K. H., Lee, Y. M., Ku, Y., & Seol, Y. J. (2021). Oral microbiome and host health: review on current advances in genome-wide analysis. *Applied sciences*. 11(9), 4050.
45. Kumari VB, C., Huligere, S. S., Alotaibi, G., Al Mouslem, A. K., Bahauddin, A. A., Shivanandappa, T. B., & Ramu, R. (2023). Antidiabetic activity of potential probiotics *limosilactobacillus* spp., *levilactobacillus* spp., and *lacticaseibacillus* spp. isolated from

- fermented sugarcane juice: a comprehensive in vitro and in silico study. *Nutrients*. 15(8), 1882.
46. Huligere, S. S., Kumari VB, C., Desai, S. M., Wong, L. S., Firdose, N., & Ramu, R. (2023). Investigating the antidiabetic efficacy of dairy-derived *Lacticaseibacillus paracasei* probiotic strains: modulating α -amylase and α -glucosidase enzyme functions. *Frontiers in microbiology*. 14, 1288487.
 47. Sajal, H., Patil, S. M., Raj, R., Shbeer, A. M., Ageel, M., & Ramu, R. (2022). Computer-aided screening of phytoconstituents from *Ocimum tenuiflorum* against diabetes mellitus targeting DPP4 inhibition: A combination of molecular docking, molecular dynamics, and pharmacokinetics approaches. *Molecules*. 27(16), 5133.
 48. Mahadev, M., Nandini, H. S., Ramu, R., Gowda, D. V., Almarhoon, Z. M., Al-Ghorbani, M., & Mabkhot, Y. N. (2022). Fabrication and evaluation of quercetin nanoemulsion: A delivery system with improved bioavailability and therapeutic efficacy in diabetes mellitus. *Pharmaceuticals*. 15(1), 70.
 49. Kumar, V., Shetty, P., HS, A., Chandra K, S., Ramu, R., Patil, S. M., ... & Poojary, B. (2022). Potential Fluorinated Anti - MRSA Thiazolidinone Derivatives with Antibacterial, Antitubercular Activity and Molecular Docking Studies. *Chemistry & Biodiversity*. 19(2), e202100532.
 50. Ramu, R., Shirahatti, P. S., Dhanabal, S. P., Zameer, F., Dhananjaya, B. L., & Prasad, M. N. (2017). Investigation of Antihyperglycaemic Activity of Banana (*Musa sp.* Var. Nanjangud rasa bale) Flower in Normal and Diabetic Rats. *Pharmacognosy magazine*. 13(Suppl 3), S417-S423.
 51. Martiz, R. M., Patil, S. M., Ramu, R., MK, J., P, A., Ranganatha, L. V., ... & Achar, R. R. (2022). Discovery of novel benzophenone integrated derivatives as anti-Alzheimer's agents targeting presenilin-1 and presenilin-2 inhibition: A computational approach. *PLoS One*. 17(4), e0265022.
 52. Ramu, R., Shirahatti, P. S., Zameer, F., Ranganatha, L. V., & Prasad, M. N. (2014). Inhibitory effect of banana (*Musa sp.* var. Nanjangud rasa bale) flower extract and its constituents Umbelliferone and Lupeol on α -glucosidase, aldose reductase and glycation at multiple stages. *South African Journal of Botany*. 95, 54-63.
 53. Ramu, R., Shirahatti, P. S., Devi, A. T., Prasad, A., Kumuda, J., Lochana, M. S., ... & Prasad, N. (2015). Bacteriocins and their applications in food preservation. *Critical reviews in food science and nutrition*. 0.
 40. health benefits of having a probiotic oral microbiome, <https://leemingdental.com.au/10-health-benefits-of-having-a-probiotic-oral-microbiome/>

41. Current Methods for Studying the Human Microbiome. Environmental Chemicals, the Human Microbiome, and Health Risk: A Research Strategy. Environmental Chemicals, the Human Microbiome, and Health Risk: A Research Strategy. Washington (DC): National Academies Press (US); 2017 Dec 29. 4, Current Methods for Studying the Human Microbiome. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK481559/>
42. Zarco, M. F., Vess, T. J., & Ginsburg, G. S. (2012). The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral diseases*. 18(2), 109-120.
44. Zhang, Z., Yu, W., Li, G., He, Y., Shi, Z., Wu, J., ... & Gao, Z. (2022). Characteristics of oral microbiome of healthcare workers in different clinical scenarios: a cross-sectional analysis. *BMC oral health*. 22(1), 481.
50. D'Aiuto, F., Gable, D., Syed, Z., Allen, Y., Wanyonyi, K. L., White, S., & Gallagher, J. E. (2017). Evidence summary: The relationship between oral diseases and diabetes. *British dental journal*. 222(12), 944-948.
52. Griffen, A. L., Thompson, Z. A., Beall, C. J., Lilly, E. A., Granada, C., Treas, K. D., ... & Fidel Jr, P. L. (2019). Significant effect of HIV/HAART on oral microbiota using multivariate analysis. *Scientific reports*. 9(1), 19946.
53. Gao, S. S., Chu, C. H., & Young, F. Y. F. (2020). Oral health and care for elderly people with Alzheimer's disease. *International journal of environmental research and public health*, 17(16), 5713.
54. Johansson, G., Andersson, G., Attstöm, R., & Edwardsson, S. (2000). Oral mucous membrane flora in patients using saliva substitutes. *Gerodontology*, 17(2), 87-90.
56. Bridges and partial dentures, Bridges and partial dentures - Oral Health Foundation (dentalhealth.org)
57. Alghamdi, S. (2022). Isolation and identification of the oral bacteria and their characterization for bacteriocin production in the oral cavity. *Saudi journal of biological sciences*. 29(1), 318-323.