

<https://doi.org/10.33472/AFJBS.6.6.2024.6191-6197>



African Journal of Biological Sciences

Journal homepage: <http://www.afjbs.com>



Research Paper

Open Access

Gait Impediment: A Rare Ca(U)Se

Dr. Thakur Keerthi¹, Dr. K Smeeta Merlin², Ravanagomagan^{3*}

¹Post Graduate, Department of Paediatrics, Sree Balaji Medical College and Hospital

²Post Graduate, Department of Paediatrics, Sree Balaji Medical College and Hospital

^{3*}Associate Professor, Department of Paediatrics, Sree Balaji Medical College and Hospital

Corresponding Author: ^{3*}Ravanagomagan
Chennai-600044

Article Info

Volume 6, Issue 6, June 2024

Received: 21 April 2024

Accepted: 31 May 2024

Published: 25 June 2024

doi: [10.33472/AFJBS.6.6.2024.6191-6197](https://doi.org/10.33472/AFJBS.6.6.2024.6191-6197)

ABSTRACT:

12year old girl, second born of second degree consanguineous marriage, with normal birth history presented with gradual onset of difficulty in walking since two years, currently being unable to walk without support. No other domains of development were affected. She was on antiepileptics from 3 to 6 years of age. On examination, B/L Ptosis was present with hypertelorism, tower skull, strabismus, microcephaly and generalized muscle wasting. CNS examination revealed hypertonia of both upper and lower limbs with brisk deep tendon jerks, right sided ankle clonus and extensor plantars. MRI Brain revealed large posterior fossa cyst communicating with slit like opening with fourth ventricle and hypoplasia of left cerebellar hemisphere. MRI C-Spine demonstrated anterior displacement of tip of odontoid process and severe compression of spinal cord, suggestive of Dandy Walker Malformation with CV Junction anomaly.

Keywords: Posterior Fossa Cyst, Left Cerebellar Hypoplasia, CV junction anomaly

© 2024 Dr. Thakur Keerthi, This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made

1. Case Report:

12 year old girl, second born of second degree consanguineous marriage, presented with gradual onset of difficulty in walking since 10 years of age, currently being unable to walk without support.No other domains of development were affected. She was currently studying in Class 7 and there was no history poor scholastic performance.Significant past history included use of antiepileptics (Tab. Phenytoin) from 3 - 6 years of age, given for Seizure disorder. Last known seizure was at 4 years of age.She was delivered by normal vaginal delivery, cried immediately at birth and there was no history of NICU admission. Her mother had a history of one previous IUD.

Her general examination revealed presence of bilateral ptosis, hypertelorism, tower skull strabismus,microcephaly, glossitis and pallor. It was observed that her mother had strabismus (Esotropia) too since childhood. The child's CNS examination demonstrated no abnormalities with regards to the higher functions. Her motor examination revealed hypertonia of the upper and lower limbs. Brisk deep tendon jerks and right sided ankle clonus were elicited along with upgoing plantar reflexbilaterally. The child was currently unable to walk without support. Her sensations were intact.

Primary work-up included complete blood profile and urine routine, which revealed no abnormalities. Her CPK levels were in the normal range. Ophthalmologist consultation was sought in view of ptosis and strabismus and the patient was advised spectacle usage.



Bilateral Ptosis and Strabismus



Examination of mother revealed
Strabismus (Esotropia).

Neurologist was consulted and further evaluation in terms of MRI Brain and MRI Cervical Spine were done. MRI Brain revealed large posterior fossa cyst communicating with slit like opening with fourth ventricle and hypoplasia of left cerebellar hemisphere. MRI C-Spine demonstrated anterior displacement of tip of odontoid process and severe compression of spinal cord with compressive myelopathy changes within the cord, suggestive of Dandy Walker Complex with CV Junction anomaly.

Neurosurgeon was consulted in view of craniovertebral junction anomaly, causing compressive myelopathy. The patient was advised decompression and stabilization surgery. However, patient's parents were not willing for surgery when the risk of quadriplegia was explained to them. The patient was advised physiotherapy and occupational therapy. She still continues to go to school and is performing well academically. She was advised to come for regular follow up and advised to use a cervical collar.

2. Discussion:

Dandy-Walker complex is a congenital group of disorders that have overlapping symptoms including Dandy-Walker malformation having a small cerebellar vermis, large fourth ventricle, and enlarged posterior fossa Isolated cerebellar vermis hypoplasia, Mega-cisterna magna, Posterior fossa arachnoid cyst. It is more common in females than males. The incidence is 1 in every 30,000 births. The longterm outcome and prognosis depend on the underlying malformation and symptoms it causes. Most of the cases of Dandy Walker are associated with hydrocephalus. However, this case was not associated with hydrocephalus. Mental retardation occurs in fewer than half, most often in those with severe hydrocephalus, chromosome abnormalities or other birth defects. Seizures occur in 15-30% of those affected. DWM results from defects in early embryonic development of the cerebellum and surrounding structures. A few patients have chromosome abnormalities including deletion of chromosome 3q24.3 (the location of the first DWM genes, known as ZIC1 and ZIC4), 6p25 or 13q32.2-q33.2, or duplication of 9p. In the remainder, it is probably due to other more complex genetic and perhaps environmental factors (teratogens) as the recurrence risk in siblings less than 5%. DWM may also occur as part of a genetic syndrome that includes


multiple birth defects, such as the PHACES syndrome of facial hemangioma, heart and sternal defects and DWM. Many other syndromes and chromosome abnormalities have been reported with DWM, but most of these appear to have CVH rather than typical DWM.


3. Conclusion:

A supportive team approach for children with Dandy-Waller malformation is often warranted and may include special education, physical therapy and other medical, social or vocational services. Genetic counselling is recommended for families that have a child with Dandy Walker malformation.



Patient standing supported against the wall







4. References:

1. Barkovich A, Kjos B, Norman D, et al. Revised classification of posterior fossa cysts and cystlike malformations based on the results of multiplanar MR imaging. *AJNR* 1989; 10:977-988.
2. Bordarier C, Aicardi J. Dandy-Walker syndrome and agenesis of the cerebellar vermis: diagnostic problems and genetic counseling. *Dev Med Child Neurol* 1990;32(4):285-94.
3. Ecker J Shipp T, Bromley B, et al. The sonographic diagnosis of Dandy-Walker and Dandy-Walker variant: associated findings and outcomes. *Prenat Diagn* 2000;20(4):328-32.
4. Grinberg I, Northrup H, Ardinger H, et al. Heterozygous deletion of the linked genes *ZIC1* and *ZIC4* is involved in Dandy-Walker malformation. *Nature Genetics* 2004;36:1053-1055.
5. Kalidasan V, Carroll T, Allcutt D. et al. The Dandy-Walker syndrome-a 10 year experience of its management and outcome. *Eur J Pediatr Surg* 1995;5 suppl 1:16-8.
6. Kolble N, Wisser J, Kurmanavicius J, et al. Dandy-Walker malformation: prenatal diagnosis and outcome. *Prenat Diagn* 2000;20(4):318-27.
7. Incesu L and Khosia A. Dandy-Walker Malformation. *eMedicine*; last updated 10/31/03.
8. McKusick VA ed. *Online Mendelian Inheritance in Man (OMIM)*. Baltimore, MD: The Johns Hopkins University; Entry No. 220200; Last updated 2/22/05.
9. Ecker J Shipp T, Bromley B, et al. The sonographic diagnosis of Dandy-Walker and Dandy-Walker variant: associated findings and outcomes. *Prenat Diagn* 2000;20(4):328-32.

10. Aldinger KA, Lehmann OJ, Hudgins L, Chizhikov VV, Bassuk AG, Ades LC, Krantz ID, Dobyns WB, Millen KJ. FOXC1 is required for normal cerebellar development and is a major contributor to chromosome 6p25.3 Dandy-Walker malformation. *Nat Genet.* 2009 Sep;41(9):1037-42. doi: 10.1038/ng.422. Epub 2009 Aug 9.