



African Journal of Biological Sciences



Research Paper

Open Access

Pollen grain efficiency of two male date palm (*Phoenix dactylifera* L.) genotypes with exogenous application of GA₃ and NAA; *in vitro* and field study

Mohammad S. Al-Tarawneh¹, Khalid M. Al-Absi², Muawya A. Al Asasfa³

¹Graduated student, Department of plant production, Faculty of Agriculture, Mutah University, Karak, Jordan

²Professor, Department of plant production, Faculty of Agriculture, Mutah University, Karak, Jordan

³Assistant Professor, Department of plant production, Faculty of Agriculture, Mutah University, Karak, Jordan

Corresponding author (*): Mohammad S. Al-Tarawneh

Article Info

Volume 6, Issue 8, April 2024

Received: 12 Feb 2024

Accepted: 19 March 2024

Published: 07 April 2024

Abstract

Two male date palm cultivars, "Ghanami akhdar" and "Jarvis," were studied in order to determine the effects of gibberellic acid (GA₃) and naphthalene acetic acid (NAA) on their pollen grain viability, germination, and pollen tube elongation. Different level of each GA₃ (10, 50 and 100 mg L⁻¹) and NAA (1, 2 and 3 mg L⁻¹) were used. The viability and fertility of pollen grains were tested by applying same levels to the aqueous pollen suspension using in date palm pollination in the field. The result showed that GA₃ at 100mg/L improved date palm pollen germination through accelerating germination rate and pollen tube growth which reflected on the speed of fertilization. Application of 100mg/L GA₃ upon pollination, also, led to the best results in term pollen tube growth and fruit set at Khelal stage; this may refer to a relationship between the amount of GA₃ in pollen grains and its role in fertilization, fruit growth and development. NAA at these (1, 2 and 3 mg L⁻¹) concentrations can present an effective thinning method in "Medjool". On other hand, no significant differences between cultivars were observed *in vitro*, but "Ghanami akhdar" outperformed "Jarvis" *in vivo*. "Ghanami akhdar" can be considering as a recommendable pollinator for "Medjool".

Key words: Date palm, Pollen germination, Pollen viability, Fruit set, Gibberellic Acid, Naphthalene Acetic Acid

© 2024 Mohammad S. Al-Tarawneh, this is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made

Introduction

The date palm is considered one of the most domesticated fruit trees in terms of moral value because of its religious importance for many different global societies, as well as for its health benefits and its productive capacity in semi-arid and arid environments (Jain and Johnson, 2015; Krueger, 2021). Date palm plants are categorized as dioecious fruit trees since their male and female blooms grow on different trees. Successful fruit set in fruits bearing seeds is contingent upon the effectiveness of pollination. A fruit's ability to form kernels and grow is dependent on

successful pollination, particularly in the case of nuts and drupe fruits with a single seed, such as stone fruits and dates. In date palms, pollination is the process by which pollen grains are transferred from male to female trees, either naturally or by manual or mechanical means. It is deemed sufficient when 60–80% of the female flowers are pollinated, and this often results in a decent fruit (Alasasfa, M., 2021). Pollination can adversely be affected by, temperature; humidity, pollen grain viability, and ability of female flowers to receipt pollen grain (Alasasfa, 2019). Khalifa et al, (1983), stated that, the big difference in fruit set from one year to another is mainly due to climatic conditions. Several studies on male cultivars have indicated their pollen efficiency. "Jarvis" is an American seedling male date palm that was recognized for its valuable qualities before 1954 and is now included of the Date Palm Germplasm Repository's holdings. Jarvis produces several inflorescences that give enormous amounts of pollen (Krueger, 2021). Pollen from "Jarvis" was found to be the best source for improving fruit physical characteristics. (Mohammadi et al., 2017). From its side "Ghanami akhdar" is a well-known Iraqi cultivar showed the highest pollen germination percentage (87%) and pollen tube growth (323 μ m) compared to other five cultivars (Aljibouri et al., 1990). In order to increase the efficiency of pollen grains, it was necessary to increase the factors that lead to the effectiveness of pollen grains, and among these studies was the introduction of hormones as a stimulating factor to increase the efficiency of pollen grains.

Gokbayrak and Engin (2017) discovered that GA₃ was the most effective plant growth regulator for enhancing pollen germination *in vitro*. Wu et al. (2008), proposed that GA₃ promotes pollen tube growth. According to Prajapati and Jain (2011), GA₃ might play different roles depending on the species tested and the dosage used; it can either help or inhibit *in vitro* pollen germination and tube development. Radovic et al. (2016) found that GAs caused an increase in pollen germination and pollen tube length for many cultivars of almonds. In addition, it was found that GA₃ was suitable for one of two examined species of Bauhinia with related to germination rate (Sanjay Kumar et al., 2016). In term of GA₃ dose, several studies indicated that the germination rate and pollen tube growth were directly stimulated by the high concentrations of GA₃; Qi et al. (2010) investigated the inducing effect of high gibberellin concentrations in kiwifruit, they found that 60-90mg/L GA₃ stimulated pollen germination and pollen tube growth, and 90mg/L was the best concentration. Using GA₃, at high concentrations, significantly improved the percentage of germination as well as pollen tube lengths of two almond (*Prunus dulcis*) cvs (Maita and Sotomayor, 2015). In Chaohong peach, the most suitable concentrations that stimulated pollen germination and tube growth of GA₃ were 25-100 ppm (Xue, et al, 2008), while the most suitable concentration of GA₃ for pollen germination and tube growth of *Syringa oblata* was 100mg/L (Liu, et al 2011). Other studies indicated an inverse relationship; low concentrations stimulate germination and that the concentrations altitude has an inhibitory effect on germination measurements in pomegranate (Gokbayrak and Engin (2018), pistachio (Acar and Sarpkaya, 2010), cotton (Almeida, 1990) and *Arbutus unedo* (Gökbayrak, et al 2020). According to Shenxi, et al, (2004), there were a negative relationship trend between pollen viability and the content of GA₃ and the mediate concentration of GA₃ increased germination and pollen tube growth.

On the other hand, auxin is a hormone that accumulates in pollen grains and governs several aspects of plant flora growth, including apical meristematic differentiation, stamen lengthening, anther maturation, and pollen grain development (Salinas-Grenet et al., 2018). However, a reduction in free auxin levels, particularly in pollen grains, can have a significant impact on pollen grain growth and elongation of the pollen tube, which affects reproduction and the rate of egg fertilization, and consequently seed yield (Gao et al., (2019). NAA is a type of synthetic auxin hormone that is commonly utilized for the vegetative propagation of plants through stem and cutting. The impact of NAA on plant development is significantly influenced by the concentration and timing of its application (Suman, et al., 2017).

The objectives of this study is to examine different level of GA₃ and NAA on germination percentage, germination rate, and pollen tube elongation of two male date palm cultivars "Ghanami akhdar" and "Jarvis", in addition to studying the viability and fertility of pollen grains by the direct method in the field, which is expressed through the fruit set and to test the metazenic effect of two pollen sources with regard to fruit set parameter.

Materials and Methods

Location

The field work for this research was performed during the year 2021 at the experimental date palm orchard in Ghor Alsafi Station, National Agriculture Research Center (NARC). The station is located at latitude 31° 02' N and longitude 35° 3E. It is about 380 meters below sea level. The soil texture is of the soil is sandy loam type. Furthermore, all laboratory works were carried out at the Department of Plant Production, Faculty of Agriculture, Mutah University, Jordan.

Pollen grains collection.

Mature spathes were collected from eight years old date palm males "Jarvis" and "Ghanami Akhdar". The selected male cultivars are traditionally used as pollinators of commercial date cultivars. Fresh mature pollen grains were collected and incubated at 25°C.

Germination medium preparation

Fourteen treatments were used including two control treatments for each cultivar. *In vitro* germination basal medium consisted of Calcium nitrate dihydrate (0.417 g), Boric acid, (0.200 g), Potassium Nitrate (0.101g), Magnesium sulfate, 7H₂O (0.217 g), agar (10 g) and sucrose (200g) per liter. The pH of the media was adjusted at 5.7. This media was sterilized in an autoclave at 121°C and 15 psi for 20 minutes. After sterilization, NAA at concentration of 1, 2 and 3 mg/L and GA₃ at concentration of 10, 50 and 100 mg/L were added to the media. All Petri dishes were incubated at 27 ± 2 °C for 1, 2 and 3 days.

Experimental set up

In vitro pollen viability, germination and pollen tube growth

Two factorial experiment using Randomized complete block design (RCBD) was carried out. The first involved two different sources of pollinators; "Jarvis" and "Ghanami Akhdar" as the first factor, whereas the second factor was the two plant hormones; NAA at concentration of 1, 2 and 3 mg/L and GA₃ at concentration of 10, 50 and 100 mg/L. Total fourteen treatments, each one contained four petri dishes (replicates), one slides per each petri dish was used as sampling unit at 24, 48 and 72 hours. Slides with germinated pollens were observed under 10× magnification objective of an Evos microscope paired with a Nikon DS – Ri1 camera. The photographs of the observed microscopic fields were taken at a resolution of 1920 × 1536 pixels. Five microscopic fields per slide were photographed.

Pollen viability was determined by staining pollen grain sample taken from each petri dish with 1% acetocarmine. Pollen grain from each microscopic field were counted and graded as viable and aborted in two classes. The pollen grains looking normal and stained red were considered viable, whereas poorly stained or colorless were record as nonviable. The pollen viability percentage was calculated by the following formula according to Iqbal et al. (2019).

$$\text{Pollen viability (PV \%)} = (\text{Viable pollen} / \text{Total pollen}) \times 100$$

Total number of pollen grain and the germinated pollen on each microscopic field were counted at 24, 48 and 72 hours. Pollen grain was considered as germinated if the length of pollen tube is larger than the diameter of the pollen grain. Pollen germination percentage was calculated by using the following formula:

$$\text{Pollen germination (PG \%)} = (\text{Germinated pollen} / \text{Total pollen}) \times 100$$

Absolute pollen viability or the effective germination capacity was calculated using the formula of Visser et al. (1977).

$$\text{Absolute pollen viability (APV \%)} = (\text{PV\%} \times \text{PG \%}) / 100$$

For determination of pollen grain tube length (PTL), 10 pollen tubes were measured from each microscopic field. The length of pollen tubes was measured with a scale of 0.61 μM/ pixel with the help of the software, AxioVision version 4.2. 8.

Fruit set of "Medjool" date palm

The same treatments examined *in vivo* were conducted at pollination time (March, 13, 2021) by using the suspension medium used in aqueous pollination of date palm tree which consist of 2 gm pollen + 3 gm sugar + 1L of distilled water. Fourteen uniform "Medjool" date palm trees, 20 years old were chosen to be pollinated. The fourteen treatments were applied on 42 bunches, 3 bunches (replicates) per treatment, 5 strands were marked to measure fruit set percentage at two different stages; initial fruit set 7 days after pollination (DAP) and fruit set at Khelal stage (July 11). The number of fruit and the number of flower scars were recorded and then the fruit set was expressed as the following equation:

$$\text{Fruit set \%} = \frac{\text{Number of fruit set in the strand}}{\text{Total number of flower per strand}} \times 100.$$

Statistical analysis.

The MSTATC software was utilized to conduct an analysis of variance (ANOVA) on the experimental data. During the ANOVA, the impacts of various treatments and their interactions were evaluated. To compare means (P=0.05), the Fisher's protected LSD test was employed.

Results and Discussion

Effect of pollen source (*in vitro*) on pollen viability, pollen germination (PG %), absolute pollen viability (APV %) and pollen tube length.

The effect of pollen source on viability (PV %), germination (PG %), absolute pollen viability (APV %) and pollen tube length (PTL) is shown in Fig.1. The results indicated were no significant differences between the two pollen sources; "Ghanami akhdar" and "Jarvis" in terms of PG% and APV% during the first 48 hours of incubation. but a slight significant superiority of "Jarvis" were noticed in PG and APV values at the third period which recorded 73.60% and 62.58%, respectively; however, "Ghanami akhdar" recorded 66.97% and 57.31% (Fig. 1.b and c). The other two study parameters (PV% and PTL) didn't show any significant differences between the two cultivars along the incubation period (Fig. 1.a and d). This close response of the two studied cultivars may be due to their high vigourity and fertility, as they are among the most widely used cultivars for pollinating female date palm trees in the largest dates producing countries.




Figure 1: Effect of two pollen sources; "Ghanami akhdar" and "Jarvis" on pollen viability (PV %) (a), pollen germination (PG %) (b), absolute pollen viability (APV %) (c) and pollen tube length (PTL (µm)) (d) at three incubation periods; 24, 48 and 72 hours

Effect of plant hormones (*in vitro*)

Effect of GA3 (*in vitro*).

Exogenous hormones effects in this study indicated that GA₃ at 100mg/L didn't present any inhibitory or enhancement effects at the end compared to control (Fig.2.a~d). However, it accelerated whole in vitro germination process in the early stage. The concentration of 100mg/L was a stimulus to germination, even if it was partially, and it itself was without effect also in other cases, especially in the last stages. There was also inhibition of the germination process by the concentrations of 10 and 50mg/L. Hence, the great importance of such experiments is determining the critical limit for the appropriate hormone concentration for a particular variety.

The use of concentrations 10 ~ 50 mg/L GA₃ led to emergence of an inhibitory role of GA₃ in the three germination characteristics (PG%, APV% and PTL) in "Ghanami akhdar". The level of 50 mg/L was critical in some cases in "Jarvis". Application of 100 mg/L showed a high effectiveness in accelerating the germination rate and growth rate of the pollen tube in the first 24 hours and did not have any inhibitory role in the later periods (Fig.2.a~d). The germination percentage after applying 100mg/L GA₃ reached 66.88% compared to 43.77% in the control treatment during the first 24 hours of incubation (Table 1); and the absolute viability rate reached 59.59% compared to 39.91% (Table 2), while the length of the pollen tube reached 232.5µm compared to 133.9 µm (Table 2). It could be concluded that this treatment increased the speed of germination by 50% and the speed of growth of the pollen tubes by 75%.

According to the findings of Samiee Rad et al. (2016), the impact of growth regulators on the percentage of pollen germination was found to be inconsistent and influenced by the concentration of the material and the compounds present in the medium. The presence of GA₃ and NAA without boric acid significantly hindered the germination of pollen. An ideal concentration for one genotype may be toxic for another genotype, and vice versa. According to Patel and Mankad (2014), plant growth regulators may contribute to the growth promoting effects that enhance early emergence of pollen tubes. This leads us to wonder about what would have happened if the GA₃ concentration rose to a higher level than 100mg/L? At least in this study, the concentration of 100mg/L can be considered as the lowest limit that can be built upon in the future with regard to date palm pollen. The higher concentration may lead to an increase in the germination rate and the length of the pollen tube at a constant rate, not only in the early stages of germination. Other factor could be vital; the relationship between medium components and applied hormones. It is known almost that pollen germination mediums were tested on the base of free plant hormones; application of plant hormones may lead to unbalance.

Effect of NAA (in vitro).

In general, it can be concluded from the results that all concentrations of NAA used in this study led to the inhibition of the germination process compared to the control, whether through the germination percentage and rate or through pollen tube growth (Fig.2.a~d). The absence of a definite or linear pattern associated with concentration makes it difficult to predict whether the inhibitory role of NAA is due to toxicity or insufficient concentration, as Samiee Rad et al. (2016) pointed out. According to Gao et al. (2019), the insufficient concentration can greatly decrease growth pollen grains germination and pollen tube elongation. However, excluding the anomalies in the results of the concentration of 2 mg/L, it can be considered that the pattern is mostly proportional to the inhibitory role of NAA directly with the increase in its concentration. Similar results were obtained by Gokbayrak and Engin (2017), who found NAA at 0.5, 1.0 and 2.5 mg L⁻¹ did not induce pollen grain germination for three *Vitis vinifera* L. cultivars. Furthermore, Xue et al. (2008) and Sotomayor et al. (2012) observed that NAA inhibited pollen germination and tube growth of Chaohong peach and "Carmel" almond, respectively. NAA was found to hinder the peaches pollen germination, and along with the increase of concentration, the inhibition enhanced (Mi, et al., 2012). On the other hand, it was concluded that NAA is necessary to be used; the minimum concentration of 0.5 mg L⁻¹ was sufficient for pollen germination of *Tulipa greigii* 'Pinocchio' (Akçal et al., 2016). The results of the current investigation, in addition to the previous studies may lead to unclear vision because the examined levels were close. However, in *Syringa oblata* (Liu et al., 2011) and *Castanea mollissima* (Shigang et al., 2017), it was reported that the optimum concentration of NAA for pollen germination and tube growth of was 10 and 15mg/L, respectively. The last two studies indicated that higher levels may be more effective.

Fig.2.d showed higher significant response of PTL to all GA₃ concentrations in the first 24 hours of incubation compared to control (133.94µm) with superiority of 100mg/L GA₃ (232.48µm). However, a slow significant PTL was shown by NAA treatments compared to GA₃ treatments but not with control. As in PG and APV, after 48 hours of incubation, PTL of control accelerated to the adequate limit (270.01 µm) that removes the significant differences with 100mg/L GA₃ (285.57µm) and 50mg/L GA₃ (241.85 µm), and significantly overcomes 10mg/L GA₃ (210.32µm).

NAA treatments continued with significant slow of PTL compared to the control. With the end of the 72 hours of incubation, the longest PTL was recorded at treatment of 100mg/L GA₃ (295.53µm) and control (280.35µm).




Figure 2: Effect of three concentrations of GA₃ at 10, 50 and 100 mg/L and NAA at 1, 2 and 3 mg/L on pollen viability (PV%) (a), pollen germination (PG %) (b), absolute pollen viability (APV %) (c) and pollen tube length (PTL (µm)) (d) at three incubation periods; 24, 48 and 72 hours.

Effect of interaction between pollen grain source and plant hormones (*in vitro*).

At the end incubation period (72 hours), PV of "Ghanami akhdar" was negatively affected with the increase in the concentration of NAA (Table 1). PV decreased to 83.39% and 83.27% by 2mg/L NAA and 3mg/L NAA, respectively, while all concentrations of NAA negatively affected PV of "Jarvis" without exception. PV of "Ghanami akhdar" was significantly affected by adding 100 mg/L GA₃ (81.64%) compared to control (89.19%). The negative effect of GA₃ was shown at the concentration of 50mg/L (83.58%) in the case of "Jarvis" (Table 1).

The higher concentration (100 mg/L) of GA₃ showed a high significant ability to accelerate PG in both cultivars after 24 hours of incubation (Table 1). After 72 hours of incubation, "Ghanami akhdar" showed a higher significant response in PG% with control and 100mg/L GA₃ treatments; 75.15% and 81.62% respectively. Other two concentrations of GA₃ and all NAA concentrations

had a significant negative effect compared to the control. In contrast, "Jarvis" showed a different behavior with wider response involved beside control treatment (82.41%). The two higher concentrations of GA₃; 100mg/L GA₃ (86.94%) and 50mg/L GA₃ (75.93%) in addition to 1mg/L NAA (79.13%) were the best significantly.

Table 1: Interactive effect of NAA and GA₃ on *in vitro* pollen viability and pollen germination of two date palm pollen sources

Treatment	Pollen viability (PV%)			Pollen germination (PG %)		
	24h	48h	72h	24h	48h	72h
Ghanami Akhdar						
Control	89.30 ^{a-c}	89.33 ^{a-c}	89.19 ^{ab}	40.24 ^{c-f}	70.82 ^{a-c}	75.15 ^{a-e}
NAA 1 mg/L	86.40 ^c	85.79 ^{b-d}	86.58 ^{a-d}	40.81 ^{c-e}	60.68 ^{c-f}	69.28 ^{b-f}
2 mg/L	86.87 ^{bc}	88.83 ^{a-d}	83.39 ^{cd}	30.86 ^{e-g}	68.16 ^{a-c}	64.64 ^{d-f}
3 mg/L	87.75 ^{bc}	85.44 ^{cd}	83.27 ^d	20.12 ^g	58.27 ^{c-f}	61.87 ^{ef}
GA ₃ 10 mg/L	88.10 ^{bc}	88.56 ^{a-d}	85.37 ^{a-d}	43.61 ^{c-e}	57.79 ^{c-f}	57.37 ^f
50 mg/L	89.92 ^{a-c}	89.18 ^{a-c}	88.50 ^{a-c}	49.31 ^{cd}	54.41 ^{d-f}	58.86 ^f
100 mg/L	88.15 ^{bc}	90.16 ^a	81.64 ^d	69.53 ^a	77.23 ^{ab}	81.62 ^{a-c}
Jarvis						
Control	92.49 ^a	88.24 ^{a-d}	90.00 ^a	47.30 ^{cd}	79.94 ^a	82.41 ^{ab}
NAA 1 mg/L	88.73 ^{bc}	85.54 ^{b-d}	82.57 ^d	29.59 ^{e-g}	48.35 ^f	79.13 ^{a-c}
2 mg/L	86.96 ^{bc}	89.71 ^{ab}	84.64 ^{b-d}	26.42 ^{fg}	50.80 ^{ef}	58.47 ^f
3 mg/L	87.92 ^{bc}	85.14 ^{cd}	83.02 ^d	38.44 ^{d-f}	64.05 ^{b-e}	63.44 ^{d-f}
GA ₃ 10 mg/L	88.43 ^{bc}	85.56 ^{b-d}	86.05 ^{a-d}	53.03 ^{bc}	65.81 ^{b-d}	68.90 ^{c-f}
50 mg/L	87.89 ^{bc}	84.65 ^d	83.58 ^{cd}	49.67 ^{cd}	65.50 ^{b-d}	75.93 ^{a-d}
100 mg/L	90.12 ^{ab}	87.18 ^{a-d}	84.86 ^{a-d}	64.23 ^{ab}	81.03 ^a	86.94 ^a

Values followed by the different letter(s) in a column are significantly different at p = 0.05 according to Least Significant Difference (LSD).

After 24 hours, APV looked like the trend in PG but with one difference, the absence the high significance of combination of 50mg/L GA₃ with "Ghanami akhdar". APV% was accelerated by 100mg/L GA₃ treatments to be 61.27% compared to 35.99% by control in "Ghanami akhdar" and 57.91% compared to 43.83% in case of "Jarvis". Finally, 100mg/L GA₃ treatments recorded the highest APV% values in "Jarvis" (73.69%) and "Ghanami akhdar" (66.71%) without significant differences compared to control treatments in both cultivars that recorded 74.18% and 67.00%, respectively (Table 2).

It is clear from table 2 that GA₃ concentrations showed a considerable ability to accelerate the elongation of the pollen tubes in the first 24 hours. It increased with increasing concentration to reach 260.95µm in "Ghanami akhdar" and 204.00µm in "Jarvis" at concentration 100mg/L compared to 92.13µm and 175.76µm at the control treatments in both cultivars (Table 2), respectively. Results at 72 hours indicated that most treatments had a negative effect on PTL compared to control treatments except in case of 100mg/L GA₃ and 3mg/L NAA in "Ghanami akhdar" and 50 and 100mg/L GA₃ in case of "Jarvis".

Table 2: Interactive effect of different levels of NAA and GA₃ on *in vitro* absolute pollen viability and pollen tube length of two date palm pollen sources.

Treatment	Absolute pollen viability (APV) %			Pollen tube length (PTL) (µm)		
	24h	48h	72h	24h	48h	72h
Ghanami Akhdar						

Control	35.99 ^{c-e}	63.26 ^{ab}	67.00 ^{ab}	92.13 ^{ef}	240.00 ^{a-d}	279.97 ^{ab}
1 mg/L NAA	35.29 ^{c-f}	52.21 ^{b-e}	60.04 ^{b-e}	125.79 ^{de}	164.20 ^{d-g}	216.03 ^{b-d}
2 mg/L NAA	26.79 ^{e-g}	60.59 ^{a-c}	54.25 ^{c-e}	93.97 ^{ef}	133.60 ^{fg}	148.97 ^d
3 mg/L NAA	17.64 ^g	49.76 ^{c-e}	51.73 ^e	100.96 ^{ef}	103.32 ^g	264.46 ^{ab}
10 mg/L GA ₃	38.36 ^{c-e}	51.14 ^{b-e}	49.18 ^e	193.89 ^{bc}	220.75 ^{c-e}	215.85 ^{b-d}
50 mg/L GA ₃	44.51 ^{cd}	48.44 ^{c-e}	52.21 ^e	195.87 ^{bc}	230.79 ^{b-e}	229.28 ^{b-d}
100 mg/L GA ₃	61.27 ^a	69.61 ^a	66.71 ^{a-c}	260.95 ^a	310.75 ^a	323.74 ^a
Jarvis						
Control	43.83 ^{cd}	70.56 ^a	74.18 ^a	175.76 ^{b-d}	300.02 ^{ab}	280.72 ^{ab}
1 mg/L NAA	26.24 ^{e-g}	41.38 ^e	65.29 ^{a-d}	58.22 ^f	160.42 ^{e-g}	216.92 ^{b-d}
2 mg/L NAA	23.15 ^{fg}	45.61 ^{de}	49.37 ^e	94.60 ^{ef}	164.39 ^{d-g}	163.66 ^{cd}
3 mg/L NAA	33.79 ^{d-f}	54.25 ^{b-d}	52.85 ^{de}	134.23 ^{c-e}	232.38 ^{b-e}	214.26 ^{b-d}
10 mg/L GA ₃	46.78 ^{bc}	56.24 ^{b-d}	59.37 ^{b-e}	165.81 ^{b-d}	199.90 ^{c-f}	204.99 ^{b-d}
50 mg/L GA ₃	43.68 ^{cd}	55.63 ^{b-d}	60.83 ^{b-e}	175.88 ^{b-d}	252.91 ^{a-c}	248.08 ^{a-c}
100 mg/L GA ₃	57.91 ^{ab}	70.92 ^a	73.69 ^a	204.00 ^{ab}	260.39 ^{a-c}	267.32 ^{ab}

Values followed by the different letter(s) in a column are significantly different at $p = 0.05$ according to Least Significant Difference (LSD).

Effect of NAA and GA₃ and pollen source on fruit set of "Medjool" date palm

In this field study, it appears from (Fig.3.a) that "Ghanami akhdar" has clearly and significantly outperformed over "Jarvis" in the percentage of set fruits in both growth stages; 7DAP and 120DAP (Khelal stage), where the percentage in "Ghanami akhdar" was 88.17% and 43.26% in both stages, respectively, compared to 85.07% and 39.95% in "Jarvis", respectively. Which may be due to the stronger metazinic effect of "Ghanami akhdar". The variations in genetic heritage and growth circumstances could also contribute to the differences between the two date palm pollinators. According to many studies, fruit set in date palm is one of the characteristics associated with the metazinic effect of the pollen source, (Sarrwy et al. (2014), Hafiz et al. (2014), Omaima et al., (2015), Salomon-Torres et al. (2017), Shahid et al. (2017), Mohammadi et al. (2017), Iqbal et al., (2019) and El-Hamady, et al. (2010).

The current study findings indicated that the general genotype effect were absent *in vitro* except in case of slightly superiority of "Jarvis" over "Ghanami akhdar" started just at the end of incubation period with related only to PG% and APV%, but genotype effects appeared clearly *in vivo* with superiority of "Ghanami akhdar" over than "Jarvis". The absence of this influence in most parameters *in vitro* and their appearance *in vivo* may be attributed to the higher metazenic effect of "Ghanami akhdar" pollen on characteristics of the tested female date palm which happened due to pollen-pistil reaction as (Swingle, 1928). According to Patel and Mankad (2014), Pollen's germination efficiency could be critical not just in fruit set but also in flower-flower and flower-pollinator interactions. On the other hand, this discrepancy between *in vitro* and *in vivo* may show a relative deficiency in the method of *in vitro* and its ability to simulate the real conditions in the field, and may make it a somewhat invalid method, especially in plants in which the phenomenon of metazinia is a limited factor such as date palm and may be this is what Read et al. (1993) meant when he indicated that even if the conditions for pollen germination were ideal, they do not fully mimic *in vivo* growth.




Figure 3: Effect of two pollen grain sources; "Ghanami Akhdar" and "Jarvis" (a) as well as NAA at 1, 2 and 3mg/L and GA₃ at 10, 50 and 100mg/L on fruit set % of "Medjool" date palm at two growth stages; 7DAP and Khelal (b).

Looking in some detail at the results of this experiment, the general effect of both genotypes may not reflect the reality. "Jarvis" is correct that it lags behind "Ghanami akhdar", but it had a better response to GA₃, and the reason for its lower general effect is its greater sensitivity to NAA (Table 2), and it may be the most appropriate to compare the performance of the two cultivars with regard to the metazenic effect of pollen is the adoption of the results of the two control treatments for both cultivars, which showed superiority to "Ghanami akhdar" in the early stages of setting fruits, but they showed clear equivalence at Khelal stage.

Table 3: Interactive effect of different levels of NAA and GA₃ and two date palm pollen sources on fruit set % of "Medjool" at two growth stages

Treatment	Ghanami Akhdar		Jarvis	
	7 DAP	Khelal	7 DAP	Khelal
Control	93.52 a	45.57 ^{bc}	84.05 ^{d-f}	46.94 ^{bc}
NAA 1 mg/L	91.78 ab	37.34 ^{de}	88.36 ^{b-d}	31.49 ^{ef}
2 mg/L	86.08 ^{c-f}	38.21 ^d	82.12 ^{ef}	28.06 ^f
3 mg/L	86.91 ^{b-e}	44.58 ^c	81.77 ^f	31.95 ^{ef}
GA ₃ 10 mg/L	91.00 a-c	41.67 ^{cd}	89.71 a-c	30.96 ^f
50 mg/L	84.55 ^{d-f}	44.44 ^c	86.55 ^{c-f}	55.41 a
100 mg/L	86.24 ^{ef}	50.99 ab	84.90 ^{ef}	54.84 a

Values followed by the different letter(s) in a column are significantly different at p = 0.05 according to Least Significant Difference (LSD).

With regard to the effect of hormone, there were clear differences in the results of the fruit set % between the early stage and the late stage of growth has been shown (Fig. 3b). The control treatment (88.78%) and the lowest concentration of both GA₃ and NAA (90.36% and 90.07%, respectively) was significantly superior to the rest of the treatments after 7 days of pollination. The results were different in the Khelal stage with the superiority of the two treatments of the

highest concentration of GA₃ on the rest of the treatments, where 100mg/L GA₃ recorded 52.91% and 50 mg/L GA₃ treatment 49.93% compared to 46.25% for the control. Auxins and gibberellins are commonly utilized to regulate fruit abscission and enhance the fruit's quality from the time of fruit set to ripening and eventual delivery to the consumer (Suman et al 2017). De Jong et al., (2009) suggested that the levels of AUX and GA genes in ovules are increased After pollination, leading to the activation of AUX and GA response genes. Consequently, these response genes will initiate the growth and development of fruits by controlling both cell division and cell expansion.

In our current investigation, the lack of effect of lower concentrations of both hormones on the percentage of setting fruits in the early stages after pollination may be due to their insufficiency to reach the toxicity level or to the extent that an imbalance occurs in the endogenous hormonal balance and vice versa with regard to the higher concentrations of them (Samiee Rad et al., 2016). Endogenous GA₃ at a definite level was a prerequisite for the successful germination of pollen grains. All NAA concentrations significantly reduced the fruit setting percentage, while it was increased by 50~100 mg/l GA₃. Ozga and Reinecke (2003) noted that the hormonal balance fluctuates throughout various stages of fruit development, whereas a single hormone can serve distinct roles at each stage.

Shahsavar and Shahhosseini (2021) found that pollen grains containing higher amounts of gibberellin during pollination on "Piarom" dates can trigger more effective fertilization followed by seed production and larger amount of gibberellin in the fruit as a result. The utilization of exogenous GA₃ + diphenylurea + NAA during full bloom has demonstrated its effectiveness in enhancing fruit set by increasing the quantity of pollen tubes that successfully reached the ovaries of pollinated flowers (Williams and Flook, 1980). Al-Samaraie and Al-Falahy (2020) determined that GA₃ has a beneficial impact on decreasing the fruit drop percentage and enhancing the weight of the "Braum" date palm fruits. This positive effect can be attributed to the increased rate of assimilation materials moving towards the fruits, resulting in an increase in the overall weight of the fruit bunch. The current study showed that NAA treatments led to a decrease in the percentage of fruit setting as the concentration increased and with a regular behavior during the two growth stages. Fruit set % was about 37~44% by "Ghanami akhdar" pollen and 28~31% by "Jarvis". NAA is commonly used as chemical thinner at petal fall and earlier phenological stages of apple fruitlets, and it is effective at low rates (Wertheim, 2000), it is almost inhibitor at the applied concentrations by hindering most of study parameters *in vitro*. Similar behavior appeared *in vivo*, NAA decreased fruit set percentage over 1mg/L 7 DAP and by all levels 120 DAP (Khelal stage) (Fig.2.a).

The interaction effect between the pollinator and the hormonal treatment is shown in (Table3). 7DAP, the percentage of fruit set in "Ghanami akhdar" significantly decreased at the higher two concentrations of each hormone compared to the control. The treatment 50mg/L GA₃ led to the lowest fruit set (84.55%). On other hand, the lowest concentration of NAA and 10 mg/L GA₃ contributed to a significant improvement in fruit set of "Jarvis" (88.36% and 89.71% respectively) compared with the control (84.05%); the highest percentage of fruit set was observed at concentration 10 mg/L GA₃ (89.71%). The treatment of 3mg/L NAA showed the lowest fruit set (81.77%) at Khelal stage. The percentage of fruit set in "Ghanami akhdar" increased significantly at the concentration of 100 mg/L GA₃ (50.99%) compared to the control treatment (45.57%).

Conclusion

It is concluded from this study that GA₃ at 100mg/L can improve date palm pollen germination. Application of this concentration upon pollination also led to the best results in term of fruit set at Khelal, the stage of full fruit size. These findings open the doors to investigate the interactions between indigenous and exogenous GA₃ in the future. This relationship was clear in term of NAA, same behavior *in vitro* and *in vivo* appeared. NAA at these concentrations can present an effective thinning method in "Medjool". On other hand, "Ghanami akhdar" can be considering as a recommendable pollinator for "Medjool". It is advisable to apply GA₃ at rate of 100mg/L to modified BK to accelerate date palm pollen germination and pollen tube growth.

References

- Acar, I., Ak, B. E., & Sarpkaya, K. (2010). Effects of boron and gibberellic acid on *in vitro* pollen germination of pistachio (*Pistacia vera* L.). African Journal of Biotechnology, 9(32), 5126-5130.

- Alasasfa, Muawya. 2019. Effects of Water Spray Timing after Pollination on Fruit Set, Yield, Physical and Chemical Properties of Two Date Palm Cultivars. *Journal of Biology, Agriculture and Health care*.9 (10):74-80.
- Alasasfa, Muawya. 2021. Effect of Pollination Methods on Fruit Set, Yield, Physical and Chemical Properties of Hayani Date Palm Cultivar. *International Journal of Environmental & Agriculture Research (IJOEAR)*. 7(1): 24-30.
- Al-Jibouri, A. A. M., Fattah, F. A., Alsaadawi, I. S., Selbi, M., & Kgazal, M. (1990). Morphometrics and scanning electron microscopy of pollen of date palm (*Phoenix dactylifera* L.). *Arab Gulf Journal of Scientific Research*, 8(2), 109-117.
- Almeida, F. C. G., da Silva, J. F., Alves, J. F., da Silva, F. P., & Almeida, F. A. G. (1990). Germination of cotton pollen in vitro. III. Effect of gibberellic acid. *Pesquisa Agropecuária Brasileira*, 25(11), 1603-1605.
- Al-Samaraie, O. H., & Al-Falahy, T. H. (2020). Effect of Bunch Covering with Different Colors of Polyethylene and GA3 Spray in Some Fruits Characteristics of Date Palm cv. Braim. *Indian Journal of Ecology*, 47(Special Issues (10)), 132-137.
- De Jong, M., Mariani, C. and Vriezen, W. H. (2009). The role of auxin and gibberellin in tomato fruit set. *Journal of Experimental Botany* 60 (5):1523-1532
- El-Hamady, M., Hamdia, M., Ayaad, M., Salama, M. E., & Omar, A. K. (2010). Metaxenic effects as related to hormonal changes during date palm (*Phoenix dactylifera* L.) fruit growth and development. In IV International Date Palm Conference 882 (pp. 155-164).
- Gao, C., Wang, Y., & Qu, H. (2019). Study of auxin regulation of pollen tube growth through calcium channels in *Pyruspyrifolia*. *Plant Growth Regulation*, 89(1), 99-108.
- Gökbayrak, Z., & Engin, H. (2018). Effects of foliar-applied brassinosteroid on viability and in vitro germination of pollen collected from bisexual and functional male flowers of pomegranate. *International Journal of Fruit Science*, 18 (2), 226-230.
- Gokbayrak, Z., & Engin, H. (2017). Brassinosteroids and gibberellic acid: effects on in vitro pollen germination in grapevine. *OENO One*, 51(3), 303-303
- Gökbayrak, Z., Engin, H., Akçal, A., & Kiraz, H. (2020). Pollen of *Arbutus unedo*: Effects of plant growth regulators. *Botanica Serbica*, 44(1), 55-59.
- Hafiz, O., M.A. Saleh, E.M.A. Mostafa, A.M. Naguib and N.E. Ashour (2014). Effect of pollen grain source on yield and fruit quality of Samany date palm. *Int. J. Agric. Res.* 1-5.
- Iqbal, M., Usman, K., Jatoi, S. A., & Munir, M. (2019). Fruit yield and quality of Dhakki date as affected by fresh and stored pollens of potential male genotypes. *Pak. J. Bot*, 51, 5.15.
- Jain, S. M., & Johnson, D. V. (2015). Date palm genetic resources and utilization (Vol. 1). J. M. Al-Khayri (Ed.). Africa and the Americas: Springer.
- Khalifa, T., Jwanh, M.Z., & Alsalem, M. I., (1983). Palms and dates in Kingdom Saudi Arabia. Kingdom of Saudi Arabia.
- Krueger, R. R. (2021). Date Palm (*Phoenix dactylifera* L.) Biology and Utilization. In *The Date Palm Genome*, Vol. 1 (pp. 3-28). Springer, Cham.
- Liu, X., Yang, Q., Qin, J., & Liu, L. (2011). Effects of plant growth regulators on in vitro pollen germination of *Syringa oblata*. *Journal of Northeast Forestry University*, 39(10), 54-55.
- Maita, S., & Sotomayor, C. (2015). The effect of three plant bioregulators on pollen germination, pollen tube growth and fruit set in almond *Prunus dulcis* (Mill.) DA Webb] cvs. Non Pareiland Camel. *Electronic Journal of Biotechnology*, 18(5), 381-386.
- Mi, Y., Wang, J., & Hu, H. (2012). Effect of plant growth regulators on pollen germination of peaches. *Southwest China Journal of Agricultural Sciences*, 25(3), 1023-1026.
- Mohammadi, N., Rastgoo, S., & Izadi, M. (2017). The strong effect of pollen source and pollination time on fruit set and the yield of tissue culture-derived date palm (*Phoenix dactylifera* L.) trees cv. Barhi. *Scientia Horticulturae*, 224, 343-350.
- Omaima, M. Hafaz, Malaka A. Saleh, N.E. Ashour, E.A. Mostafa and M.M. Naguib (2015). Evaluation of some pollen grain sources on yield and quality of Sanany Date palm cv. (*Phoenix dactylifera* L.). Vol. 4 Issue: 1 Jan-Mar 2015 pages: 27-30.
- Ozga, J. A. and Reinecke, D. M. (2003). Hormonal interaction in fruit development, *Journal of Plant Growth Regulation* 22:73-81.
- Patel, R. G., & Mankad, A. U. (2014). In vitro pollen germination-A review. *International Journal of Science and Research*, 3(5), 304-307.

- Prajapati, P., & Jain, B. (2011). 84-221. Effects of Gibberallic Acid GA₃ On in Vitro Pollen Germination and Pollen Tube Growth Inluffa Aegyptica Mill. Byp. P. Prajapati and b. K. Jain. Life Sciences Leaflets, 221-to.
- Radovic, A., Nikolic, D., Milatovic, D., Živkovic, B., & Stevanovic, N. (2016). The effect of plant hormones on pollen germination and pollen tube growth of almond cultivars. *Acta Horticulturae*, 1139, 375-380.
- Salinas-Grenet, H., Herrera-Vásquez, A., Parra, S., Cortez, A., Gutiérrez, L., Pollmann, S., ... & Blanco-Herrera, F. (2018). Modulation of auxin levels in pollen grains affects stamen development and anther dehiscence in Arabidopsis. *International journal of molecular sciences*, 19 (9), 2480.
- Salomon-Torres, R., Ortiz-Urbe, N., Villa-Angulo, R., Villa-Angulo, C., Norzagaray-Plasencia, S., & GARCIA-Verdugo, C. R. I. S. T. I. A. N. (2017). Effect of pollenizers on production and fruit characteristics of date palm (*Phoenix dactylifera* L.) cultivar Medjool in Mexico. *Turkish Journal of Agriculture and Forestry*, 41(5), 338-347.
- Samiee Rad, Z., Imani, A., & Salmani, M. (2016). The interaction effects of boron and plant growth regulators on pollen germination of almond. *Journal of Nuts*, 7(01), 45-50.
- Sanjay Kumar, K., Prasad Khanduri, V., Kar, K., Mohan Sharma, C., & Kumar Riyal, M. (2016). Effect of growth regulators and time on in vitro pollen germination in three ornamental tropical tree species. *Journal of Agricultural Science and Technology*, 18(5), 1247-1255.
- Sarrwy, S.M.A., A.A.A. Haiba., S.A.A. Attia and Enas A.M. Ali (2014) Influence of Different pollen Grains Sources on yield and fruit quality of siwi date palm and their phylogenetic relationships using RAPD markers. *Middle East Journal of Agric Res*, 3(2):330-337,(2014).
- Shahid, M. A., Iqbal, M., & Niamatullah, M. (2017). Response of male pollinizers in fruit set, yield and quality of date palm (*Phoenix dactylifera* L.). Cv. Dhakki. *Sarhad Journal of Agriculture*, 33(1).
- Shahsavar, A. R., & Shahhosseini, A. (2021). Pollen grain hormones of date palm pollinator cultivars and their relationship with hormones of different stages of 'Piarom'date fruit growth. *Scientia Horticulturae*, 288, 110389.
- Shenxi, X., Xianshi, L., Yuechang, W., & CJ, L. (2004). Pollen viability of Asian pear and effect of PGR, B and sucrose on germination and pollen tube development. *Journal of Fruit Science*, 4, 289-294.
- Sotomayor, C., Castro, J., Velasco, N., & Toro, R. (2012). Influence of seven growth regulators on fruit set, pollen germination and pollen tube growth of almonds. *Journal of Agricultural Science and Technology. B*, 2(9B), 1051.
- Suman, M., Sangma, P. D., Meghwal, D. R., & Sahu, O. P. (2017). Effect of plant growth regulators on fruit crops. *Journal of Pharmacognosy and Phytochemistry*, 6(2), 331-337.
- Swingle, W. T. (1928). Metaxenia in the date palm: possibly a hormone action by the embryo or endosperm. *Journal of Heredity*, 19(6), 257-268.
- Visser, T., De Vries, D. P., Scheurink, J. A. M., & Welles, G. W. H. (1977). Hybrid tea-rose pollen. II. Inheritance of pollen viability. *Euphytica*, 26(3), 729-732.
- Wertheim, S. J. (2000). Developments in the chemical thinning of apple and pear. *Plant growth regulation*, 31(1), 85-100.
- Williams, R. R., & Flook, V. A. (1980). The mode of action of the hormone apple fruit-setting mixture PP 341B applied to Cox's Orange Pippin. *Journal of horticultural science*, 55(3), 275-277.
- Wu, J., Qin, Y., & Zhao, J. (2008). Pollen tube growth is affected by exogenous hormones and correlated with hormone changes in styles in *Torenia fournieri* L. *Plant Growth Regulation*, 55(2), 137-148.
- Xue, X. M., Wang, J. Z., Zhang, A. N., & Lu, C. (2008). Effects of plant growth regulating substances on pollen germination and tube growth in Chaohong peach. *Journal of Northwest A & F University (Natural Science Edition)*, 4. 66-82.