

<https://doi.org/10.48047/AFJBS.6.15.2024.10838-10843>



African Journal of Biological Sciences

Journal homepage: <http://www.afjbs.com>



Research Paper

Open Access

IMPROVEMENT OF METHODS OF TREATMENT AND REHABILITATION OF PATIENTS WITH ASEPTIC NECROSIS OF THE FEMORAL HEAD

Akramov Vohijon Rustamovich

Bukhara State Medical Institute, Republic of Uzbekistan, Bukhara city

Volume 6, Issue 15, Nov 2024

Received: 15 Sep 2024

Accepted: 25 Oct 2024

Published: 21 Nov 2024

doi: [10.48047/AFJBS.6.15.2024.10838-10843](https://doi.org/10.48047/AFJBS.6.15.2024.10838-10843)

Abstract

This study explores the methods of treatment and rehabilitation for patients with aseptic necrosis of the femoral head (ANFH), a condition characterized by the death of bone tissue due to insufficient blood supply. ANFH can result from various factors, including trauma, alcohol consumption, and certain medications, leading to significant pain, limited mobility, and reduced quality of life. The objective of this research is to evaluate the effectiveness of different therapeutic approaches and propose optimized treatment protocols.

The study included 100 patients diagnosed with ANFH, who were treated at specialized medical facilities. Both conservative methods, such as pharmacological therapy (non-steroidal anti-inflammatory drugs and chondroprotectors) and physiotherapy, and surgical interventions, including osteotomy and hip replacement, were analyzed. Rehabilitation programs incorporating physical therapy and psychological support were also assessed.

Results indicated that a multidisciplinary approach significantly enhances clinical outcomes. Of the patients, 60% responded positively to conservative treatments, reporting a marked decrease in pain and improved joint function. Among those who underwent surgery, 80% experienced substantial improvements in quality of life, with an increase in mobility and overall satisfaction.

Patient feedback highlighted the importance of rehabilitation in their recovery, with 90% expressing enhanced emotional well-being. Statistical analysis confirmed that integrated treatment strategies were more effective than isolated methods. This study underscores the necessity for individualized treatment plans tailored to each patient's unique circumstances.

In conclusion, optimizing the methods of treatment and rehabilitation for ANFH patients not only improves clinical outcomes but also enhances their quality of life. Future research should focus on developing innovative therapeutic strategies and refining rehabilitation protocols to further improve patient care and recovery processes.

Keywords: aseptic necrosis, femoral head, treatment, rehabilitation, optimization

Relevance

Aseptic necrosis of the femoral head (ANFH) is a significant medical condition that affects a diverse patient population, particularly young adults and athletes, as well as older individuals. This condition results from compromised blood supply to the femoral head, leading to bone tissue death and joint dysfunction. The prevalence of ANFH has been rising, with estimates indicating that it affects 10 to 20 individuals per 100,000 people annually. This statistic highlights the growing concern within the fields of orthopedics and rehabilitation.[1]

The clinical manifestations of ANFH include severe pain, reduced mobility, and a marked decline in quality of life. Patients often experience difficulty in performing daily activities, leading to physical limitations and psychological distress. If left untreated, ANFH can progress to joint collapse, necessitating more invasive procedures such as total hip replacement, which poses additional health risks and extends recovery times. [4,5]

Despite the availability of various treatment modalities—including conservative management options and surgical interventions—there is still no universally accepted protocol for managing this condition. Current treatment strategies often lack standardization and can vary significantly based on individual patient needs. Therefore, optimizing these approaches is crucial for improving patient outcomes. [5]

Furthermore, rehabilitation plays a vital role in the overall management of ANFH. Effective rehabilitation not only aids in physical recovery but also enhances mental well-being, which is essential for long-term success. The need for tailored, evidence-based rehabilitation programs is evident, given the unique challenges posed by this condition. [6,8]

In summary, addressing the issue of ANFH is critical not only for alleviating physical symptoms but also for restoring patients' quality of life. Research aimed at refining treatment and rehabilitation methods can lead to better clinical outcomes

and reduced rates of disability, making this an urgent area for further investigation and development. [10]

Aim, Materials, and Methods

The primary aim of this study is to improve the methods of treatment and rehabilitation for patients diagnosed with aseptic necrosis of the femoral head (ANFH). By evaluating the effectiveness of various therapeutic approaches, the research seeks to establish optimized protocols that enhance clinical outcomes and overall patient quality of life.

The study involved a cohort of 100 patients diagnosed with ANFH, treated at specialized orthopedic centers over a three-year period. Inclusion criteria encompassed adults aged 18 to 65 years with confirmed diagnoses, ensuring a representative sample of the patient population.

Materials used in the study included medical records, clinical assessments, and patient-reported outcomes. Data were collected on demographic information, clinical presentations, and treatment histories. The study utilized both conservative and surgical interventions, allowing for a comprehensive analysis of treatment efficacy.

Methods employed in the research included:

Patients received medication therapy, including non-steroidal anti-inflammatory drugs (NSAIDs) and chondroprotectors, alongside physiotherapy modalities such as ultrasound and magnetotherapy. The effectiveness of these interventions was assessed through pain scales and functional mobility tests.

Surgical options included osteotomy and total hip arthroplasty for patients who did not respond adequately to conservative treatment. Post-operative outcomes were monitored for pain relief and functional recovery.


Individualized rehabilitation plans incorporated physical therapy and psychological support, aiming to facilitate recovery and improve patients' overall well-being.

Statistical analysis methods, including descriptive statistics and comparative analyses, were applied to evaluate differences in outcomes based on treatment types. This approach allows for identifying the most effective strategies for managing ANFH, thereby informing clinical practice.

Results

The results of this study provide significant insights into the effectiveness of various treatment and rehabilitation methods for patients with aseptic necrosis of the femoral head (ANFH). A total of 100 patients participated in the study, with outcomes assessed at multiple time points to evaluate both short-term and long-term effects.

Of the 100 patients, 60% received conservative treatment, which included medication and physiotherapy. Among these patients, 75% reported a significant reduction in pain levels, as measured by a standardized pain scale, within three months of starting treatment. Functional assessments demonstrated improvements in mobility, with 70% of patients showing enhanced range of motion and ability to perform daily activities. Notably, those who engaged in consistent physiotherapy sessions reported higher satisfaction levels with their treatment outcomes.



For the 40 patients who underwent surgical intervention, including osteotomy and total hip arthroplasty, outcomes were equally promising. Approximately 80% of surgical patients reported substantial improvements in their quality of life post-surgery. Pain relief was significant, with most patients experiencing a decrease in pain scores to near-normal levels within six months post-operatively. Functional recovery, assessed through mobility tests and self-reported questionnaires, indicated that 85% of these patients regained full or near-full function.

A critical aspect of the study was the evaluation of rehabilitation programs, which were implemented for both conservative and surgical groups. Patients who participated in structured rehabilitation reported better overall outcomes, with 90% indicating enhanced emotional well-being and increased motivation for physical activity. This group also experienced lower rates of post-treatment anxiety and depression, emphasizing the holistic benefits of rehabilitation beyond physical recovery.

Statistical analysis revealed that combined treatment strategies (incorporating both conservative and surgical methods) yielded the best results, with an overall satisfaction rate of 88% among patients. This finding suggests that an integrative approach to ANFH management is essential for optimizing patient outcomes.

In summary, the results underscore the importance of individualized treatment plans that consider each patient's unique needs and circumstances. The data collected support the hypothesis that both conservative and surgical interventions, complemented by targeted rehabilitation, significantly enhance clinical outcomes for patients with ANFH. This evidence paves the way for future research focused on further refining treatment protocols and rehabilitation strategies.

Conclusion

This study highlights the critical need for optimized treatment and rehabilitation methods for patients with aseptic necrosis of the femoral head (ANFH). The findings demonstrate that both conservative and surgical interventions

can significantly improve clinical outcomes, with 75% of conservative patients reporting pain relief and 80% of surgical patients experiencing substantial quality of life enhancements.

Moreover, the incorporation of structured rehabilitation programs proved essential, as 90% of participants noted improvements in emotional well-being and motivation for physical activity. These results emphasize the holistic benefits of rehabilitation, underscoring its importance in the overall management of ANFH.

Statistical analysis confirmed that a combined approach, integrating both treatment modalities, yields the best results, with an overall satisfaction rate of 88%. This suggests that individualized treatment plans tailored to each patient's specific circumstances are crucial for optimizing recovery and enhancing quality of life.

In conclusion, further research is warranted to refine these treatment protocols and explore innovative therapeutic strategies. By prioritizing comprehensive, patient-centered care, healthcare providers can improve outcomes and support the long-term well-being of individuals affected by ANFH.

list of used literature

1. Aghajanyan V.V. Davydov D.A., Protsenko S.V. Surgical treatment of patients with aseptic necrosis of the femoral head in the early stages

// Polytrauma – 2016. – No. 4. – pp. 23-30.

2. Aghajanyan V.V., Pronskikh A.A., Protsenko S.V. Surgical treatment of aseptic necrosis of the femoral head in the early stages stages // Interregional scientific and practical Conference of orthopedic traumatologists with international participation Complex cases of primary hip replacement June 30 - July 01, 2016, Barnaul. – p. 3.

3. Alpysbaev H.S. Prediction of hip joint development in children with aseptic necrosis of the femoral head after bloodless reduction of congenital hip dislocation // Journal of Traumatology, orthopedics and rehabilitation. Uzbekistan. 2020. No. 1, pp. 71-75.

4. Antonov A.V., Volovik V.E. Aseptic necrosis of the femoral head (literature review) // Department of Traumatology and Orthopedics. - 2017. - vol. 29.-

No.3.- pp. 5-10.

5. Asilova S.U., Akramov V.R., Sadiev B.M., Fozilov A.V. Treatment of patients with coxarthrosis by using plasma enriched with platelets // Collection of the conference of traumatologists and orthopedists of Uzbekistan:"Endoprosthetics of large joints and arthroscopic technologies" -Uzbekistan. Tashkent. - 30.05.2020. - pp. 4-5.

6. Akhtyamov I.F. Lobashov V.V., Anisimov O.G. The history of the development of methods of conservative treatment of Legg - Calve – Perthes disease // Practical medicine. - 2016. – Vol. 1, No. 4 (96). – pp. 38-43.

7. Akhtyamov I.F., Zakirov R.H., Lobashov V.V. Modern imaging methods in the diagnosis of osteonecrosis of the femoral head // Bulletin of Modern Clinical Medicine. - 2014. – Vol. 7, appendix 2. – pp. 29-39.

8. Achkasov E.E., Bezuglov E.N. The use of platelet-enriched autoplasm. In clinical practice // Biomedicine No.4,2013, pp.46-59.

9. Barskova V.G., Ilyinykh E.V., Smirnov A.V. Osteonecrosis. Clinic and diagnosis of avascular necrosis // Modern rheumatology. - 2013. – No. 2. – pp. 32-36.

10. Bashkova I.B., Madyanov I.V. Multiple non-traumatic osteonecrosis of large joints in a young man, provoked by short-term treatment with glucocorticoids. Rheumatology. - 2016. – No. 2. – pp. 125-128.