



## Comparative impairing effects of selected arthropod venoms on the main body metabolites of *Galleria mellonella* (Lepidoptera: Pyralidae)

Mohammad Tanani<sup>1</sup>, Karem Ghoneim<sup>2\*</sup>, Khalid Hamadah<sup>3</sup>, and Dyaa Emam<sup>4</sup>

<sup>1</sup>Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt. E-mail: [tanani484@gmail.com](mailto:tanani484@gmail.com)

<sup>2</sup>Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt. E-mail: [karemghoneim@gmail.com](mailto:karemghoneim@gmail.com)

<sup>3</sup>Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt. E-mail: [khalidhamadah@gmail.com](mailto:khalidhamadah@gmail.com)

<sup>4</sup>Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt. E-mail: [Dyaa2015@gmail.com](mailto:Dyaa2015@gmail.com)

### Article Info

Volume 3, Issue 3, July 2021

Received : 19 February 2021

Accepted : 06 June 2021

Published : 05 July 2021

doi: [10.33472/AFJBS.3.3.2021.64-79](https://doi.org/10.33472/AFJBS.3.3.2021.64-79)

### Abstract

The greater wax moth, *Galleria mellonella* (Lepidoptera: Pyralidae) is widely distributed in the world. It is the most destructive pest of honey bee, *Apis mellifera* (Hymenoptera: Apidae), throughout the world. The present study was conducted to estimate the disturbance of main body metabolites in the tissues of 5<sup>th</sup> and 7<sup>th</sup> instar larvae, as well as early-, mid- and late-aged pupae of *G. mellonella*, by certain arthropod venoms. The 3<sup>rd</sup> instar larvae were treated, via an artificial diet, with LC<sub>50</sub> of death stalker scorpion *Leiurus quinquestriatus* venom, oriental Hornet (wasp) *Vespa orientalis* venom, or Apitoxin of honey bee *Apis mellifera* (3428.9, 2412.6, and 956.16 ppm, respectively). The important results could be summarized as follows. The total protein content in larvae was drastically reduced in larvae, regardless the tested venom or the larval instar. Also, all venoms predominantly prevented the successfully developed pupae to attain normal protein level. All venoms exhibited prevalent inducing effects on larvae to gain more lipids than control congeners of both 5<sup>th</sup> and 7<sup>th</sup> instars. Also, all tested venoms unexceptionally enhanced the pupae to gain more lipids than the control congeners. Treatment of the 3<sup>rd</sup> instar larvae with each of the tested venoms resulted in a dramatic reduction of carbohydrate content in 5<sup>th</sup> and 7<sup>th</sup> instar larvae. The carbohydrate content in treated pupae had been drastically reduced, regardless the tested venom or the pupal age.

**Keywords:** Apitoxin, Carbohydrate, Hornet, Larva, Lipid, Protein, Pupa, Scorpion

© 2021 Mohammad Tanani et al. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## 1. Introduction

The greater wax moth *Galleria mellonella* L. (Lepidoptera: Pyralidae) is widely distributed throughout the world. Although the adults do not feed, because they have atrophied mouth parts, the voracious nature of larval feeding and tunneling lead to the destruction of the honeycomb, and subsequently to the death of weak colonies (Chandel et al., 2003; Awasthi and Sharma, 2013; Mohamed et al., 2014; and Kwadha et al., 2017). For the control of *G. mellonella*, various physical methods have been adopted; including freezing, heating, CO<sub>2</sub>, Ozone gas and sulphur fumigation against larvae and pupae (Owayss and Abd-Elgayed, 2007; Christen et al., 2008;

\* Corresponding author: Karem Ghoneim, Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt. E-mail: [karemghoneim@gmail.com](mailto:karemghoneim@gmail.com)

Akyol et al., 2009; and James, 2011). Conventional insecticides of different categories had been used for controlling *G. mellonella* (Durmuş and Büyükgüzel, 2008; and Sak and Uçkan, 2009). Several biological control agents, such as the natural enemies, predators and parasitoids, along with entomopathogenic nematodes, viruses and fungi, had been assessed for controlling this pest (Dindo et al., 2001; Armendariz et al., 2002; Hussaini, 2003; Ellis et al., 2013; and George et al., 2019). The sterile insect technique (or inherited sterility) has been assessed against this pest (Ebadi et al., 2001; Carpenter et al., 2005; and El-Kholly and Mikhael, 2008). Also, insect hormone analogues, insect growth regulators had been assessed against it (Izzetoglu and Karacali, 2003; Unsal et al., 2004; Awasthi and Sharma, 2012; and Pamita and Priyanka, 2013). Natural compounds of the plant origin may be efficient alternatives to conventional fumigants against *G. mellonella* (Abbasipour et al., 2009; Mahmoudvand et al., 2011; Basedow et al., 2012; Elbeherly et al., 2016; and Er et al., 2017).

In the last few decades, a great interest of investigation by agrochemical companies is the development of highly selective biopesticides derived from animals. Natural products of the animal origin have been described as very good alternative agents to the conventional insecticides for controlling some insect pests. These animal-derived biopesticides include the venom-derived peptides from different sources including the venomous arthropods, such as spiders (Harrison and Bonning, 2000; Tedford et al., 2004; and Nicholson, 2006), scorpions (Froy et al., 2000; and Taniai et al., 2002), wasps (Dahlman et al., 2003), as well as cone snails (Olivera, 2002) and some marine animals (Whetstone and Hammock, 2007; Windley et al., 2012; and Nakasu et al., 2014). In addition, the arthropod hormones and neuropeptides may be effective control agents against various insect pests (Altstein et al., 2000; and Altstein, 2004).

Scorpion is a mysterious creature in the animal world. It has poisonous venom (Possani et al., 2000) which has increasingly attracted the scientists' attention throughout the world (Cao et al., 2013; and Ma and Shi, 2014). The death stalker scorpion, or yellow scorpion, *Leiurus quinquestriatus* (Buthidae: Arachnida) can be found in desert and scrubland habitats ranging from North Africa through to the Middle East. In Egypt, Saleh et al. (2017) reported the occurrence of this scorpion species in six eco-geographical regions. Among different scorpion venoms, venom of *L. quinquestriatus* exhibited the most potent toxicity against the meal worm *Tenebrio molitor* (Valk and Meijden, 2014). As reported by some authors (Gurevitz, 2010; and Leng et al., 2011), the scorpion toxins contain active toxins against insects and are valuable as leads for the development and synthesis of eco-friendly insecticides, since they exhibited no effect on beneficial insects or mammals (Fabiano et al., 2008; and Gurevitz, 2010). However, Joseph and George (2012) reviewed the insecticidal activities of scorpion toxins on a broad range of insect pests and concluded that the scorpion toxins provide safe biopesticides.

Workers and queen of the honey bee *Apis mellifera* (Apidae: Hymenoptera) produce the venom in a special long and thin branched acid gland at the end of their abdomens. This venom or toxin can be called Apitoxin; since the word was originated from the Latin *apis* (bee) and *toxikon* (venom) (Cruz-Landim and Abdalla, 2002). In a recent review, Azam et al. (2019) compiled information on the history, chemical composition and scientific evidence concerning the Apitoxin pharmaceutical research and different medical uses. The honey bee venom had been studied for its action on mammals although little is known about its action on insects (Piek, 1987; and Quistad et al., 1988). This venom exhibited toxic effects on some insects, such as the corn earworm *Heliothis zea* (Ross et al., 1987), the tobacco hornworm *Manduca sexta* (Quistad et al., 1988) and the lesser wax moth *Achroia grisella* (Mahgoub et al., 2018). Recently, Ghoneim et al. (2019a) recorded a dose-dependent toxicity of Apitoxin on larvae and pupae of *G. mellonella*, as well as the reduction of larval weight gain and growth rate. In another recent study, Apitoxin blocked the adult emergence, prohibited the fecundity and fertility of *G. mellonella* (Ghoneim et al., 2019b).

The oriental hornet *Vespa orientalis* (Hymenoptera: Vespidae), is a social wasp in the Middle East (Khodairy and Awad, 2013; and Abdelaal and El-defrawy, 2014). The world distribution of this wasp species comprises, also, North Africa, southeastern Europe, Southwest Asia across Turkey and Arabian Peninsula to India and Nepal (Carpenter and Kojima, 1997; and Archer, 1998). In addition, it was accidentally introduced into Madagascar and China (Carpenter and Kojima, 1997; and Archer, 1998) and recorded in Mexico (Dvorak, 2006). Many studies had been conducted for examining the toxicity of wasp venoms on insects (Libersat et al., 1999; Coudron et al., 2000; Haspel and Libersat, 2003; and Nadolski, 2013). In Egypt, many studies had been conducted on *V. orientalis* focusing on ecology, biology, control and its dangerous effect on apiculture (Khater et al., 2001; AbdAl-Fattah and Ibrahim, 2009; and Taha, 2014).

In insects, the main body metabolites, viz., Carbohydrates, protein and lipids, have an essential role in different biological and physiological activities, such as body size, growth rate, fecundity and fertility (Fagan et al., 2002). Therefore, the contents of these macromolecules are good indicators of the level of metabolism in insects after treatment with chemicals or other materials (Zhu et al., 2012).

As reported by many authors (Hassan, 2002; Chapman, 2012; Cohen, 2010; and Sugumaran, 2010), proteins perform many functions within the living organisms, such as catalyzing metabolic reactions, replicating DNA, synthesis of ATP, the cell division, enzymes and hormones controlling many chemical reactions in the cell metabolism. In insects, protein metabolism plays a key role in rebuilding adult structures during the transformation of larvae/pupae into adults (Resmitha et al., 2014) as well as energy production and insect reproduction (Taşkin and Aksoylar, 2011).

Lipids represent an important source of energy, hormone precursors and structural members for insects and are transported from their synthesis site of storage via the haemolymph towards the user organs, in particular the vitellogenesis (Zhou and Miesfeld, 2009) and cuticular synthesis (Dapporto et al., 2008). Also lipids located in the egg play an important role in meeting the energy needs for developing embryo (Boz and Gülel, 2012). Quantity of lipids available for the reserves seems to be the result of a balance between the catch of food and the requests for reserves by processes, such as maintenance, growth and reproduction, and this balance is disturbed by any toxic product (Canavoso et al., 2001).

Carbohydrates play an important role in the structure and function of all tissues during metamorphosis as well as for the normal functioning of the male and female reproductive organs and embryonic development (cf. Chippendale, 1978). They increase during the rest periods, like metamorphosis, and decrease during the growth periods, like the stages of maturation of the gonads in insects (Bouaziz et al., 2011). On the other hand, the carbohydrate content in the haemolymph is an important indicator of the level of metabolism in insects, and a dynamic balance of the absorption, metabolism, and utilization by different tissues (Zhu et al., 2012). According to the currently available literature, many studies had been conducted for examining the toxicity of arthropod venoms against insects, but very little research attention had been paid to their effects on the physiological criteria. Therefore, the present study was conducted aiming at the estimation of disturbing effects of *L. quinquestriatus* venom, *V. orientalis* venom and Apitoxin of *A. mellifera* on the main body metabolites in larvae and pupae of *G. mellonella*.

## 2. Materials and methods

### 2.1. Experimental insect

A culture of the greater wax moth *G. mellonella* L. (Lepidoptera: Pyralidae) was maintained in the laboratory of Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt, under controlled conditions ( $27 \pm 2^\circ\text{C}$ ,  $65 \pm 5\%$  R.H., photoperiod 14 h L and 10 h D). This culture was originated by a sample of larvae kindly obtained from a culture of susceptible strain maintained for several generations in Plant Protection Unit, Desert Research Center, Cairo, Egypt. Larvae were transferred into glass containers, tightly covered with muslin cloth secured with rubber bands. After reviewing different techniques of the artificial diet described by some authors (Metwally et al., 2012; and Nitin et al., 2012), *G. mellonella* larvae in the present culture had been provided with an artificial diet as described by Bhatnagar and Bareth (2004). It contained maize flour (400 g), wheat flour, wheat bran and milk powder, 200 g of each. Also, the diet was provided with glycerol (400 g), bee honey (400 g), yeast (100 g). The full grown larvae metamorphosed into pupae. The resulting pupae were collected and transferred into clean jars provided with a layer of moistened saw dust on the bottom. Then, the emerged adult moths were kept in glass containers provided with white paper scraps, as oviposition sites. After mating, female moths were allowed to lay eggs. The egg patches were collected daily, and transferred into Petri dishes containing a layer of artificial diet for feeding of the hatching larvae.

### 2.2. Collection and preparation of arthropod venoms

#### 2.2.1. Scorpion collection and obtaining of venom

Sixty five adult individuals of the Death stalker scorpion *Leiurus quinquestriatus* Hemprich and Ehrenberg (Buthidae: Scorpiones: Arachnida) were collected from Garf Hessin at 23.289024N32.776828E, west of Nasser Lake, Aswan, Egypt, during October 2014. Scorpions were collected at daytime by random searching their hiding places, mostly under rocks and other favorable shelters (Williams, 1968). The collected specimens were



### 2.5. Statistical data analysis

Data obtained were analyzed by the Student's *t*-distribution, and refined by Bessel correction (Moroney, 1956) for the test significance of difference between means.

## 3. Results

### 3.1. Effects of arthropod venoms on the total protein content of *G. mellonella*

Depending on the data assorted in Table 1, the protein level in body tissues of control larvae gradually elevated along the larval stage ( $2.90 \pm 0.50$  and  $5.35 \pm 0.40$  mg/g in 5<sup>th</sup> instar and 7<sup>th</sup> instar larvae, respectively). On the other hand, the total protein content was drastically reduced in the treated larvae, regardless the tested venom or the larval instar. The strongest reducing potency was exhibited by Apitoxin in the 5<sup>th</sup> instar larvae (19.80% reduction) and exhibited by wasp venom in the 7<sup>th</sup> instar larvae (34.77% reduction). The least reducing potency was exhibited by scorpion venom (17.45 and 15.7% reductions, in 5<sup>th</sup> and 7<sup>th</sup> instar larvae, respectively).

**Table 1: Influenced total protein content (mg/g) in the larval tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Larval instar	
		5 <sup>th</sup>	7 <sup>th</sup>
<i>A. mellifera</i>	Mean ± SD	2.39 ± 1.33 d	3.77 ± 2.19 d
	Change (%)	-19.80	-29.53
<i>V. orientalis</i>	Mean ± SD	2.47 ± 0.24 d	3.49 ± 1.48 d
	Change (%)	-17.11	-34.77
<i>L. quinquestratus</i>	Mean ± SD	2.64 ± 0.03 d	4.51 ± 1.16 c
	Change (%)	-17.45	-15.7
Control	Mean ± SD	2.98 ± 0.50	5.35 ± 0.40

**Note:** Mean ± SD followed with c: highly significantly different ( $p < 0.01$ ). d: very highly significantly different ( $p < 0.001$ ).

With regard to the disturbance of protein content in the body tissues of the successfully developed pupae, data of Table 2 obviously revealed that the protein content as gradually declined throughout the pupal stage of controls ( $1.96 \pm 0.19$ ,  $1.42 \pm 1.41$  and  $0.99 \pm 0.31$  mg/g in early-, mid- and late-aged pupae, respectively). In general, all arthropod venoms predominantly prevented the pupae to attain normal protein level. In the early-aged pupae, the most powerful suppressing action on proteins was exerted by wasp venom while scorpion venom exhibited the least reducing potency. In the mid-aged pupae, the strongest reducing action was exhibited by the scorpion venom while the weakest reducing action was exerted by Apitoxin. In the late-aged pupae, Apitoxin had the strongest reducing potency on proteins but scorpion venom had the least reducing potency (46.46 and 4.04% reductions, respectively).

### 3.2. Effects of arthropod venoms on the total lipid content of *G. mellonella*

Depending on the data arranged in Table 3, the lipid content gradually decreased with instar of normal larvae ( $265.77 \pm 18.0$  and  $247.75$  mg/g, in 5<sup>th</sup> and 7<sup>th</sup> instars, respectively). Also, these data revealed a prevalent inducing effect of all arthropod venoms on larvae to gain more lipids than control congeners. In the 5<sup>th</sup> instar larvae, Apitoxin achieved the least enhancing action. In the 7<sup>th</sup> instar larvae, the least enhancing effect was achieved by scorpion venom on lipids.

In connection with the lipid content in the pupae, data listed in Table 4 revealed a gradual decreasing content with the age ( $151.65 \pm 65.0$ ,  $149.44 \pm 25.2$  and  $67.57 \pm 11.9$  mg/g, in early-, mid- and late-aged pupae, respectively). The lipid content in pupal tissues significantly increased after treatment of 3<sup>rd</sup> instar larvae with

**Table 2: Influenced total protein content (mg/g) in the pupal tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Pupal age		
		Early	Mid	Late
<i>A. mellifera</i>	Mean ± SD	1.64 ± 0.75 d	1.34 ± 1.21 a	0.53 ± 0.84 c
	Change (%)	-16.33	-5.63	-46.46
<i>V. orientalis</i>	Mean ± SD	1.49 ± 0.36 d	1.35 ± 0.44 a	0.91 ± 1.08 c
	Change (%)	-23.98	-4.93	-8.08
<i>L. quinquestriatus</i>	Mean ± SD	1.74 ± 1.43 d	1.31 ± 0.94 a	1.03 ± 1.55 c
	Change (%)	-11.22	-7.75	4.04
Control	Mean ± SD	1.96 ± 0.19	1.42 ± 1.49	0.99 ± 0.31

**Note:** Mean ± SD followed with a: insignificantly different ( $p > 0.05$ ). c, d: see footnote of Table (1).

**Table 3: Influenced total lipid content (mg/g) in the larval tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Larval instar	
		5 <sup>th</sup>	7 <sup>th</sup>
<i>A. mellifera</i>	Mean ± SD	345.35 ± 101.6 d	618.62 ± 148.3 c
	Change (%)	29.94	149.70
<i>V. orientalis</i>	Mean ± SD	358.86 ± 93.8 d	528.53 ± 131.8 c
	Change (%)	35.03	113.33
<i>L. quinquestriatus</i>	Mean ± SD	363.36 ± 71.6 d	414.41 ± 240.3 c
	Change (%)	36.72	67.27
Control	Mean ± SD	265.77 ± 18.0	247.75 ± 99.6

**Note:** c, d: see footnote of Table (1).

each of the tested venoms, regardless the pupal age. In other words, the tested venoms unexceptionally enhanced the pupae to gain more lipids than the control congeners. For some detail, Apitoxin exhibited the most potent inducing action but scorpion venom exhibited the least enhancing action on the early-aged pupae to gain more lipids (31.68 and 20.79% increments, respectively). With regard to mid-aged pupae, Apitoxin exhibited the strongest enhancing effect while the scorpion venom exhibited the least one (15.54 and 0.48% increments, respectively). The wasp venom had the highest promoting potency against lipids of late-aged pupae but scorpion venom had the least potency (102.22 and 51.1% increments, respectively).

**3.3. Effects of arthropod venoms on the total carbohydrate content of *G. mellonella***

According to the data distributed in Table 5, treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> value of each venom resulted in a dramatic reduction of carbohydrate content in body tissues of both 5<sup>th</sup> and 7<sup>th</sup> instar larvae. In regard of the normal larvae, carbohydrate content decreased with the instar (0.0366 ± 0.007 and 0.0314 ± 0.006 mg/g, in 5<sup>th</sup> and 7<sup>th</sup> instars, respectively). For some detail, the least reducing action on

carbohydrates in the 5<sup>th</sup> larval instar was exerted by the wasp venom while a similar action on carbohydrates in 7<sup>th</sup> larval instar was exerted by the scorpion venom (21.04 and 11.78% reductions, respectively).

**Table 4: Influenced total lipid content (mg/g) in the pupal tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Pupal age		
		Early	Mid	Late
<i>A. mellifera</i>	Mean ± SD	199.70 ± 2.6 c	172.67 ± 34.1 d	93.09 ± 13.0 d
	Change (%)	+31.68	+15.54	+37.77
<i>V. orientalis</i>	Mean ± SD	198.19 ± 11.9 c	162.16 ± 11.9 d	136.64 ± 60.0 d
	Change (%)	+30.69	+8.51	+102.22
<i>L. quinquestriatus</i>	Mean ± SD	183.18 ± 15.8 c	150.15 ± 6.9 d	102.10 ± 5.2 d
	Change (%)	+20.79	+0.48	+51.1
Control	Mean ± SD	151.65 ± 65	149.44 ± 25.2	67.57 ± 11.9

**Note:** c, d: see footnote of Table (1).

**Table 5: Influenced total carbohydrate content (mg/g) in the larval tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Larval instar	
		5 <sup>th</sup>	7 <sup>th</sup>
<i>A. mellifera</i>	Mean ± SD	0.0262 ± 0.004 d	0.0237 ± 0.006 d
	Change (%)	-28.42	-24.52
<i>V. orientalis</i>	Mean ± SD	0.0289 ± 0.007 d	0.0253 ± 0.004 d
	Change (%)	-21.04	-19.43
<i>L. quinquestriatus</i>	Mean ± SD	0.0284 ± 0.009 d	0.0277 ± 0.003 d
	Change (%)	-22.40	-11.78
Control	Mean ± SD	0.0366 ± 0.007	0.0314 ± 0.006

**Note:** d: see footnote of Table (1).

Data of the determined carbohydrate content in the body tissues of pupae were assorted in Table 6. In the light of these data, the carbohydrate content gradually decreased in normal pupae with the age (0.0125 ± 0.003, 0.0101 ± 0.003 and 0.0063 ± 0.002 mg/g in the early-, mid- and late-aged pupae, respectively). As clearly seen in the same table, the carbohydrate content in treated pupae had been drastically reduced, regardless the tested venom. In respect of the early-aged pupae, the strongest reducing effect was exhibited by wasp venom (27.20% carbohydrate reduction) while the least reducing effect was exhibited by Apitoxin (19.20% carbohydrate reduction). In the mid-aged pupae, wasp venom had the most potent reducing potency (49.50% carbohydrate reduction) but scorpion venom had the least reducing potency (6.93% carbohydrate reduction). Apitoxin could not affect the carbohydrate content in the body tissues of late-aged pupae while the wasp venom exhibited the most powerful reducing effect (46.30% carbohydrate reduction).



**Table 6: Influenced total carbohydrate content (mg/g) in the pupal tissues after treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with LC<sub>50</sub> values of certain arthropod venoms**

Venom		Pupal age		
		Early	Mid	Late
<i>A. mellifera</i>	Mean ± SD	0.0101 ± 0.001 d	0.0057 ± 0.002 c	0.0063 ± 0.008 c
	Change (%)	-19.20	-43.56	0.00
<i>V. orientalis</i>	Mean ± SD	0.0091 ± 0.001 d	0.0051 ± 0.007 c	0.0034 ± 0.001 c
	Change (%)	-27.20	-49.50	-46.30
<i>L. quinquestriatus</i>	Mean ± SD	0.0093 ± 0.001 d	0.0060 ± 0.001 c	0.0047 ± 0.002 c
	Change (%)	-25.60	-6.93	-25.40
Control	Mean ± SD	0.0125 ± 0.003	0.0101 ± 0.003	0.0063 ± 0.002

**Note:** c, d: see footnote of Table (1).

## 4. Discussion

### 4.1. Disturbed protein content in *G. mellonella* by arthropod venoms

As reported by some authors (Cohen, 2010; and Sugumaran, 2010), proteins perform many functions within the living organisms, such as catalyzing metabolic reactions, replicating DNA, synthesis of ATP, enzymes and hormones controlling many chemical reactions in the cell metabolism. Also, protein metabolism plays a key role in rebuilding adult structures during the transformation of larvae/pupae into adults in insects, (Resmitha et al., 2014) as well as energy production and insect reproduction (Tapkyn and Aksoylar, 2011). Proteins regulate and integrate several physiological and metabolic processes in the insect body through hormones, enzymes and nucleoproteins (Chapman, 2012).

After treatment of the 3<sup>rd</sup> instar larvae of *G. mellonella* with venoms of *L. quinquestriatus* venom, *V. orientalis* venom or Apitoxin of *A. mellifera*, the total protein contents in both 5<sup>th</sup> and 7<sup>th</sup> instar larvae had been drastically reduced. Also, all venoms predominantly prevented the successfully developed pupae to attain normal protein level, since remarkably reduced protein content was estimated. This reduction of protein content in larvae and pupae of *G. mellonella*, in the present study, after larval treatment with the tested arthropod venoms can be interpreted in the light of some conceivable suggestions, as follows. (1) It was suggested that the protein plays a major role in synthesis of the microsomal detoxifying enzymes against toxicants (foreign compounds) entering into the insect body (Kyung and Kim, 1990; and Hassan, 2002). In the present study, reduction of proteins might be due to their binding with the tested arthropod venoms and therefore might reflect the depressed activity of the detoxifying enzymes. (2) The toxins' stress can inhibit the total proteins owing to the breakdown of protein into amino acids, so with the entrance of these amino acids to TCA cycle as a keto acid (Schoonhoven, 1982), they will help to supply energy for the insect (Etebari and Matindoost, 2004). Therefore, protein reduction in the tissues of *G. mellonella* might play a role in compensatory mechanisms under the present toxins' stresses, to provide intermediates to the Krebs cycle (Nath et al., 1997). (3) The protein reduction in the current study might, also, be due to the interference of the tested arthropod venoms with the insect endocrine system causing a hormonal imbalance (Sugumaran, 2010) and affecting the general metabolism (De Mark and Bennett, 1989) or protein synthesis, in particular (Padmaja and Rao, 2000). The tested *L. quinquestriatus* venom, *V. orientalis* venom or Apitoxin of *A. mellifera* might either act on the hormonal regulation of the protein synthesis, degradation and inhibition or act on the neurosecretory cells which control endocrine organs (Bouaziz et al., 2011; and Djeghader et al., 2014). Further investigation should be conducted in the foreseeable future for good understanding the modes of action of some chemical constituents of the tested venoms, in the present investigation.



#### 4.2. Disturbed lipid content in *G. mellonella* by arthropod venoms

Lipids represent an important source of energy, hormone precursors and structural members for insects and are transported from their synthesis site of storage via the haemolymph towards the user organs (Dapporto et al., 2008; and Zhou and Miesfeld, 2009). In the current study, all venoms of *L. quinquestriatus*, *V. orientalis* and *A. mellifera* exhibited prevalent enhancing effects on larvae to gain more lipids than control larvae of both 5<sup>th</sup> and 7<sup>th</sup> instars. Also, all tested venoms unexceptionally induced the pupae to gain more lipids than the control congeners. This result of enhanced lipids disagreed with many results of reduced lipid content in different insect species after treatment with various toxic compounds. On the other hand, the present result was, in some extent, in agreement with the reported results of induced lipids in various insect species after larval treatment with some insect growth regulators (Ghoneim, 1994; Soltani-Mazouni et al., 1999; and Bouaziz et al., 2011).

To interpret the enhancement of lipids in larvae and pupae of *G. mellonella*, in the current work, it is important to point out that the influenced synthesis of lipids in insects has been resulted in disruptively affected physiology and subsequently deranged vital functions of growth and reproduction. The balance between quantity of lipids available for the reserves and the requests for these reserves can be disturbed by any toxic material (Canavoso et al., 2001). Therefore, induction of lipids in larvae and pupae of *G. mellonella* by the tested arthropod venoms might be due to their interference with not only the lipid synthesis but also the lipid mobilization as promoted to convert into other metabolites or fatty acids. Also, the increase of lipids might be due to the accumulation of carbohydrates which might lead to an inverse in their conversion rate to lipids as a reverse material. However, the exact interpretation of lipid induction in larvae and pupae of *G. mellonella*, in the present study, remains speculative.

#### 4.3. Disturbed carbohydrate content in *G. mellonella* by arthropod venoms

Carbohydrates play an important role in the structure and function of all tissues during metamorphosis in insects. Also, carbohydrates are prerequisite metabolites for the normal functioning of male and female reproductive organs and embryonic development (cf. Chippendale, 1978). In general, carbohydrates play a key role in the physiology of those insects subjected to foreign toxins (Kaufmann and Brown, 2008). In the present investigation, treatment of the 3<sup>rd</sup> instar larvae with each of the tested venoms, viz., scorpion *L. quinquestriatus*, wasp *V. orientalis* and *A. mellifera*, resulted in dramatic reduction of carbohydrate content in tissues of 5<sup>th</sup> and 7<sup>th</sup> instar larvae. Also, carbohydrate content in treated pupae had been drastically reduced, regardless the tested venom or the pupal age.

The production or utilization of the main body metabolites, such as carbohydrates, is controlled by juvenile hormone or related to various hormonal organs and neurosecretion (Gade et al., 1997; Gade, 2004; and Sugumaran, 2010). Thus, the prevalent reduction of carbohydrate content in larvae and pupae of *G. mellonella*, in the present study, might be due to interference of the tested arthropod venoms with the hormonal regulation of carbohydrate metabolism (Imboden and Luscher, 1976) or to their effects on the carboxylase activity (Mukherjee and Sharma, 1996). Also, it is suggested that this carbohydrate depletion might be due to utilization of the reserved glucose sources of the larval tissues as a result of venoms' stresses (Sharma et al., 2011). In addition, the present result of carbohydrate reduction in *G. mellonella* might be due to stresses of the tested arthropod venoms on proteins to be degraded into amino acids to take part in the TCA cycle of acetic acid. Subsequently, the carbohydrate metabolic functions to make up for the lower energy were altered under the stress of present toxic materials in the body of *G. mellonella* (Nath et al., 1997; Nath, 2000). In general, detoxification in the larvae required a larger portion of consumed substances to be transformed into energy after arthropod venom treatments, in the current investigation, which might be another reason for the reduction in both proteins and carbohydrates in *G. mellonella* larvae (Xu et al., 2016).

Finally, it is important to mention that the disturbing effects of the tested arthropod venoms on the main metabolites in *G. mellonella*, in the present study, should be achieved by certain constituent (s) in each of these venoms; since venom of the scorpion *L. quinquestriatus* contains different compounds, such as polypeptide neurotoxins and basic proteins with low molecular weight bound to disulphide bridges especially (Hamon et al., 2002; and Kopygan et al., 2006); venom of the wasp *V. orientalis* contains a mixture of proteins and chemical constituents, such as serotonin, adrenaline, nor-adrenaline, dopamine, kinins and histidine (Biggs et al., 2007; and Klochkov et al., 2008); and Apitoxin of *A. mellifera* contains a complex mixture of proteins, peptides, and

low molecular components. The chief components are apamin, melittin and phospholipase A<sub>2</sub> (Khajehpour et al., 2004; and Bogdanov, 2012).

## 5. Conclusion

Depending on the present results, scorpion *L. quinquestriatus* venom, wasp *V. orientalis* venom and Apitoxin of *A. mellifera* caused reduction in proteins and carbohydrates but induced lipids, in the tissues of larvae and pupae of *G. mellonella*. Reduction of proteins might reflect the depressed activity of the detoxifying enzymes in this insect. The accumulated lipids in these tissues indicated a disruption of the balance between quantity of available lipids for the reserves and the requests for these reserves by processes, which caused by the tested venoms. Thus, these venoms seriously disturbed the main body metabolism. Therefore, the tested arthropod venoms may be good control agents, in an integrated management program against the present pest, *G. mellonella*. However, novel delivery systems should be explored in order to the insectotoxins can find their way into commercial applications in foreseeable future.

## References

- Abbasipour, H., Mahmoudvand, M., Deylami, A. and Hosseinpour, M.H. (2009). Fumigant toxicity of essential oils of *Rosemarinus officinalis* L. and *Eucalyptus camodulensis* Deh. against some stored products pests, Proceeding of the 6<sup>th</sup> Asia-Pasific Congress of Entomology, Entomology in Health, Agriculture and Environment, Beijing, China.
- AbdAl-Fattah, A.M.A. and Ibrahim, Y.Y. (2009). The serious effects of the dangerous insect predator (*Vespa orientalis* L.) on honey bee colonies in Giza Governotate, Egypt. The 4<sup>th</sup> Conference on Recent Technologies in Agriculture, Faculty of agriculture, Cairo University, 9 September, 2009.
- Abdelaal, A.A.A. and El-defrawy, B.M. (2014). Efficacy of new designed traps for controlling the oriental hornet (*Vespa orientalis*) in Egyptian apiaries and its measurements. *International Journal of Advanced Research*. 2(10), 1-8.
- Akyol, E., Yeninar, H., ahinler, N. and Ceylan, D.A. (2009). The using of carbon dioxide (CO<sub>2</sub>) on controlling of the greater wax moth's *Galleria mellonella* L. (Lepidoptera: Pyralidae) damages. *Uludag Bee Journal*. 9(1), 26-31.
- Altstein, M. (2004). Novel insect control agents based on neuropeptide antagonists: the PK/PBAN family as a case study. *Journal of Molecular Neuroscience*. 22 (1–2), 147–157. doi: 10.1385/JMN:22:1-2:147.
- Altstein, M., Ben-Aziz, O., Scheffler, I., Zeltser, I. and Gilon, C. (2000). Advances in the application of neuropeptides in insect control. *Crop Protection*. 19 (8–10), 547–555. [https://doi.org/10.1016/S0261-2194\(00\)00071-5](https://doi.org/10.1016/S0261-2194(00)00071-5)
- Archer, M.E. (1998). Taxonomy, distribution and nesting biology of *Vespa orientalis* L. (Hym., Vespidae). *Entomologist's Monthly Magazine*. 134, 45–51.
- Armendariz, I., Downes, M.J. and Griffin, C.T. (2002). Effect of timber condition on parasitization of pine weevil (*Hylobius abietis* L.) larvae by entomopathogenic nematodes under laboratory conditions. *Biocontrol Science and Technology*. 12, 225-233. DOI: 10.1080/09583150120124487
- Awasthi, P. and Sharma, P. (2012). Docking Study of synthesized juvenile hormone analogues as an Insect Growth Regulators. 2012 UK Sim 14<sup>th</sup> International Conference on Computer Modelling and Simulation. Cambridge, Cambridgeshire United Kingdom, March 28-March 30.
- Awasthi, P. and Sharma, P. (2013). Designing and binding mode prediction of juvenile hormone analogues as potential inhibitor for *Galleria mellonella*. *Journal of Computer Science and Systems Biology*. 6(3), 106-111. DOI: 10.4172/jcsb.1000107
- Azam, N.K., Ahmed, N., Biswas, S., Ara, N., Rahman, M., Hirashima, A. and Hasan, N. (2019). A review on bioactivities of honey bee venom. *Annual Research & Review in Biology*. 30(2), 1-13. DOI: 10.9734/ARRB/2018/45028
- Badry, A.; Younes, M.; Sarhan, M. M. and Saleh, M. (2018): On the scorpion fauna of Egypt, with an identification key (Arachnida: Scorpiones). *Zoology in the Middle East*. 64(1), 75-87. <https://doi.org/10.1080/09397140.2017.1414976>

- Basedow, T., Shafie, H., Abo-El-Saad, M. and Al Ajlan, A. (2012). Evaluation of *Bacillus thuringiensis* aizawi and neem for controlling the larvae of the greater wax moth, *Galleria mellonella* (Lepidoptera: Pyralidae). *International Journal of Agriculture and Biology*. 14, 60-63.
- Bhatnagar, A. and Bareth, S.S. (2004). Development of low cost, high quality diet for greater wax moth, *Galleria mellonella* (Linnaeus). *Indian Journal of Entomology*. 66(3), 251-255.
- Biggs, J.S., Rosenfeld, Y., Shai, Y., Olivera, B.M. and Conolysin, M.T. (2007). Conolysin-Mt: a conus peptide that disrupts cellular membranes. *Biochemistry*. 46, 12586-12593. DOI: 10.1021/bi700775p
- Bogdanov, S. (2012): Bee venom: composition, health, medicine: a review. *Journal of Peptides*. (1), 1-20.
- Bouaziz, A., Boudjelida, H. and Soltani, N. (2011). Toxicity and perturbation of the metabolite contents by a chitin synthesis inhibitor in the mosquito larvae of *Culiseta longiareolata*. *Annals of Biological Research*. 2(3), 134-143.
- Boz, A. and Gülel, A. (2012). The effects of temperature and time after parasitization on total amount of protein, lipid and carbohydrate in hemolymph of host larvae, *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae). *Turkish Entomology Journal*. 36(2), 239-247.
- Canavoso, L.E.; Jouni, Z.E.; Karnas, K.J.; Pennington, J.E. and Wells, M.A. (2001): Fat metabolism in insects. *Annual Review of Nutrition*. 21: 23-46. doi: 10.1146/annurev.nutr.21.1.23
- Cao, Z., Yu, Y. and Li, W. (2013). The genome of *Mesobuthus martensii* reveals a unique adaptation model of arthropods. *Nature Communications*. 4, 2602.
- Carpenter, J.M. and Kojima, J. (1997). Checklist of the species in the subfamily Vespinae (Insecta: Hymenoptera: Vespidae). *Natural History Bulletin of Ibaraki University*. 1, 51-92.
- Carpenter, J., Bloem, S. and Marec, F. (2005). Inherited sterility in insects. In: "Sterile Insect Technique" (Dyck, V.A., Hendrichs, J., Robinson, A.S., eds.). pp: 115-146.
- Chandel, Y.S., Sharma, S. and Verma, K.S. (2003). Comparative biology of the greater wax moth, *Galleria mellonella* L. and lesser wax moth, *Achoria grisella*. *Forest Pest Management and Economic Zoology*. 11, 69-74.
- Chapman, R.F. (2012). The insects: structure and function, 5th ed. Cambridge University Press, Cambridge, United Kingdom, 595pp.
- Chippendale G.M. (1978). The functions of carbohydrates in insect life processes. In: "Biochemistry of Insects" (Rockstein, M., ed.). Academic Press, New York, pp: 2-54.
- Christen, J.M., Campbell, J.F., Zurek, L., Shapiro-Ilan, D.I., Lewis, E.E., Ramaswamy, S.B. (2008). Role of symbiotic and non-symbiotic bacteria in carbon dioxide production from hosts infected with *Steinernema riobrave*. *Journal of Invertebrate Pathology*. 99(1), 35-42. doi: 10.1016/j.jip.2008.05.008.
- Cohen, E. (2010). Chitin biochemistry: synthesis, hydrolysis and inhibition. *Advances in Insect Physiology*. 38, 5-74. [https://doi.org/10.1016/S0065-2806\(10\)38005-2](https://doi.org/10.1016/S0065-2806(10)38005-2)
- Coudron, T.A., Wright, M.M., Puttler, B., Brandt, S.L. and Rice, W.C. (2000). Effect of the ectoparasite *Necremnus breviramulus* (Hymenoptera: Eulophidae) and its venom on natural and factitious hosts. *Annals of Entomological Society of America*. 93(4), 890-897. [https://doi.org/10.1603/0013-8746\(2000\)093\[0890:EOTENB\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2000)093[0890:EOTENB]2.0.CO;2)
- Cruz-Landim, C. and Abdalla, F.C. (2002). Bees exocrine glands. FUNPEC-Publisher, 194pp.
- Dahlman, D.L., Rana, R.L., Schepers, E.J., Schepers, T., Diluna, F.A. and Webb, B.A. (2003). A teratocyte gene from a parasitic wasp that is associated with inhibition of insect growth and development inhibits host protein synthesis. *Insect Molecular Biology*. 12, 527-534. doi: 10.1046/j.1365-2583.2003.00439.x.
- Dantas, C.G., Nunes, T.L.G.M. and Gomes, M. (2013). Apitoxina: coleta, composição química, propriedades biológicas e atividades terapêuticas. *Revista Ibero-Americana de Ciências Ambientais*. 4(2), 127-150. DOI: <https://doi.org/10.6008/ESS2179-6858.2013.002.0009>
- Dapporto, L., Lambardi, D. and Turillazzi, S. (2008). Not only cuticular lipids: first evidence of differences between foundresses and their daughters in polar substances in the paper wasp *Polistes dominulus*. *Journal of Insect Physiology*, 54, 89-95. doi: 10.1016/j.jinsphys.2007.08.005.

- De Mark, I.J. and Bennett, G.W. (1989). Efficacy of chitin synthesis inhibitors on nymphal German cockroaches (*Dictyoptera: Blattidae*). *Journal of Economic Entomology*. 82, 1633-1637. DOI: 10.1093/jee/82.6.1633
- Dindo, M.L., Verdinelli, M., Baronio, P., Serra, G.E. (2001). Laboratory and field performance of *in vitro* and *in vivo* reared *Exorista larvarum* (L.), a natural enemy of cork oak defoliators. In: "Integrated protection in oak forests" (Villemant, C. and Sousa, E., eds.). Proceedings of the IUBC-WPRS working group at Oeiras Lisbon, Portugal, 01st-04<sup>th</sup> Oct., 2001. *Bulletin Oib-Srop*, 25, 147-150.
- Djehghader, N.E.H., Aïssaoui, L., Amira, K. and Boudjelida, H. (2014). Impact of a chitin synthesis inhibitor, Novaluron, on the development and the reproductive performance of mosquito *Culex pipiens*. *World Applied Sciences Journal*. 29(7), 954-960. DOI: 10.5829/idosi.wasj.2014.29.07.82190
- Durmu<sup>o</sup>, Y. and Büyükgüzel, K. (2008). Biological and immune response of *Galleria mellonella* (Lepidoptera: Pyralidae) to sodium tetraborate. *Journal of Economic Entomology*. 101(3), 777-783. DOI:10.1603/0022-0493(2008)101[777:BAIROG]2.0.CO;2
- Dvorak, L. (2006). Oriental Hornet *Vespa orientalis* Linnaeus, 1771 found in Mexico (Hymenoptera, Vespidae, Vespinae). *Entomological Problems*, 36(1), 80.
- Ebadi, R., Jafari, R., Majd, F., Tahmasbi, G. and Zolphagharieh, H. (2001). Effect of gamma-ray male sterilization on the integrated control management of greater wax moth, *Galleria mellonella* L. (Lep., Pyralidae). *JWSS - Isfahan University of Technology*. 5(3), 191-199.
- Elbehery, H., Abd El-Wahab, T.E. and Dimetry, N.Z. (2016). Management of the greater wax moth *Galleria mellonella* with Neemazal-T/S, in the laboratory and under semi-field conditions. *Journal of Apiculture Science*. 60(2), 69-76. DOI: 10.1515/jas-2016-0018
- El-Hennawy, H.K. (1987). A simplified key to Egyptian scorpion species (Arachnida: Scorpionida). *Serket*, 1, 15-17.
- El-Kholy, E.M.S. and Mikhael, A.A. (2008). Scanning electron microscopy on the male antennae of the greater wax moth, *Galleria mellonella* (L.), treated with gamma radiation. *Isotope and Radiation Research*. 40(3), 603-613.
- Ellis, J.D., Graham, J.R. and Mortensen, A. (2013). Standard methods for wax moth research. *Journal of Apiculture Research*. 52, 1–17. <https://doi.org/10.3896/IBRA.1.52.1.10>
- Er, A., Tapkýran, D. and Sak, O. (2017). Azadirachtin-induced effects on various life history traits and cellular immune reactions of *Galleria mellonella* (Lepidoptera: Pyralidae). *Archives of Biological Sciences*. 69(2), 335-344. <https://doi.org/10.2298/ABS160421108E>.
- Etebari, K. and Matindoost, L. (2004). The study on effects of larval age and starvation stress on biochemical macromolecules abundance of haemolymph in silkworm *Bombyx mori*. In: Proceedings of the Sixteenth Iranian Plant Protection Congress, General Entomology Symposium, August 28–September 1, University of Tabriz, Iran, p. 435.
- Fabiano, G., Pezzolla, A., Filograna, M.A. and Ferrarese, F. (2008). Traumatic shock-physiopathologic aspects. *Giornale di chirurgia*, 29, 51-57.
- Fagan, W.F., Siemann, E., Mitter, C., Denno, R.F., Huberty, A.F., Woods, H.A. and Elser, J.J. (2002). Nitrogen in insects: implications for trophic complexity and species diversification. *American Naturalist*. 160, 784-802. doi:10.1086/343879
- Folch, J., Less, M. and Sloane-Stanley, G.H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*. 26, 497-509.
- Friedman, J. and Ishay, J.S. (1987). Inhibition of protein synthesis by an extract of the venom sac of the oriental hornet (*Vespa orientalis*). *Toxicon*. 25(6), 673-676. [https://doi.org/10.1016/0041-0101\(87\)90114-0](https://doi.org/10.1016/0041-0101(87)90114-0)
- Froy, O., Zilberberg, N., Chejanovsky, N., Anglister, J., Loret, E.P., Shaanan, B., Gordon, D. and Gurevitz, M. (2000). Scorpion neurotoxins: structure/function relationships and application in agriculture. *Pest Management Science*. 56, 472–474. DOI: 10.1002/(SICI)1526-4998(200005)56:5<472::AID-PS148>3.3.CO;2-6



- Gade, G. (2004). Regulation of intermediary metabolism and water balance of insects by neuropeptides. *Annual Review of Entomology*. 49, 93-113. doi: 10.1146/annurev.ento.49.061802.123354.
- Gade, G., Hoffmann, K.H. and Spring, J.H. (1997). Hormonal regulation in insects: facts, gaps, and future directions. *Physiological Review*. 77, 963-1032. doi: 10.1152/physrev.1997.77.4.963.
- George, J., Devi, G. and Bhattacharyya, B. (2019). Survival and infectivity of entomopathogenic nematode *Oscheius rugaensis* in different formulations against wax moth, *Galleria mellonella*. *Journal of Entomology and Zoology Studies*. 7 (3), 241-244.
- Ghoneim, K.S. (1994). Synergistic and antagonistic action of Chlorfluazuron and mevalonic acid against the main body metabolism of the Egyptian cotton leafworm *Spodoptera littoralis* Boisd. (Lepidoptera: Noctuidae). *Journal of Egyptian German Society of Zoology*. 14(D), 89-115.
- Ghoneim, K., Hamadah, Kh., Tanani, M., Abdel-Khaliq, A. and Emam, D. (2019a). Toxicity and disruptive impacts of the honeybee Apitoxin on growth and development of the greater wax moth, *Galleria mellonella* (Lepidoptera: Pyralidae). *Egyptian Academic Journal of Biological Sciences (F. Toxicology & Pest Control)*. 11(2), 97-106.
- Ghoneim, K., Tanani, M., Hamadah, Kh., Abdel-Khaliq, A. and Emam, D. (2019b). Deteriorated adult performance and reproduction of the greater wax moth *Galleria mellonella* (Lepidoptera: Pyralidae) by the honey bee Apitoxin. *Egyptian Academic Journal of Biological Sciences (A. Entomology)*. 12(4), 95-108.
- Gurevitz, M. (2010). A deadly scorpion provides a safe pesticide. *Science News*, <http://www.sciencedaily.com>
- Hamon, A., Gilles, N., Sautiere, P., Kopeyan, C., Ulens, C., Tytgat, J., Lancelin, J. and Gordon, D. (2002). Characterization of Scorpion a-like toxin group using two new toxins from the Scorpion *Leiurus quinquestriatus* Hebraeus. *European Journal of Biochemistry*. 269(16), 3920-3933. doi: 10.1046/j.1432-1033.2002.03065.x.
- Harrison, R.L. and Bonning, B.C. (2000). Use of scorpion neurotoxins to improve the insecticidal activity of *Rachiplusia ou* multicapsid nucleopolyhedrovirus. *Biological Control*. 17(2), 191-201. <https://doi.org/10.1006/bcon.1999.0792>
- Haspel, G. and Libersat, F. (2003). Wasp venom blocks central cholinergic synapses to induce transient paralysis in cockroach prey. *Journal of Neurobiology*, 54(4), 628-637. doi: 10.1002/neu.10195.
- Hassan, H.A. (2002). Biological and biochemical studies on the effect of some botanical extracts on cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). M.Sc. Thesis, Faculty of Science, Ain Shams University, Egypt.
- Hussaini, S.S. (2003). Progress of research work on entomopathogenic nematodes in India. In: "Current status of research entomopathogenic Nematodes in India (Hussaini, S.S., Rabindra, R.J. and Nagesh, M., eds.). PdbC, publication, Bangalore, India, pp. 27-69.
- Imboden, H. and Luscher, M. (1976). Allatectomy in adult worker *Apis mellifera* (Hym., Apidae). *Review of Suisse Zoology*. 82, 964-989.
- Izzetoglu, S. and Karacali, S. (2003). The effects of 20-hydroxyecdysone on haemocytes of *Galleria mellonella* (Lepidoptera) *in vitro* conditions. *Gazi University Journal of Science (Turkey)*, 16(2), 233-238.
- James, R.R. (2011). Potential of ozone as a fumigant to control pests in honey bee (Hymenoptera: Apidae) hives. *Journal of Economic Entomology* 104, 353-359. doi: 10.1603/ec10385.
- Joseph, B. and George, J. (2012). Scorpion toxins and its Applications. *International Journal of Toxicological and Pharmacological Research*. 4(3), 57-61.
- Kaufmann, C. and Brown, M.R. (2008). Regulation of carbohydrate metabolism and flight performance by a hypertrehalosaemic hormone in the mosquito *Anopheles gambiae*. *Journal of Insect Physiology*. 54, 367-377. doi: 10.1016/j.jinsphys.2007.10.007
- Khajehpour, M., Troxler, T., Nanda, V. and Vanderkooi, J.M. (2004): Melittin as model system for probing interactions between proteins and cyclodextrins. *Proteins: Structure, Function and Bioinformatics*. 55, 275-287. DOI: 10.1002/prot.20036

- Khater, A.M., Ebada, I.M.A. and Khalil, I. (2001). The seasonal activity of oriental wasps, *Vespa orientalis* L. populations attacking honey bee colonies. *Arab Universities Journal of Agricultural Sciences, Ain Shams University, Cairo*, 9, 447-455.
- Khodairy, M.M. and Awad, A.A. (2013). A study on the sensory structure, in relation to some behavioral ecology of the oriental hornet (*Vespa orientalis* L.) (Hymenoptera: Vespidae). *Life Science Journal*. 10(2), 1207-1215.
- Klochkov, S.G., Pikhteleev, A.R. and Vskii, V.I.K. (2008). A new peptide from venom of the oriental hornet *Vespa orientalis*. *Mass Spectrometric Chemistry of Natural Compounds*. 44(1), 44-55.
- Knight, J.A., Anderson, S. and Jams, M.R. (1972). Chemical basis of the sulfo vanillin reaction of estimating total serum lipid. *Journal of Clinical Chemistry*. 18(3), 199-202. <https://doi.org/10.1093/clinchem/18.3.199>
- Kwadha, C.A., Ong'amo, G.O., Ndegwa, P.N., Raina, S.K. and Fombong, A.T. (2017). The biology and control of the greater wax moth, *Galleria mellonella*. *Insects*, 8, 61. doi: 10.3390/insects8020061
- Kyung, Y.H. and Kim, H.R. (1990). Characterization, of haemolymph protein from *Hyphantria cunea* (Drwry.). *The Korean Journal of Entomology*. 20(4), 239-246.
- Leng, P., Zhang, Z.; Pan, G. and Zhao, M. (2011). Applications and development trends in biopesticides. *African Journal of Biotechnology*. 10, 19864-19873. DOI: 10.5897/AJBX11.009
- Libersat, F., Haspel, G., Casagrand, J. and Fouad, K. (1999). Localization of the site of effect of a wasp's venom in the cockroach escape circuitry. *Journal of Comparative Physiology A*. 184, 333-345. DOI: 10.1007/s003590050331
- Ma, J. and Shi, Y.B. (2014). The *Mesobuthus martensii* genome reveals the molecular diversity of scorpion toxins. *Cell & Bioscience*. 4, 1. doi:10.1186/2045-3701-4-1
- Mahgoub, M.O., Lau, W.H., Bin Omar, D. and El Naim, A.M. (2018). Evaluation the toxicity of honey bee venom on *Achroia grisella* developmental stages. *World Journal of Agricultural Research*. 6(1), 5-9. doi: 10.12691/wjar-6-1-2.
- Mahmoudvand, M., Abbasipour, H., Hosseinpour, M.H., Rastegar, F. and Basij, M. (2011). Using some plant essential oils as natural fumigants against adults of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *Munis Entomology & Zoology*. 6(1), 150-154.
- Metwally, H.M.S., Hafez, G.A., Hussein, M.A., Salem, H.A. and Saleh, M.M.E. (2012). Low cost artificial diet for rearing the greater wax moth, *Galleria mellonella* L. (Lepidoptera: Pyralidae) as a host for entomopathogenic nematodes. *Egyptian Journal of Biological Pest Control*. 22(1), 15-22. <http://www.esbcp.org/index.asp>
- Mohamed, H.F., El-Naggar, S.E.M., Elbarky, N.M., Ibrahim, A.A. and Salama, M.S. (2014). The impact of each of the essential oils of marjoram and lemon grass in conjunction with Gamma irradiation against the greater wax moth, *Galleria mellonella*. *IOSR Journal of Pharmacy and Biological Sciences*. 9(5), 92-106. DOI: 10.9790/3008-095492106
- Moroney, M.J. (1956). Facts from figures (3<sup>rd</sup> ed.). Penguin Books Ltd., Harmondsworth. Middle Sex.
- Mukherjee, S.N. and Sharma, R.N. (1996). Azadirachtin induced changes in feeding, dietary utilization and midgut carboxylesterase activity of the final instar larvae of *Spodoptera litura* (Fabricius)(Lepidoptera: Noctuidae). *Journal of Environmental Science and Health*. B31, 1307-1319. <https://doi.org/10.1080/03601239609373069>
- Nadolski, J. (2013). Effects of the European hornet (*Vespa crabro* Linnaeus 1761) crude venom on its own species. *Journal of Venomous Animals and Toxins Including Tropical Diseases*, 19, 4. doi:10.1186/1678-9199-19-4.
- Nakasu, E.Y.T., Williamson, S.M., Edwards, M.G., Fitches, E.C., Gatehouse, J.A. and Wright, G.A. (2014). Novel biopesticide based on a spider venom peptide shows no adverse effects on honeybees. *Proceeding of Biological Science*. 281, 1787 doi: 10.1098/rspb.2014.0619
- Nath, B.S. (2000). Changes in carbohydrate metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* L., exposed to organophosphorus insecticides. *Pesticide Biochemistry and Physiology*. 68, 127-137. <https://doi.org/10.1006/pest.2000.2509>

- Nath, B.S., Suresh, A., Varma, B.M. and Kumar, R.S. (1997). Changes in protein metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* (Lepidoptera: Bombycidae) in response to organophosphorus insecticides toxicity. *Ecotoxicology and Environmental Safety*. 36, 169–173. DOI: 10.1006/eesa.1996.1504
- Neuman, M.G., Cotariu, D., Eshchar, J., Barr-Nea, L. and Ishay, J.S. (1987). Liver damage induced by Oriental hornet venom sac extract at the level of subcellular fractions. *Clinical Biochemistry*. 20(2), 85-90. DOI: 10.1016/S0009-9120(87)80105-4
- Nicholson, G.M. (2006). Spider venom peptides. In: *The Handbook of Biologically Active Peptides* (Kastin, A.J., ed.). Elsevier, San Diego, CA, pp: 389–399.
- Nitin, K., Kumar, K.D., Kumar, M.V. and Sanjay, P. (2012). Effect of economical modification in artificial diet of greater wax moth *Galleria mellonella* (Lepidoptera: Pyralidae). *Indian Journal of Entomology*, 74(4), 369-374.
- Olivera, B.M. (2002). Conus venom peptides: reflections from the biology of clades and species. *Annual Review of Ecology and Systematics*. 33, 25-47. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150424>
- Owayss, A.A. and Abd-Elgayed, A.A. (2007). Potential efficacy of certain plant volatile oils and chemicals against greater wax moth *Galleria mellonella* L. (Lepidoptera: Pyralidae). *Bulletin of Entomological Society of Egypt (Economic Series)*. 33, 67-75.
- Padmaja, P.G. and Rao, P.J. (2000). Effect of plant oils on the total haemocyte count (THC) of final instar larvae of *Helicoverpa armigera* Hübner. *Pesticide Research Journal*. 12(1), 112-116.
- Pamita, A. and Priyanka, S. (2013). Designing and binding mode prediction of juvenile hormone analogues as potential inhibitor for *Galleria mellonella*. *Journal of Computer Science and Systems Biology*. 6, 106-111. DOI: 10.4172/jcsb.1000107
- Piek, T. (1987). Synergistic effects of agonists and antagonists in insect venoms-a natural way of insecticidal action. *Pesticide Science*. 19, 317-22.
- Possani, L.D., Merino, E., Corona, M., Bolivar, F. and Becerril, B. (2000). Peptides and genes coding for scorpion toxins that affect ion-channels. *Biochimie*. 82(9), 861-8. [http://dx.doi.org/10.1016/S0300-9084\(00\)01167-6](http://dx.doi.org/10.1016/S0300-9084(00)01167-6)
- Quistad, G.B., Skinner, W.S. and Schooley, D.A. (1988). Venoms of social hymenoptera- toxicity to the lepidopteran, *Manduca sexta*. *Journal of Insect Biochemistry*. 18(6), 511-514. [https://doi.org/10.1016/0020-1790\(88\)90001-7](https://doi.org/10.1016/0020-1790(88)90001-7)
- Resmitha, C., Reshma, R.M., Punathumpambath, B. and Vadakkadath Meethal, K. (2014). The ecdysone mimic, methoxyfenozide, alters the level of major haemolymph proteins in the larvae of *Spodoptera mauritia* Boisdu. (Lepidoptera: Noctuidae). *Acta Biologica Indica*. 3(2), 726-730.
- Ross, D.C., Herzog, G.A., Brady, L.J.E. and Joe, W.C. (1987). Differential effects of *Vespula* wasp venom and its boll components, mastoparan, and bradykinin: toxicity and feeding tests with the cotton worm. *Comparative Biochemical Physiology*. 89C(2), 299-303.
- Sak, O. and Uckan, F. (2009). Effects of cypermethrin on the pupation and mortality of *Galleria mellonella* L. (Lepidoptera: Pyralidae). *Uludag Bee Journal*. 9(3), 88-96.
- Saleh, M., Younes, M., Badry, A. and Sarhan, M. (2017). Zoogeographical analysis of the Egyptian scorpion fauna. *Al Azhar Bulletin of Science*. 28(1), 1- 14.
- Sarhan, M., Maged, M.F., Alaa, M.H., Hamdy, A.M. and Ahmed, B.S. (2012). Variation of protein profile among consecutive stings of the scorpion *Parabuthus leiosoma* (Family: Buthidae) from Egypt, supports the venom-metering hypothesis in scorpions. *Al-Azhar Bull. Sci.*, 23(1), 61-71.
- Schoonhoven, L.M. (1982). Biological aspects of antifeedants. *Entomologia Experimentalis et Applicata*. 31, 57-89. <https://doi.org/10.1111/j.1570-7458.1982.tb03119.x>
- Sharma, P., Mohan, L., Kumar, K.D. and Srivastava, C.N. (2011). Status of carbohydrate, protein and lipid profile in the mosquito larvae treated with certain phytoextracts. *Asian Pacific Journal of Tropical Medicine*. 4, 301-304. DOI: 10.1016/S1995-7645(11)60090-4



- Singh, N.B. and Sinha, R.N. (1977). Carbohydrate, lipid and protein in the development stages of *Sitopholes orzae* and *Sitophilus granarius*. *Annals of Entomological Society of America*. 70, 107-111.
- Soltani-Mazouni, N., Khebbeb, M.E.H. and Soltani, N. (1999). Production d'ecdysteroïdes ovariens durant la maturation des oocytes chez *Tenebrio molitor*. *Annals of Entomological Society, France*, 35, 82-86.
- Sugumaran, M. (2010). Chapter 5 - Chemistry of cuticular sclerotization. *Advances in Insect Physiology*, 39, 151-209. <https://doi.org/10.1016/B978-0-12-381387-9.00005-1>
- Taha, A.A. (2014). Effect of some climatic factors on the seasonal activity of oriental wasp, *Vespa orientalis* attacking honeybee colonies in Dakahlia governorate, Egypt. *Egyptian Journal of Agricultural Research*. 92(1), 43-51.
- Taniai, K., Inceoglu, A.B. and Hammock, B.D. (2002). Expression efficiency of a scorpion neurotoxin, AaHIT, using baculovirus in insect cells. *Applied Entomology and Zoology*. 37(2), 225-232. DOI: 10.1303/aez.2002.225
- Taşkın, A.D. and Aksoylar, M.Y. (2011). *Itoptectis melanocephala* (Gravenhorst, 1829) (Hymenoptera: Ichneumonidae)'nın ergin öncesi dönemleri ile erginlerinin total lipid ve total yağ asidi yüzdeleri. *Turkish Entomology Journal*. 35(4), 641-649.
- Tedford, H.W., Sollod, B.L., Maggio, F. and King, G.F. (2004). Australian funnel-web spiders: master insecticide chemists. *Toxicon*. 43, 601-618. <https://doi.org/10.1016/j.toxicon.2004.02.010>
- Unsal, S., Ozparlak, H. and Aktumsek, A. (2004). Effects of diflubenzuron on the integument of fifth instar *Galleria mellonella* larvae. *Phytoparasitica*. 32 (1), 43-51. DOI:10.1007/BF02980858
- Vachon, M. (1966). Liste des scorpions connus en Égypte, Arabie, Israël, Liban, Syrie, Jordanie, Turquie, Irak, Iran. *Toxicon*. 4(3), 209-218. [https://doi.org/10.1016/0041-0101\(66\)90052-3](https://doi.org/10.1016/0041-0101(66)90052-3)
- Valk, T. and Meijden, A. (2014). Toxicity of scorpion venom in chick embryo and mealworm assay depending on the use of the soluble fraction versus the whole venom. *Toxicon*. 88, 38-43. doi: 10.1016/j.toxicon.2014.06.007
- Weichselbaum, T.E. (1946). Photometric colorimetric test for total proteins. *American Journal of Clinical Pathology*. 16, 40-48.
- Whetstone, P.A. and Hammock, B.D. (2007). Delivery methods for peptide and protein toxins in insect control. *Toxicon*. 49, 576-596. doi:10.1016/j.toxicon.2006.11.009
- Williams, S.S.C. (1968). Scorpion preservation for taxonomic and morphological studies. *The Wasmann Journal of Biology*. 26(1), 133-136.
- Windley, M.J., Herzig, V., Dziemborowicz, S.A., Hardy, M.C., King, G. F. and Nicholson, G.M. (2012). Spider-venom peptides as bioinsecticides. *Toxins (Basel)*, 4(3), 191-227. doi: 10.3390/toxins 4030191
- Xu, C., Zhang, Zh., Cui, K., Zhao, Y., Han, J., Liu, F. and Mu, W. (2016). Effects of sublethal concentrations of Cyantraniliprole on the development, fecundity and nutritional physiology of the black cutworm *Agrotis ipsilon* (Lepidoptera: Noctuidae). *PLOS ONE*, 19pp. <https://doi.org/10.1371/journal.pone.0156555>
- Zhou, G. and Miesfeld, R.L. (2009). Energy metabolism during diapause in *Culex pipiens* mosquitoes. *Journal of Insect Physiology*. 55, 40-46. doi: 10.1016/j.jinsphys.2008.10.002
- Zhu, Q., He, Y., Yao, J., Liu, Y.; Tao, L. and Huang, Q. (2012). Effects of sublethal concentrations of the chitin synthesis inhibitor, hexaflumuron, on the development and hemolymph physiology of the cutworm, *Spodoptera litura*. *J. Insect Sci.*, 12(27), 1-13. doi: 10.1673/031.012.2701

**Cite this article as:** Mohammad Tanani, Karem Ghoneim, Khalid Hamadah and Dyaa Emam (2021). Comparative impairing effects of selected arthropod venoms on the main body metabolites of *Galleria mellonella* (Lepidoptera: Pyralidae). *African Journal of Biological Sciences*. 3(3), 64-79. doi: 10.33472/AFJBS.3.3.2021.64-79.