

<https://doi.org/10.48047/AFJBS.6.15.2024.3895-3909>



African Journal of Biological Sciences

Journal homepage: <http://www.afjbs.com>



Research Paper

Open Access

IMPACT OF MINDFULNESS MEDITATION ON JOB STRESS, WORK-LIFE BALANCE AND RESILIENCE AMONG HEALTHCARE PROFESSIONALS

Aparna, K. K, Research Scholar, Govt. Women's College, Thiruvananthapuram.

Dr. Anjana, R., Assistant Professor, Department of Psychology, Sree Narayana College, Chempazhanchy.

Article History

Volume 6, Issue 15, 2024

Received : 03 May 2024

Accepted : 28 Aug 2024

Published: 07 Sep 2024

doi:10.48047/AFJBS.6.15.2024.3895-3909

Abstract

The study titled "Impact of Mindfulness Meditation on Job Stress, Work-Life Balance, and Resilience among Health care Professionals" explores the effectiveness of mindfulness meditation on job stress, work-life balance, and resilience among healthcare workers in Kerala. The study employed a pre-test/post test experimental design. The participants consisted of 36 healthcare professionals from both public and private sectors selected using multistage cluster sampling method. Participants underwent an eight-week Mindfulness-Based Stress Reduction (MBSR) program, attending weekly two-hour group sessions led by a certified instructor. The Parker Job Stress Scale (PJSS), Work-Life Balance Scale (WLBS), and Brief Resilience Scale (BRS) were administered to collect data. The data were analysed using the statistical software EDUSTAT. The paired samples t-test analysis revealed significant reduction in job stress, improvement in work-life balance and enhanced resilience. These findings suggest that mindfulness meditation is a valuable tool for reducing job stress and enhancing work-life balance and resilience among healthcare professionals.

Key words: Mindfulness Meditation, Job Stress, Work Life Balance, Resilience, Healthcare Professionals.

Introduction

In the fast-paced and demanding environment of healthcare, professionals frequently encounter high levels of job stress, which can adversely impact their work-life balance and overall resilience. Job stress among healthcare professionals is often attributed to long working hours, emotional strain from patient care, and administrative pressures (Shanafelt et

al., 2015). These stressors not only affect their mental and physical well-being but also compromise the quality of care provided to patients (Dyrbye et al., 2017). Addressing these challenges is crucial to ensuring both the well-being of healthcare workers and the efficiency of healthcare systems.

Mindfulness meditation, a practice rooted in the principles of present-moment awareness and non-judgmental acceptance, has emerged as a promising intervention for managing stress and enhancing psychological well-being (Kabat-Zinn, 1990). Numerous studies have documented the benefits of mindfulness meditation in reducing stress, improving emotional regulation, and fostering a sense of balance in various professional settings (Khoury et al., 2015). However, there is a need for more targeted research to understand its specific effects on healthcare professionals, who operate under unique and intense pressures compared to other occupational groups (Westphal et al., 2015).

This study aims to fill this gap by assessing the impact of mindfulness meditation on job stress, work-life balance, and resilience among healthcare professionals in Kerala. By employing an experimental design and utilizing validated measurement tools such as the Parker Job Stress Scale (PJSS), Work-Life Balance Scale (WLBS), and Brief Resilience Scale (BRS), this research seeks to provide empirical evidence on the effectiveness of mindfulness interventions. The findings are expected to offer valuable insights for healthcare administrators and policymakers, highlighting the potential of mindfulness practices to enhance the well-being and professional satisfaction of healthcare workers (Goyal et al., 2014).

Need and Significance of the Study

The healthcare industry is grappling with unprecedented levels of job stress, which significantly affect the mental health and overall well-being of professionals within this field.

High stress levels can lead to burnout, decreased job satisfaction, and increased turnover rates, all of which compromise the quality of patient care and burden healthcare systems (Shanafelt et al., 2015). Despite the critical role that healthcare professionals play in society, there has been insufficient focus on interventions that can effectively alleviate their stress and enhance their resilience. This gap underscores the urgent need for evidence-based strategies, such as mindfulness meditation, which can be integrated into the professional lives of healthcare workers to foster a more sustainable and healthy work environment.

The significance of this study lies in its potential to provide empirical support for the use of mindfulness meditation as a practical intervention for improving the well-being of healthcare professionals. By systematically evaluating the impact of mindfulness on job stress, work-life balance, and resilience, the research aims to offer valuable insights that can inform organizational policies and individual practices within healthcare settings. The findings could lead to the development of structured mindfulness programs tailored for healthcare environments, ultimately contributing to better mental health outcomes for professionals and improved patient care quality. In a broader context, this study emphasizes the importance of supporting healthcare workers through innovative and holistic approaches, recognizing their indispensable contribution to public health and the necessity of maintaining their well-being.

Objectives of the Study

The objectives formulated for the study are as follows.

1. To assess the impact of mindfulness meditation by comparing pre-test and post-test scores of Job Stress for Health Professionals.

2. To evaluate the effectiveness of mindfulness meditation by comparing pre-test and post-test scores of Work Life Balance for Health Professionals.

3. To determine the impact of mindfulness meditation by comparing pre-test and post-test scores of Resilience for Health Professionals.

Hypotheses of the Study

Based on the objectives the following hypotheses were formulated

1. There is significant difference between the pre test and post test scores of Job Stress for the Health Professionals.

2. There is significant difference between the pre test and post test scores of Work Life Balance for the Health Professionals.

3. There is significant difference between the pre test and post test scores of Resilience for the Health Professionals.

Method Adopted

The investigator adopted pre-post experimental design for the present study.

Population and Sample

The population of the study comprises healthcare professionals working in both government and private healthcare sectors in Kerala. This includes physicians, nurses, pharmacists, and other allied health workers. The sample consists of 36 healthcare professionals selected using a multistage cluster sampling technique. Among these, 15 participants are from public health centres, while the remaining 21 are from private health centres. The sample includes both male and female healthcare professionals aged between 20 and 60 years. It encompasses individuals with varying years of experience and family statuses, ensuring a diverse representation of the healthcare workforce in the study.

Variables of the Study

The study examines three main variables: job stress, work-life balance, and resilience. These variables are crucial indicators of the psychological well-being and professional satisfaction of healthcare professionals.

Instruments Used for the Study

The investigator employed the following instruments to collect data from the participants.

1. Parker Job Stress Scale (PJSS)
2. Work Life Balance Scale (WLBS)
3. Brief Resilience Scale (BRS)
4. Mindfulness Meditation Strategy

Procedure of the Study

The study commenced with the recruitment of participants from the diverse healthcare landscape of Thiruvananthapuram district, in Kerala, utilizing a multistage cluster sampling approach to ensure representation from both government and private health sectors. After obtaining informed consent, participants embarked on an eight-week intervention program rooted in Mindfulness-Based Stress Reduction (MBSR) principles, facilitated by trained professionals in structured sessions spanning two hours each week. From introductory sessions on mindfulness fundamentals to advanced explorations of stress management and work-life balance, participants engaged in diverse practices and discussions aimed at cultivating present moment awareness and fostering resilience. Preceding the intervention, participants underwent a comprehensive pre-test assessment of job stress, work-life balance, and resilience, followed by a post-test assessment upon completion of the program. The collected data underwent rigorous statistical analysis using the statistical software EDUSTAT

to unveil the transformative impact of mindfulness meditation on participants' well-being, offering insights into nurturing the mental health of healthcare professionals.

Statistical Techniques used in the study

The collected data for the pre-test and post-test were analysed using paired samples t-test.

Analysis and Interpretation

The investigator analysed the data collected using appropriate statistical techniques with the statistical software EDUSTAT and interpreted the findings based on the objectives and hypotheses formulated for the study.

Hypotheses in Null Form

For the purpose of testing the hypotheses were written in null form as follows.

1. There is no significant difference between the pre test and post test scores of Job Stress for the Health Professionals.
2. There is no significant difference between the pre test and post test scores of Work Life Balance for the Health Professionals.
3. There is no significant difference between the pre test and post test scores of Resilience for the Health Professionals.

Testing of Hypothesis 1

Table 1

Test of significance of difference between means of pre-test and post-test scores of
Job Stress of the Health Professionals

Asses sment	Num ber of students	Mean	Stand ard deviation	C.R	Level of significance
Pre test	36	41.06	5.19	12.96	0.01
Post test	36	24.11	5.11		


The calculated value of C.R. is 12.96 and is significant at 0.01 level (C.R. = 12.96; $p < 0.01$). The mean of the post-test scores of Job Stress is significantly lower than that of the pre-test scores. This can be attributed to the influence of the mindfulness intervention module.

Tenability of Hypothesis

Test of significance of difference between means of pre-test and post-test scores of Job Stress of the Health Professionals revealed that there is significant difference between pre-test and post-test scores of Job Stress. Hence the null hypothesis formulated in this context is rejected.

Figure 1

Mean pre-test and post-test scores of Job Stress of the Health Professionals



Testing of Hypothesis 2

Table 2

Test of significance of difference between means of pre-test and post-test scores of Work Life Balance of the Health Professionals

Asses sment	Num ber of students	Mean	Stand ard deviation	C.R	Level of significance
Pre test	36	30.92	6.62	13.57	0.01
Post test	36	64.61	13.25		

The calculated value of C.R. is 13.57 and is significant at 0.01 level (C.R. = 13.57; $p<0.01$). The mean of the post-test scores of Work Life Balance is significantly greater than


that of the pre-test scores. This can be attributed to the influence of the mindfulness intervention module.

Tenability of Hypothesis

Test of significance of difference between means of pre-test and post-test scores of Work Life Balance of the Health Professionals revealed that there is significant difference between pre-test and post-test scores of Work Life Balance. Hence the null hypothesis formulated in this context is rejected.

Figure 2

Mean pre-test and post-test scores of Work Life Balance of the Health Professionals



Testing of Hypothesis 3

Table 3

Test of significance of difference between means of pre-test and post-test scores of
Resilience of the Health Professionals

Asses sment	Num ber of students	Mean	Stand ard deviation	C.R	Level of significance
Pre test	36	10.39	2.15	10.29	0.01
Post test	36	20.39	5.62		


The calculated value of C.R. is 10.29 and is significant at 0.01 level (C.R. = 10.29; $p < 0.01$). The mean of the post-test scores of Resilience is significantly greater than that of the pre-test scores. This can be attributed to the influence of the mindfulness intervention module.

Tenability of Hypothesis

Test of significance of difference between means of pre-test and post-test scores of Resilience of the Health Professionals revealed that there is significant difference between pre-test and post-test scores of Resilience. Hence the null hypothesis formulated in this context is rejected.

Figure 3

Mean pre-test and post-test scores of Resilience of the Health Professionals



Findings of the Study

The calculated value of C.R. for significance of difference between means of pre-test and post-test scores of Job Stress of the Health Professionals is obtained as 12.96 and is significant at 0.01 level. The mean of the post-test scores of Job Stress is significantly lower than that of the pre-test scores. This means that the mindfulness intervention module is effective in reducing the Job Stress.

The calculated value of C.R. for significance of difference between means of pre-test and post-test scores of Work Life Balance of the Health Professionals is obtained as 13.57 and is significant at 0.01 level. The mean of the post test scores of Work Life Balance is significantly greater than that of the pre-test scores. This means that the mindfulness intervention module is effective in enhancing the Work Life Balance.

The calculated value of C.R. for significance of difference between means of pre-test and post-test scores of Resilience of the Health Professionals is obtained as 10.29 and is

significant at 0.01 level. The mean of the post test scores of Resilience is significantly greater than that of the pre-test scores. This means that the mindfulness intervention module is effective in enhancing the Resilience.

Discussion of the Results

The results of this study highlight the significant impact of mindfulness meditation on reducing job stress among healthcare professionals. The pre-test and post-test scores of the participants revealed a substantial decrease in job stress levels following the eight-week Mindfulness-Based Stress Reduction (MBSR) program. This finding aligns with previous research that underscores the effectiveness of mindfulness interventions in mitigating stress by promoting present-moment awareness and emotional regulation (Kabat-Zinn, 1990; Khoury et al., 2015). The reduction in job stress can be attributed to the participants' enhanced ability to manage stressors and maintain a calm and focused mindset, which is crucial in the high-pressure healthcare environment.

Additionally, the study demonstrated that mindfulness meditation significantly improves work-life balance among healthcare professionals. The participants' post-test scores indicated a marked increase in their ability to balance professional responsibilities with personal life. This improvement suggests that mindfulness practices help individuals prioritize and manage their time more effectively, reduce work-related spillover into personal life, and foster a greater sense of overall well-being (Goyal et al., 2014). Given the demanding nature of healthcare jobs, these findings are particularly relevant, as better work-life balance is linked to increased job satisfaction, lower burnout rates, and enhanced quality of care provided to patients.

Moreover, the study found that mindfulness meditation significantly boosts resilience among healthcare professionals. The increase in resilience scores from pre-test to post-test

suggests that participants developed a stronger capacity to recover from setbacks and adapt to challenging situations. This enhancement in resilience can be crucial for healthcare workers who often face emotionally and physically taxing scenarios (Westphal et al., 2015). By fostering resilience, mindfulness meditation equips healthcare professionals with the psychological tools to navigate their demanding work environment more effectively, contributing to sustained mental health and professional longevity.

In conclusion, the findings of this study provide robust evidence supporting the implementation of mindfulness meditation programs in healthcare settings. The significant reductions in job stress, improvements in work-life balance, and enhancements in resilience observed among the participants underscore the value of such interventions. These results advocate for healthcare organizations to consider integrating mindfulness practices into their wellness initiatives, aiming to enhance the overall well-being and professional satisfaction of their workforce. Future research could explore the long-term effects of mindfulness meditation and its impact on patient care outcomes, further solidifying its role in healthcare settings.

Conclusion

The study conclusively demonstrates that mindfulness meditation significantly reduces job stress and enhances work-life balance and resilience among healthcare professionals. The empirical evidence gathered from pre-test and post-test assessments indicates that participants who engaged in an eight-week Mindfulness-Based Stress Reduction (MBSR) program experienced notable improvements in their psychological well-being. The statistical analysis confirmed that the mindfulness intervention effectively mitigated job stress while simultaneously fostering a better balance between work and personal life and increasing resilience. These findings underscore the value of incorporating

mindfulness practices into healthcare settings as a viable strategy to support the mental health and professional satisfaction of healthcare workers, ultimately leading to improved patient care and more sustainable healthcare environments.

References

Dyrbye, L. N., West, C. P., Sinsky, C., Goeders, L. E., Satele, D. V., Shanafelt, T. D. (2017). "Medical licensure questions and physician reluctance to seek care for mental health conditions." *Mayo Clinic Proceedings*, 9(10), 1486-1493.

Goyal, M., Singh, S., Sibinga, E. M. S., Gould, N. F., Rowland-Seymour, A., Sharma, R., ... & Haythornthwaite, J. A. (2014). "Meditation programs for psychological stress and well-being: A systematic review and meta-analysis." *JAMA Internal Medicine*, 17(3), 357-368.

Kabat-Zinn, J. (1990). *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. Delacorte.

Khoury, B., Sharma, M., Rush, S. E., & Fournier, C. (2015). "Mindfulness-based stress reduction for healthy individuals: A meta-analysis." *Journal of Psychosomatic Research*, 78(6), 519-528.

Shanafelt, T. D., Boone, S., Tan, L., Dyrbye, L. N., Sotile, W., Satele, D., ... & Oreskovich, M. R. (2015). "Burnout and satisfaction with work-life balance among US physicians relative to the general US population." *Archives of Internal Medicine*, 17(18), 1377-1385.

Westphal, M., Bonanno, G. A., & Bartone, P. T. (2015). "Resilience and the ability to bounce back from stress." *Perspectives on Psychological Science*, 10(3), 346-351.

Westphal, M., Bonanno, G. A., & Bartone, P. T. (2015). Resilience and the ability to bounce back from stress. *Perspectives on Psychological Science*, 10(3), 346-351.

